

Original Research Article

Effect of Ripening Stage on Biochemical Composition of Seeds from three Species of *Canavalia* spp Consumed as Protein Substitutes in Côte d'Ivoire

ABSTRACT

Aims: The present study was aimed at investigating the nutritional properties of legume-seeds from *Canavalia* spp.

Methodology: Seeds of three species of *canavaliaseeds* notably *Canavaliagladiata*, *Canavaliaensiformis* and *Canavaliarosea*.

Results: The present study was aimed at investigating the nutritional properties of legume-seeds from *Canavalia* genus as a function of their ripening stage. Results reveal that the profiles of nutrients in the three *Canavalia* species varied as a function of ripening stages. The highest fat (3.27 ± 0.01 %), protein (33.20 ± 0.17), fibre (7.51 ± 0.06 %), ash (6.67 ± 0.01 %) and total sugar (25.40 ± 0.05 %) contents were obtained at the last ripening stage. Also, all the studied legume-seeds are excellent sources of proteins mainly consisting of high amounts of essential amino acids such as lysine (100.23 ± 0.04 mg/g prot), leucine (103.51 ± 0.04 mg/g prot), valine (83.68 ± 0.04 mg/g prot), methionine (27.36 ± 0.04 mg/g prot), tryptophan (74.32 ± 0.04 mg/g prot), cysteine (44.24 ± 0.04 mg/g prot), isoleucine (63.61 ± 0.04), threonine (40.22 ± 0.04 mg/g prot) and histidine (17.76 ± 0.04 mg/g prot). As regards mineral contents of these seeds, they were higher at penultimate ripening stage and, they partially meet dietary allowance of infants.

Conclusion: These seeds could be ranked as protein rich food for human and then, used as valuable substitutes for meat.

Keywords: Physicochemical composition, amino acid profil, legume-seeds, *Canavaliagladiata*, *Canavaliaensiformis*, *Canavaliarosea*

1. INTRODUCTION

Today, in many parts of the world, the exploitation of local resources is certainly a way to achieve the objective of food security [1], particularly for a fast-growing population and facing an increasing demand for protein and other nutrient-rich foods [2]. It seems imperative to improve and develop crops such as neglected or underutilized and called orphan or minor crops, by both national and international research programs [3, 4]. Many seeds generally used as vegetables or leguminous belong to this group of little-known plants. However, it is well known that legume seeds provide less expensive and important

Comment [TECNO L9 1]: repetition

protein sources to combat malnutrition, especially in developing countries where protein-rich foods of animal origin are not available for the people in low socio-economic groups. Otherwise, legumes are rich in many nutrients such as proteins, starch, dietary fibre, oils, vitamins, minerals and phytochemicals that are protected by nature. They provide substantial protein intake support to a significant proportion of the world's population, particularly in most developing countries [5]. Because of their high nutritional value, abundance of minerals and secondary metabolites, grain legumes have become valuable components of staple and functional foods [6]. The *Canavalia* species are categorized as a member of these crops. The genus of *Canavalia* is considered the third largest family among flowering plants [7]. It comprises approximately 50 species of tropical vines widely distributed in tropical and subtropical regions all over the world [8].

This genus was used traditionally as a food due to its nutritional significance. Sridhar and Seena [9] envisaged a comparative account of nutritional and functional properties of *Canavalia* species. *Canavaliagradiata* and *Canavaliaensiformis* are the common legume species having the potential to be a rich protein source, like edible legumes. Regarding *Canavalia* species notably, *Canavaliagradiata* (sword bean otherwise called "magic bean"), *Canavaliaensiformis* (jack bean) and *Canavaliarosea* (bay bean), they can be used to cushion the high cost of proteinaceous feed ingredients in the livestock industry [2]. It is pink coloured bean that originated from Asia and Africa. It is a leguminous, annual crop grown as a green manure, or cover crop and used as a fodder for livestock [10]. The high crude protein content (20 – 32%) and amino-acid profile of jack bean seed have been reported to make it suitable for use as a substitute for fish feed, while the fully ripened seeds are occasionally used as a coffee substitute [11]. Furthermore, the leguminous plant *Canavalia* species, though not eaten frequently, traditionally possess more medicinal properties yet to be scientifically proven. According to Soetan and Antia [12], jack bean and sword bean are rich in phytochemicals like flavonoids and saponins which should be exploited as medicinal foods for the benefits of human and animals.

Despite the aforementioned important nutritional and medicinal characteristics, in Côte d'Ivoire, *Canavalia* seeds are legume crops that have not been fully exploited. Thus, to our knowledge, no or very few studies have been devoted to biochemical and nutritional properties of these leguminous seeds.

2. MATERIAL AND METHODS

2.1. RAW MATERIAL COLLECTION AND SAMPLE PREPARATION

The three species of *canavaliaseeds* notably *Canavaliagradiata* (sword bean), *Canavaliaensiformis* (jack bean) and *Canavaliarosea* (bay bean) was obtained from an experimental plot located in Bouaké area's (7°41'00" N, 5°01'59" W) in Côte d'Ivoire. The seeds of these three species were collected at different ripening stages: 30 days (S1), 40 days (S2), 50 days (S3), 60 days (S4) and 80 days (S5) after fertilization.

After thoroughly drying in the sun, the pods were thrashed to remove seeds. The seeds, after thoroughly clearing and removal of broken seeds and foreign materials, were stored in airtight plastic containers at room temperature (25°C ± 2°C). The air-dried seeds (nearly 50 g from each accession) were powdered in a Wiley mill to pass a 60-mesh screen and stored in screw-capped bottles at room temperature for further analysis [13].

2.2. Proximate Composition Analysis

Dry matters were determined by drying in an oven at 105°C during 24 h to constant weight [13]. Crude protein was calculated from nitrogen (Nx6.25) obtained using the Kjeldahl method by AOAC [13]. Crude fat was determined by continuous extraction in a Soxhlet apparatus for 8 h using hexane as solvent [13]. Carbohydrate content was determined through the method

Comment [TECNO L9 2]: were

used by Samant and Rege[14]. Total ash was determined by incinerating in a furnace at 550°C [13]. Method described by Dubois et al.[15] was used to determine total sugars while reducing sugars were analysed according to the method of Bernfeld[16] using 3,5 dinitrosalicylic acids (DNS). The crude fibre contents were determined according to standard method [13]. The energy values of Canavalia seeds were evaluated using formula described by Crisan and Sands[17]. Energy value (kcal/100g) = (2.62 × % protein) + (8.37 × % fat) + (4.2 × % carbohydrate)

2.3. Minerals analysis

Minerals were determined employing AOAC[13]method. Flour was digested with a mixture of concentrated nitric acid (14.44 mol/L), sulfuric acid (18.01 mol/L) and perchloric acid (11.80 mol/L) and analysed using an atomic absorption spectrophotometer. The total phosphorus was determined as orthophosphate by the ascorbic acid method after acid digestion and neutralization using phenolphthalein indicator and combined reagent [18].

2.4. Amino acid composition

Total amino acid composition of samples was determined after hydrolysis in 6 M HCl with phenol (1%) at 150°C for 60 min, in Pico-Tag system (Waters, Milford, Mass., U.S.A.). The phenylisothiocyanate (PITC) amino acid derivatives were eluted on HPLC Applied Biosystems Model 172 A (Applera Corp, Foster City, Calif., U.S.A.) equipped with a PTC RP-18 column (2.1 mm × 22 cm). Sodium acetate (45 mM, pH 5.9) and sodium acetate (105 mM, pH4.6; 30%), and acetonitrile (70%) were used as buffers.

2.5. Statistical analysis

All analyses were performed in triplicates. Results are expressed as the mean ± standard deviation of several sample with Ky plot (version 2.0 beta 15, ©1997-2001, Koichi Yoshioka) statistical software. The data were statistically analysed by one way analysis of variance (ANOVA). Means were compared by Turkey's test. Differences were considered statistically significant at P < 0.05.

3. RESULTS AND DISCUSSION

3.1. Proximate Composition

The proximate composition of flours from the seeds of three Canavalia species was presented in **Table 1** for different ripening stages. Generally, the results reveal that fat, protein, fibre and ash contents increase as a function of ripening stages while the moisture and carbohydrate contents decrease in the same manner. Thus, at the last ripening stage (stage 5), highest fat, protein, fibre, ash and total sugar contents were obtained. As regards the moisture contents, all the studied seeds exhibited values about 5% at the last ripening stage suggesting that the studied beans are slightly perishable due to low moisture content which inhibits susceptibility to microbial growth and enzyme activity and, therefore could decelerates spoilage [19].

Canavalia ensiformis seeds were found to have the best protein, fibre, ash and total sugar contents with values of 33.20 ± 0.17, 7.51 ± 0.06, 6.67 ± 0.01 and 25.40 ± 0.05 %, respectively. At the same ripening stage, *Canavaliarosea* and *Canavaliagladia* exhibited protein contents of 32.05 ± 0.11 and 25.28 ± 0.29, respectively. These protein contents are higher than those reported for beans (ranged from 20.8 ± 1.3 to 23.5 ± 1.0%) [20], and for wild edible mushroom *Termitomyces heimii* (23.75 ± 0.04%) and *Amillariamellea* (21.32 ± 0.86 %) [21, 22]. Moreover, crude protein contents obtained in this study were greater than

Comment [TECNO L9 3]: increased

Comment [TECNO L9 4]: decreased

Comment [TECNO L9 5]: were

Comment [TECNO L9 6]: highest

Comment [TECNO L9 7]: which beans

Comment [TECNO L9 8]: compare with other grain legumes instead unrelated materials

those of some edible insects such as *Analeptestrifasciata* (20.1 %) and *Cirinafordalarva* (20 %) which are categorized as good protein sources [23, 24]. Therefore, *Canavalia* seeds could be ranked as protein rich food for human and then, used as valuable substitutes for meat or fish especially in developing countries for malnourished children suffering from kwashiorkor (a protein deficiency condition) and for pregnant women.

For crude fats, both *Canavaliarosea* and *Canavaliagladiata* showed the highest content of 3.25 ± 0.01 and $3.27 \pm 0.01\%$ (no significant difference) at the last ripening stage. These values are greater than those previously reported for some 52 common bean landraces (range of 0.6 to 2.9%) [25], and Yellow kidney beans had (2.7%) [20]. Los et al. [26] reported crude fat contents of 0.7 ± 0.01 to $2.7 \pm 0.09\%$ for common beans (*Phaseolusvulgaris* L.). However, these fat contents are low compared to high-fat legumes such as peanut and soybeans which have fat contents of about 25.3% and 22.19%, respectively [27, 28]. In fact, the studied seeds could be useful for preparation of some low-fat foodstuffs and snacks and in preventing hypertension and hypercholesterolemia [6]. Thus, they would be ideal foods for obese persons and recommended as good source of food supplement for patients with cardiac problems or at risk with lipid induced disorders as commented by Pedrosa et al. [29]. Concerning carbohydrates, the highest values (decreasing from 74.42 ± 0.08 to 58.17 ± 0.28 %) were obtained with *Canavaliagladiata* seeds and the lowest contents (decreasing from 66.94 ± 0.12 to 49.73 ± 0.22 %) with *Canavaliaensiformis* seeds. Butkute et al. [30] found carbohydrate contents ranged from 38.0 to 46.0 % for seven perennial legume seed species from Lithuania. So, values obtained in the present study were sited in the range of 58.8 to 77.4 % as reported by Los et al. [26] for several common bean varieties. These high carbohydrate contents would be due to their low lipid contents. Also, it might imply the uses of these seeds as binder, bulking agent or thickener in soups. These results suggest that the studied legume seeds would be highly nutritious and good for human and animal consumption.

It's noteworthy that the ash content of the sample is of nutritional importance as a previous report indicated that when leaves or vegetables are to be used as food for humans, they should contain about 3.0% ash [31]. As for the ash contents the studied *Canavalia* seeds, they were all above 3.0% whatever the ripening stage. At the last ripening stage, the ash contents were of 6.67 ± 0.01 , 6.05 ± 0.01 and 5.92 ± 0.01 , respectively for *Canavaliaensiformis*, *Canavaliarosea* and *Canavaliagladiata*. It indicates that these legume seeds are good sources of mineral elements with values greater than that previously reported (2.1%) by Omowunmi et al. [32] for *Canavaliaensiformis* seeds harvested in Ogun State (Nigeria).

The fibre contents is within $6.38 \pm 0.03 - 7.35 \pm 0.03\%$, $6.61 \pm 0.03 - 7.39 \pm 0.04\%$ and $6.79 \pm 0.04 - 7.51 \pm 0.06\%$, respectively for *Canavaliagladiate*, *Canavaliarosea* and *Canavaliaensiformis*. The ripened beans showed greater values of fibres. This ranges are similar to that found in the literature for legumes that serve as good sources of fibre [33].

The low energy levels provided by the studied *Canavalia* seeds (less than 400 kcal/100g DW) give only about 18 % of the daily energy intakes recommended for a 70 kg person. It is well known that the energy content is affected by the proportion of fat, protein and carbohydrate in the vegetables. The present results suggested that *Canavalia* seeds can be used as a source of low energy diet [21].

Comment [TECNO L9 9]: remove 'had'

Comment [TECNO L9 10]: verify

Comment [TECNO L9 11]: give reasons why the decrease with ripening

Comment [TECNO L9 12]: adduce reason why... : decrease in moisture contents lead to increase in dry matter

Comment [TECNO L9 13]: These

Comment [TECNO L9 14]: known

Table 1: Proximate composition of.....

Species	Stage	P a r a m e t e r s									
		Dry matter (%)	Moisture (%)	Fats (%)	Proteins (%)	Carbohydrates (%)	Fibres (%)	Ash (%)	Total sugars (%)	Reducing sugars (%)	Energy value (Kcal)
Canavaliagladiata (CG)	S 1	87.24 ± 0.01 ^c	12.76 ± 0.01 ^p	2.05 ± 0.01 ^e	13.67 ± 0.12 ^a	74.42 ± 0.08 ^q	6.38 ± 0.03 ^a	3.47 ± 0.01 ^c	8.14 ± 0.00 ^b	0.98 ± 0.04 ^a	370.83 ± 0.11 ^p
	S 2	90.54 ± 0.01 ^f	9.46 ± 0.01 ^k	2.19 ± 0.01 ^f	15.67 ± 0.12 ^b	71.52 ± 0.11 ^p	6.68 ± 0.01 ^c	3.93 ± 0.01 ^e	14.84 ± 0.00 ^d	2.01 ± 0.04 ^h	368.51 ± 0.05 ^m
	S 3	92.56 ± 0.02 ^h	7.44 ± 0.02 ^h	2.43 ± 0.01 ^h	20.11 ± 0.12 ^d	66.26 ± 0.12 ^k	6.77 ± 0.02 ^d	4.43 ± 0.01 ^g	17.98 ± 0.18 ^e	3.58 ± 0.03 ^k	367.38 ± 0.07 ^j
	S 4	94.27 ± 0.00 ^p	5.73 ± 0.00 ^c	2.79 ± 0.01 ^k	24.70 ± 0.14 ^h	60.12 ± 0.11 ^g	6.86 ± 0.03 ^e	5.51 ± 0.01 ^k	19.77 ± 0.29 ^{ef}	4.08 ± 0.03 ^m	364.46 ± 0.12 ^f
	S 5	95.00 ± 0.00 ^r	5.00 ± 0.00 ^a	3.27 ± 0.01 ⁿ	25.28 ± 0.29 ^j	58.17 ± 0.28 ^f	7.35 ± 0.03 ^k	5.92 ± 0.01 ⁿ	20.97 ± 0.10 ^f	1.65 ± 0.03 ^e	363.27 ± 0.13 ^d
Canavaliarosea (CR)	S 1	86.43 ± 0.01 ^b	13.57 ± 0.01 ^q	1.85 ± 0.02 ^c	19.09 ± 0.18 ^c	69.30 ± 0.20 ⁿ	6.61 ± 0.03 ^b	3.15 ± 0.01 ^a	5.41 ± 0.00 ^a	1.52 ± 0.00 ^d	370.17 ± 0.09 ⁿ
	S 2	88.90 ± 0.01 ^d	11.10 ± 0.01 ⁿ	2.01 ± 0.02 ^d	22.09 ± 0.18 ^f	65.22 ± 0.23 ^j	7.01 ± 0.05 ^f	3.67 ± 0.01 ^d	7.71 ± 0.01 ^b	1.67 ± 0.00 ^e	367.32 ± 0.21 ^j
	S 3	91.77 ± 0.00 ^g	8.23 ± 0.00 ^j	2.47 ± 0.01 ^j	27.53 ± 0.18 ^k	58.26 ± 0.17 ^f	7.16 ± 0.04 ^h	4.57 ± 0.01 ^h	11.22 ± 0.08 ^c	2.01 ± 0.03 ^h	365.42 ± 0.15 ^h
	S 4	92.58 ± 0.00 ^j	7.42 ± 0.00 ^g	2.89 ± 0.01 ^m	30.88 ± 0.16 ⁿ	53.87 ± 0.17 ^d	7.25 ± 0.07 ^j	5.11 ± 0.01 ^j	15.14 ± 0.16 ^d	2.94 ± 0.04 ^j	365.03 ± 0.25 ^g
	S 5	94.23 ± 0.01 ⁿ	5.77 ± 0.01 ^d	3.25 ± 0.01 ⁿ	32.05 ± 0.11 ^q	51.25 ± 0.10 ^b	7.39 ± 0.04 ^k	6.05 ± 0.01 ^p	19.07 ± 0.25 ^e	1.34 ± 0.02 ^c	362.49 ± 0.15 ^c
Canavaliainsiformis (CE)	S 1	85.79 ± 0.01 ^a	14.21 ± 0.01 ^r	1.67 ± 0.02 ^a	21.28 ± 0.07 ^e	66.94 ± 0.12 ^m	6.79 ± 0.04 ^d	3.31 ± 0.01 ^b	5.54 ± 4.04 ^a	1.31 ± 0.02 ^c	367.93 ± 0.14 ^k
	S 2	89.34 ± 0.00 ^e	10.66 ± 0.00 ^m	1.71 ± 0.01 ^b	24.28 ± 0.07 ^g	62.75 ± 0.11 ^h	7.08 ± 0.03 ^g	4.17 ± 0.01 ^f	10.30 ± 0.01 ^c	1.51 ± 0.00 ^d	363.55 ± 0.11 ^e
	S 3	92.81 ± 0.00 ^k	7.19 ± 0.00 ^f	2.03 ± 0.01 ^e	29.72 ± 0.07 ^m	55.37 ± 0.10 ^e	7.19 ± 0.02 ^{hj}	5.69 ± 0.01 ^m	16.12 ± 0.18 ^d	1.76 ± 0.03 ^f	358.64 ± 0.08 ^b
	S 4	93.81 ± 0.01 ^m	6.19 ± 0.01 ^e	2.29 ± 0.01 ^g	31.54 ± 0.12 ^p	52.70 ± 0.13 ^c	7.35 ± 0.03 ^k	6.11 ± 0.01 ^q	21.40 ± 0.24 ^f	1.89 ± 0.02 ^g	357.63 ± 0.11 ^a
	S 5	94.56 ± 0.00 ^q	5.44 ± 0.00 ^b	2.89 ± 0.01 ^m	33.20 ± 0.17 ^r	49.73 ± 0.22 ^a	7.51 ± 0.06 ^m	6.67 ± 0.01 ^f	25.40 ± 0.05 ^g	1.10 ± 0.04 ^b	357.74 ± 0.24 ^a

Values are mean ± standard deviation of three measurements (n = 3). The different case letters (a, b, c) in the column indicate significant differences ($p < 0.05$) in the respective values. S1: Stage 1; S2: Stage 2; S3: Stage 3; S4: Stage 4; S5: Stage 5.

3.2. Mineral

The variation of mineral contents as a function of ripening stages is given in **Table 2**. Significant differences ($p < 0.05$) were noted in mineral composition as a function of *Canavalia* species and, ripening stages. Results reveal that highest mineral contents were obtained at the penultimate ripening stage (stage 4) for all the studied legume-seeds. *C. rosea* seeds contained the **highest** concentration of potassium ($43506.67 \pm 60.28 \mu\text{g } 100 \text{ g}^{-1}$), magnesium ($27656.67 \pm 30.55 \mu\text{g } 100 \text{ g}^{-1}$), manganese ($9.65 \pm 0.39 \mu\text{g } 100 \text{ g}^{-1}$), zinc ($107.45 \pm 0.22 \mu\text{g } 100 \text{ g}^{-1}$) and iron ($973.67 \pm 4.16 \mu\text{g } 100 \text{ g}^{-1}$), while calcium ($9669.37 \pm 0.54 \mu\text{g } 100 \text{ g}^{-1}$) and copper ($74.67 \pm 4.04 \mu\text{g } 100 \text{ g}^{-1}$) contents were greater in *C. gladiata* seeds. These values are higher than those previously reported by Nkwocha et al.[34] for the same legume-seeds harvested in the Kaduna state (Nigeria).

Potassium was the most abundant macro-mineral and its composition ranged from 999.85 ± 30.65 to $35400.00 \pm 45.83 \mu\text{g } 100 \text{ g}^{-1}$ in *C. gladiata*, 564.85 ± 30.65 to $43506.67 \pm 60.28 \mu\text{g } 100 \text{ g}^{-1}$ in *C. rosea* and 452.21 ± 6.98 to $13610.00 \pm 45.83 \mu\text{g } 100 \text{ g}^{-1}$ in *C. ensiformis* seeds. It partially meets requirements for infants (500-700 mg/ 100 g DW) than adults and pregnant lactating women (2000 mg/ 100 g DW) [35]. In the same manner, magnesium, manganese, zinc, iron, calcium and copper contents in the studied seeds partially meet dietary allowance of infants. It would be recommended to consume these legume-seeds with the addition of mineral-rich foods such as meat, fish or even caterpillars to meet the daily nutrient needs as describe by Akpoussan et al[36] and Assiélou et al.[37].

Comment [TECNO L9 16]: which?

UNDER PEER REVIEW

Comment [TECNO L9 17]:of three species of canva..... at different stages of ripening

Table 2: Mineral composition.....

Species	Stage	P a r a m e t e r s								
		Calcium (ug/100 g DW)	Phosphorus (ug/100 g DW)	Magnesium (ug/100 g DW)	Potassium (ug/100 g DW)	Sodium (ug/100 g DW)	Manganese (ug/100 g DW)	Zinc (ug/100 g DW)	Copper (ug/100 g DW)	Iron (ug/100 g DW)
Canavaliaglabrata (CG)	S 1	1239.37 ± 0.64 ^d	967.33 ± 41.88 ^b	234.33 ± 3.51 ^a	999.85 ± 30.65 ^c	0.97 ± 0.07 ^b	0.00 ± 0.00 ^a	9.89 ± 0.50 ^b	1.89 ± 0.12 ^a	18.37 ± 0.15 ^{ab}
	S 2	2033.37 ± 0.64 ^f	1878.33 ± 41.88 ^d	4340.33 ± 29.87 ^h	10195.95 ± 30.65 ^f	9.01 ± 0.13 ^d	0.50 ± 0.02 ^b	14.24 ± 0.11 ^c	8.98 ± 0.04 ^{cd}	23.33 ± 0.08 ^{bc}
	S 3	7588.47 ± 76.61 ^p	3483.33 ± 35.12 ^e	9523.33 ± 29.87 ^k	18161.00 ± 53.51 ^m	12.37 ± 0.35 ^f	1.55 ± 0.05 ^d	42.67 ± 1.53 ^e	27.50 ± 0.54 ^g	109.80 ± 0.19 ^d
	S 4	9669.37 ± 0.64 ^r	9536.67 ± 28.87 ⁿ	21297.33 ± 12.22 ^q	35400.00 ± 45.83 ^p	47.60 ± 0.36 ^k	4.11 ± 0.10 ^f	99.95 ± 1.00 ^h	74.67 ± 4.04 ⁿ	465.91 ± 3.91 ^j
	S 5	4746.47 ± 81.00 ^k	4200.00 ± 55.68 ^f	12390.00 ± 29.31 ⁿ	19500.00 ± 55.68 ⁿ	32.1 ± 0.30 ^h	3.10 ± 0.11 ^e	62.33 ± 1.15 ^f	39.80 ± 0.74 ^h	186.88 ± 2.90 ^f
Canavaliarosea (CR)	S 1	699.39 ± 0.62 ^a	627.67 ± 15.95 ^a	1007.31 ± 11.09 ^c	564.85 ± 30.65 ^b	0.97 ± 0.05 ^b	0.00 ± 0.00 ^a	9.83 ± 0.61 ^b	1.45 ± 0.15 ^a	18.32 ± 0.07 ^b
	S 2	1039.39 ± 0.62 ^c	1439.33 ± 41.88 ^c	4190.00 ± 36.06 ^g	2287.55 ± 30.65 ^d	6.64 ± 0.32 ^c	0.99 ± 0.04 ^c	27.23 ± 1.62 ^d	5.01 ± 0.17 ^b	23.94 ± 0.34 ^c
	S 3	3020.00 ± 70.49 ^g	5153.33 ± 45.09 ^g	11190.00 ± 36.06 ^m	14723.33 ± 49.33 ^k	14.43 ± 0.49 ^g	2.96 ± 0.25 ^e	40.67 ± 1.53 ^e	13.88 ± 0.19 ^e	204.00 ± 4.36 ^g
	S 4	6201.47 ± 3.07 ^m	9900.00 ± 45.83 ^p	27656.67 ± 30.55 ^r	43506.67 ± 60.28 ^r	84.43 ± 0.31 ⁿ	9.65 ± 0.39 ^m	107.45 ± 0.22 ^j	65.95 ± 0.04 ^m	973.67 ± 4.16 ⁿ
	S 5	3433.13 ± 57.91 ⁱ	7556.67 ± 20.82 ^k	13173.33 ± 56.86 ^p	35956.67 ± 61.10 ^q	41.08 ± 0.16 ^j	7.78 ± 0.16 ^j	76.98 ± 1.75 ^g	46.88 ± 0.12 ^j	741.67 ± 3.06 ^m
Canavaliensisformis (CE)	S 1	879.39 ± 0.62 ^b	907.33 ± 41.88 ^b	294.67 ± 3.51 ^b	452.21 ± 6.98 ^a	0.00 ± 0.00 ^a	0.25 ± 0.02 ^d	5.53 ± 0.11 ^a	7.70 ± 0.28 ^c	13.25 ± 0.07 ^a
	S 2	1949.39 ± 0.62 ^e	3546.33 ± 41.88 ^e	2035.23 ± 42.39 ^d	3891.21 ± 14.32 ^e	0.00 ± 0.00 ^a	0.27 ± 0.04 ^d	9.89 ± 0.61 ^b	10.31 ± 0.28 ^d	21.95 ± 0.09 ^{bc}
	S 3	3292.47 ± 12.53 ^h	6890.00 ± 45.83 ^h	3238.00 ± 20.30 ^f	10923.33 ± 35.12 ^g	10.13 ± 0.55 ^e	5.07 ± 0.11 ^g	13.13 ± 0.35 ^c	23.40 ± 0.35 ^f	317.33 ± 3.06 ^h
	S 4	8606.47 ± 11.72 ^q	7043.33 ± 45.09 ^j	5730.00 ± 26.46 ^j	13610.00 ± 45.83 ^j	93.77 ± 0.60 ^p	8.59 ± 0.09 ^k	78.00 ± 4.58 ^g	63.90 ± 0.13 ^k	597.33 ± 6.43 ^k
	S 5	7466.47 ± 57.56 ⁿ	8880.00 ± 20.00 ^m	2453.33 ± 15.28 ^e	11996.67 ± 65.06 ^h	70.87 ± 0.75 ^m	7.03 ± 0.15 ^h	40.99 ± 1.02 ^e	46.51 ± 0.50 ^j	164.17 ± 3.82 ^e

Values are mean ± standard deviation of three measurements (n = 3). The different case letters (a, b, c) in the column indicate significant differences (p < 0.05) in the respective mineral concentrations. S1: Stage 1; S2: Stage 2; S3: Stage 3; S4: Stage 4; S5: Stage 5.

3.3. Amino acids profiles of three species of canavalia seed proteins at different stages of ripening.

Amino acid profiles are indicators of proteins nutritional qualities and functionalities. Essential/nonessential amino acid content is one of the parameters that provide important nutritional details about the protein quality of legumes [38]. The amino acid profiles of the three *Canavalia* species as a function of ripening stages is shown in **Table 3**. Overall observation revealed significant differences ($p < 0.05$) in amino acid amounts as a function of *Canavalia* species and, ripening stages. Generally the highest values were obtained at penultimate or last ripening stage.

The essential amino acids detected were methionine, valine, tryptophan, lysine, leucine, tyrosine, cysteine, isoleucine, threonine and histidine. Among them, the highest amounts were obtained with lysine, leucine and valine. Thus, *C. ensiformis* seeds contained 103.51 ± 0.04 mg/ g protein of leucine and 100.23 ± 0.04 mg/ g protein of lysine, while *C. rosea* seeds exhibited 83.68 ± 0.04 mg/ g protein of valine. The amino acid amounts determined in this study were higher than those reported for *C. ensiformis* in Nigeria [39], three jack bean accessions [33] and for pigeon pea [40]. The relatively high content of essential amino acids in the studied *Canavalia* seeds make them potential supplement in cereal-based diets due to the fact that cereal grain-based diets have been reported to be deficient in some essential amino acid such as lysine [41]. Furthermore, as argued by Millar et al.[42], the amino acid content of the three studied legume-seeds meet dietary allowance for adults [43].

Comment [TECNO L9 18]: table 3 is bulky and needs to be summarized: sulfure, aromatic, essential & nonessential, etc

Comment [TECNO L9 19]: acid

Comment [TECNO L9 20]: complete

Comment [TECNO L9 21]: remove

Comment [TECNO L9 22]: In all your discussions, you never aduced reason or reasons on what were responsible for higher contents of nutrients either at penultimate or terminal end

Comment [TECNO L9 23]: why?

Comment [TECNO L9 24]: what are limiting amino acids and most abundant, compare with soybean proteins

Comment [TECNO L9 25]: reason

Comment [TECNO L9 26]: compare with WHO/FAO standards

UNDER PEER REVIEW

Comment [TECNO L9 27]: as in other tables, give complete title of table content

Table 3: Amino acid composition.....

S p e c i e s	Stage	Arginine (mg/g proteins)	Methionine (mg/g prot)	Valine (mg/g prot)	Alaline (mg/g prot)	Tryptophane (mg/g prot)	Proline (mg/g prot)	Lysine (mg/g prot)	Glutamic acid (mg/g prot)
Canavaliagladiaata (CG)	S 1	20.36 ± 0.04 ^a	9.27 ± 0.04 ^e	0.00 ± 0.00 ^a	6.32 ± 0.04 ^a	0,00 ± 0,00 ^a	30.57 ± 0.04 ^e	48.33 ± 0.04 ^a	55.62 ± 0.04 ^h
	S 2	27.97 ± 0.04 ^b	17.39 ± 0.04 ^h	0.00 ± 0.00 ^a	12.87 ± 0.04 ^b	3.27 ± 0.04 ^b	39.31 ± 0.04 ^g	61.51 ± 0.04 ^c	77.11 ± 0.04 ⁿ
	S 3	33.76 ± 0.04 ^c	24.32 ± 0.04 ^p	6.88 ± 0.04 ^b	18.93 ± 0.04 ^c	9.35 ± 0.04 ^c	50.92 ± 0.04 ^p	72.68 ± 0.04 ^g	87.29 ± 0.04 ^p
	S 4	42.18 ± 0.04 ^f	23.76 ± 0.04 ⁿ	10.44 ± 0.04 ^c	23.77 ± 0.04 ^e	14.16 ± 0.04 ^e	62.15 ± 0.04 ^r	84.53 ± 0.04 ⁿ	98.35 ± 0.04 ^q
	S 5	40.74 ± 0.04 ^e	27.36 ± 0.04 ^q	12.58 ± 0.04 ^d	22.34 ± 0.04 ^d	13.48 ± 0.04 ^d	59.37 ± 0.04 ^g	80.17 ± 0.04 ^j	110.54 ± 0.04 ^r
Canavaliarosea (CR)	S 1	51.35 ± 0.04 ^g	2.58 ± 0.04 ^b	52.72 ± 0.04 ^f	68.53 ± 0.04 ^m	22.67 ± 0.04 ^f	20.16 ± 0.04 ^b	51.65 ± 0.04 ^b	32.15 ± 0.04 ^d
	S 2	59.62 ± 0.04 ^h	7.94 ± 0.04 ^d	60.48 ± 0.04 ^h	81.58 ± 0.04 ⁿ	30.22 ± 0.04 ^g	28.55 ± 0.04 ^c	69.11 ± 0.04 ^e	44.71 ± 0.04 ^g
	S 3	70.17 ± 0.04 ^k	11.84 ± 0.04 ^g	72.61 ± 0.04 ^k	100.22 ± 0.04 ^r	37.59 ± 0.04 ⁱ	34.41 ± 0.04 ^f	82.84 ± 0.04 ^m	57.36 ± 0.04 ^j
	S 4	85.59 ± 0.04 ⁿ	17.33 ± 0.04 ^h	80.02 ± 0.04 ^m	98.78 ± 0.04 ^q	44.36 ± 0.04 ^k	40.72 ± 0.04 ^h	80.77 ± 0.04 ^k	70.92 ± 0.04 ^k
	S 5	92.34 ± 0.04 ^r	22.47 ± 0.04 ^m	83.68 ± 0.04 ^q	97.22 ± 0.04 ^p	50.34 ± 0.04 ⁿ	45.92 ± 0.04 ^k	79.39 ± 0.04 ^h	73.21 ± 0.04 ^m
Canavaliensisiformis (CE)	S 1	39.51 ± 0.04 ^d	0.00 ± 0.00 ^a	37.66 ± 0.04 ^e	25.28 ± 0.04 ^f	32.12 ± 0.04 ^h	17.63 ± 0.04 ^a	68.15 ± 0.04 ^d	13.67 ± 0.04 ^a
	S 2	60.57 ± 0.04 ^j	6.11 ± 0.04 ^c	52.91 ± 0.04 ^g	41.51 ± 0.04 ^g	49.23 ± 0.04 ^m	28.68 ± 0.04 ^d	71.23 ± 0.04 ^f	17.15 ± 0.04 ^b
	S 3	73.14 ± 0.04 ^m	10.02 ± 0.04 ^f	70.17 ± 0.04 ⁱ	50.49 ± 0.04 ^h	61.25 ± 0.04 ^p	41.15 ± 0.04 ^j	88.13 ± 0.04 ^p	29.98 ± 0.04 ^c
	S 4	87.21 ± 0.04 ^p	18.77 ± 0.04 ^j	81.40 ± 0.04 ^p	60.77 ± 0.04 ^j	74.32 ± 0.04 ^r	48.16 ± 0.04 ⁿ	100.23 ± 0.04 ^r	38.47 ± 0.04 ^e
	S 5	88.70 ± 0.04 ^q	20.91 ± 0.04 ^k	80.63 ± 0.04 ⁿ	62.35 ± 0.04 ^k	72.77 ± 0.04 ^q	46.38 ± 0.04 ^m	97.68 ± 0.04 ^q	42.87 ± 0.04 ^f

Values are mean ± standard deviation of three measurements (n = 3). The different case letters (a, b, c) in the column indicate significant differences ($p < 0.05$) in the respective mineral concentrations. S1: Stage 1; S2: Stage 2; S3: Stage 3; S4: Stage 4; S5: Stage 5.

S p e c i e s	Stage	Leucine (mg/g prot)	Serine (mg/g prot)	Tyrosine (mg/g prot)	Cysteine (mg/g prot)	Isoleucine (mg/g prot)	Aspartic acid (mg/g prot)	Threonine (mg/g prot)	Histidine (mg/g prot)	Glycine (mg/g prot)
Canavaliaglabrata (CG)	S 1	38.29 ± 0.04 ^a	16.71 ± 0.04 ^a	17.54 ± 0.04 ^j	5.73 ± 0.04 ^a	22.41 ± 0.04 ^b	33.53 ± 0.04 ^b	14.39 ± 0.04 ^e	0.00 ± 0.00 ^a	4.32 ± 0.04 ^a
	S 2	57.27 ± 0.04 ^c	24.77 ± 0.04 ^d	26.29 ± 0.04 ^m	15.11 ± 0.04 ^d	37.71 ± 0.04 ^g	69.37 ± 0.04 ^h	25.66 ± 0.04 ^k	1.11 ± 0.04 ^b	10.63 ± 0.04 ^b
	S 3	64.15 ± 0.04 ^f	32.22 ± 0.04 ^g	36.13 ± 0.04 ⁿ	17.17 ± 0.04 ^e	49.23 ± 0.04 ^k	80.55 ± 0.04 ^k	32.47 ± 0.04 ^m	4.43 ± 0.04 ^e	18.15 ± 0.04 ^c
	S 4	71.44 ± 0.04 ^h	40.63 ± 0.04 ^j	42.47 ± 0.04 ^p	20.73 ± 0.04 ⁱ	58.63 ± 0.04 ^p	91.84 ± 0.04 ^m	38.11 ± 0.04 ^q	11.24 ± 0.04 ^h	23.41 ± 0.04 ^f
	S 5	69.21 ± 0.04 ^g	42.53 ± 0.04 ^k	44.76 ± 0.04 ^q	18.40 ± 0.04 ^f	63.61 ± 0.04 ^q	97.35 ± 0.04 ⁿ	36.71 ± 0.04 ⁿ	15.76 ± 0.04 ^j	20.33 ± 0.04 ^d
Canavaliarosea (CR)	S 1	50.43 ± 0.04 ^b	19.21 ± 0.04 ^b	0.00 ± 0.00 ^a	13.15 ± 0.04 ^c	22.63 ± 0.04 ^c	42.33 ± 0.04 ^d	3.49 ± 0.04 ^a	0.00 ± 0.00 ^a	41.39 ± 0.04 ^h
	S 2	58.11 ± 0.04 ^d	28.19 ± 0.04 ^e	2.53 ± 0.04 ^c	20.44 ± 0.04 ^h	30.29 ± 0.04 ^e	50.15 ± 0.04 ^e	10.02 ± 0.04 ^c	0.00 ± 0.00 ^a	57.18 ± 0.04 ⁿ
	S 3	71.72 ± 0.04 ^j	40.55 ± 0.04 ^h	9.33 ± 0.04 ^e	31.50 ± 0.04 ^m	44.88 ± 0.04 ^h	54.76 ± 0.04 ^f	17.28 ± 0.04 ^f	1.81 ± 0.04 ^c	64.35 ± 0.04 ^p
	S 4	86.37 ± 0.04 ^m	51.27 ± 0.04 ⁿ	15.41 ± 0.04 ^h	40.78 ± 0.04 ^p	54.11 ± 0.04 ^m	70.44 ± 0.04 ^j	23.62 ± 0.04 ^g	7.77 ± 0.04 ^f	72.71 ± 0.04 ^q
	S 5	88.22 ± 0.04 ⁿ	58.27 ± 0.04 ^r	15.22 ± 0.04 ^g	44.24 ± 0.04 ^r	58.39 ± 0.04 ⁿ	68.61 ± 0.04 ^g	24.44 ± 0.04 ^h	11.22 ± 0.04 ^h	75.29 ± 0.04 ^r
Canavaliensisformis (CE)	S 1	60.03 ± 0.04 ^e	20.51 ± 0.04 ^c	0.00 ± 0.00 ^a	11.17 ± 0.04 ^b	10.71 ± 0.04 ^a	0.00 ± 0.00 ^a	9.21 ± 0.04 ^b	0.00 ± 0.00 ^a	21.38 ± 0.04 ^e
	S 2	72.11 ± 0.04 ^k	29.24 ± 0.04 ^f	1.02 ± 0.04 ^b	18.75 ± 0.04 ^g	23.74 ± 0.04 ^d	0.00 ± 0.00 ^a	13.88 ± 0.04 ^d	4.29 ± 0.04 ^b	30.11 ± 0.04 ^g
	S 3	89.78 ± 0.04 ^p	42.72 ± 0.04 ^m	6.77 ± 0.04 ^d	30.69 ± 0.04 ^k	31.43 ± 0.04 ^f	0.00 ± 0.00 ^a	24.58 ± 0.04 ^j	10.13 ± 0.04 ^g	42.39 ± 0.04 ^j
	S 4	100.02 ± 0.04 ^q	55.67 ± 0.04 ^q	13.18 ± 0.04 ^f	39.26 ± 0.04 ⁿ	48.60 ± 0.04 ^j	40.55 ± 0.04 ^c	37.19 ± 0.04 ^p	17.76 ± 0.04 ^m	52.26 ± 0.04 ^m
	S 5	103.51 ± 0.04 ^r	54.47 ± 0.04 ^p	21.13 ± 0.04 ^k	43.76 ± 0.04 ^q	30.29 ± 0.04 ^e	0.00 ± 0.00 ^a	40.22 ± 0.04 ^r	16.27 ± 0.04 ^k	51.55 ± 0.04 ^k

Table4: Amino acid composition in different species

4. CONCLUSION

The present study reveals that the profiles of nutrients in the three *Canavalia* species varied as a function of ripening stages. The highest fat, protein, fibre, ash and total sugar contents were obtained at the last ripening stage especially in *C. ensiformis* seeds. So, all the studied legume-seeds are excellent sources of proteins mainly consisting of high amounts of essential amino acids such as lysine, leucine, valine, methionine, tryptophan, cysteine, isoleucine, threonine and histidine. As regards mineral contents of these seeds, they were higher at penultimate ripening stage and, they partially meet dietary allowance of infants.

REFERENCES

1. KoffiDM, FauletBM, GonnetyJT, BédikouME, KouaméLP, ZoroBi IA, NiamkéSL. Biochemical characterization of two acid phosphatases purified from edible seeds of the neglected crop *Lagenariasiceraria*(Molina) Standl. Blocky-fruited cultivar. Philipp. Agric. Scientist. 2010; 93(3): 269–280.
2. Okla MK,Akhtar N,Alamri SA, Al-Qahtani SM, Ismail A,Abbas ZK, AL-Ghamdi AA,Qahtan AA, Soufan WH, Alaraidh IA. Potential Importance of Molybdenum Priming to Metabolism and Nutritive Value of *Canavali* spp. Sprouts. Plants.2021;10: 2387.<https://doi.org/10.3390/plants.10112387>.
3. RasulMG, HiramatsuM, Okubo H. Genetic relatedness (diversity) and cultivar identification by randomly amplified polymorphic DNA (RAPD) markers in teasle gourd (*Momordicadioica*Roxb). Sci. Hort. 2007; 111: 271-279.
4. EsfeldK, Plaza S, TadeleZ. Bringing highthroughput techniques to orphan crop of Africa: Highlights from the Tef TILLING Project. Tef Biotechnology project, Institute of Plant Sciences, University of Bern, Switzerland. 2009; 5 p.
5. Ojo MA, Ade-Omowaye BIO. Some Functional and Physical Properties of Selected Underutilised Hard-To-Cook Legumes in Nigeria. Amer. J. Food Sci.Nutr.2015; 2(5): 73-81.
6. Bouchenak M, Lamri-Senhadj M. Nutritional quality of legumes, and their role in cardiometabolic risk prevention: A review. J. Med. Food. 2013;16: 185–198.
7. Michael K,Sogbesan O,Onyia L. Effect of processing methods on the nutritional value of *Canavaliaensiformis*jack bean seed meal. J. Food Process. Technol. 2018;9:766.
8. YamashiroA, Yamashiro T. Utilization on extrafloralnectaries and fruit domatia of *Canavali**lineata*and *C. cathartica*(Leguminosae) by ants. Arthropod-Plant Interact. 2008;2: 1–8.
9. Sridhar KR, Seen S. Nutritional and antinutritional significance of four unconventional legumes of the genus *Canavalia*—A comparative study. Food Chem. 2006;99: 267–288.
10. Kerala K.Minor vegetables. Kirsankerala Operations centre, IIITM-K, NLLA Techno park Campus, ThiruVananthapuram. 2018;4(23): 20-24.
11. Osuigwe DI, Obiekezie AI,Onuoha GC. Effects of Jackbean Seed Meal on the Intestinal Mucosa of Juvenile *Heterobranchuslongifilis*. Afr. J. Biotech.2006; 5(13): 1294-1298.
12. SoetanKO, AntiaRE. Comparative phytochemicals and *in vitro* antioxidative effects of jack beans (*Canavaliaensiformis*) and sword beans (*Canavaliagladiata*). Annals. Food Sci. Techn. 2018; 19(3): 499-505.
13. AOAC. Official Methods of Analysis of AOAC International, (17th 505 Edition), 506 Association of Official Analytical Chemists, Inc, Gaithersburg. 2003.
14. Samant SK,Rege DV.Carbohydrate Composition of Some Cucurbit Seeds. J. Food Comp. Anal. 1989;2:149-156.
15. Dubois M, Gilles KA, Hamilton JK, Rebers PA, Smith F.Colorimetric method for determination of sugars and related substances. Anal. Chem.1956; 28:350-356.

16. Bernfeld P. Amylase α and β . Methods in enzymology 1.S. P. Colswick and N.O.K., Ed. Academic Press Inc, New-York. 1955; 149-154.
17. Crisan EV, Sands A. The biology and cultivation of edible mushrooms, Nutritional value, Academic Press, New York. 1978; 137-168.
18. APHA. Standard methods for examination of water and waste water, 19th ed., American Public Health Association, USA; 1995; 113-11 p.
19. Aremu MO, Basu SK, Gyar SD, Goyal A, Bhowmik PK, Datta BS. Proximate Composition and Functional Properties of Mushroom Flours from *Ganoderma spp.*, *Omphalotusolearius*(DC.) Sing. And *Hebelomamesophaeum*(Pers.) Quél. Used in Nasarawa State, Nigeria. Mal. J. Nutr. 2009; 15(2):233–241.
20. Anino C, Onyango AN, Imathiu S, Maina J, Onyangore F. Chemical composition of the seed and 'milk' of three common bean (*Phaseolusvulgaris L*) varieties. J. Food Measur. Charact. 2019; 13: 1242–1249.
21. Dué EA, Koffi DM, Digbeu YD. (a). Biochemical and functional properties of two wild edible mushrooms (*Volvariellavolvacea* and *Armillariamellea*) consumed as protein substitutes in South Côte d'Ivoire. J. Chem. Biol. Phys. Sci. 2016; 6(4): 1197 – 1206.
22. Dué EA, Koffi DM, Digbeu YD. (b). Physicochemical and Functional Properties of Flour from the Wild Edible Mushroom *Termitomycesheimii* Natarajan Harvested in Côte d'Ivoire. Turk. J. Agr. Food Sci. Techn. 2016; 4(8): 651–655.
23. Banjo AD, Lawal OA, Songonuga EA. The nutritional value of fourteen species of edible insects in Southwestern Nigeria. Afr J Biotechnol. 2006; 5(3): 298- 301.
24. Osasona AI, Olafe O. Nutritional and functional properties of *Cirinafordalarva* from Ado-Ekiti, Nigeria. Afr J Food Sci. 2010; 4(12): 775–777.
25. Gouveia CSS, Freitas, De Brito JH, Slaski JJ, De Carvalho MAAP. Nutritional and mineral variability in 52 accessions of common bean varieties (*Phaseolusvulgaris*L.) from Madeira Island. Agric. Sci. 2014; 5: 317–329.
26. Los FGB, Zielinski AAF, Wojeicichowski JP, Nogueira A, Demiate IM. Beans (*Phaseolusvulgaris L.*): whole seeds with complex chemical composition. Curr. Opin. Food Sci. 2018; 19: 63–71. doi:10.1016/j.cofs.2018.01.010.
27. Narasinga-Rao BS, Deosthale YG, Pant KC. *Nutritive value of Indian foods*. Indian Council of Medical Research, National Institute of Nutrition, Hyderabad, India. 1989; Pp. 24 - 48.
28. Kan L, Nie S, Hu J, Wang S, Bai Z, Wang J, Zhou Y, Jiang J, Zeng Q, Song K. Comparative study on the chemical composition, anthocyanins, tocopherols and carotenoids of selected legumes. Food Chem. 2018; doi: <https://doi.org/10.1016/j.foodchem.2018.03.148>.
29. Pedrosa MM, Cuadrado C, Burbano C, Muzquiz M, Cabellos B, Olmedilla-Alonso B, Asensio-Vegas C. Effects of industrial canning on the proximate composition, bioactive compounds contents and nutritional profile of two Spanish common dry beans (*Phaseolusvulgaris L.*). Food Chem. 2015; 166: 66-75.
30. Butkute Bronislava, Taujenis Lukas and Norkeviciene Egle. Small-Seeded Legumes as a Novel Food Source. Variation of Nutritional, Mineral and Phytochemical Profiles in the Chain: Raw Seeds-Sprouted Seeds-Microgreens. Molecules. 2019; 24: 133 doi:10.3390/molecules24010133.
31. Pivic NW, Butler JB. 1977. A simple unit leaf. Proc Nutr Soc. 12:36: 136.
32. Omowunmi OO, Oluwatosin AY, Adesola BN. Chemical analysis of the seed oil of *Canavaliaensiformis* Linn. for nutritional and industrial qualities. Afr J Sci Nat. 2018; 6: 66-72.
33. Vadivel V. The nutritional and antioxidant contents of wild jack bean (*Canavaliaensiformis L. DC.*): an under-exploited legume from south India. Int. J. Rec. Scient. Res. 2019; 10: 35502-35508

34. Nkwocha, G.A.1, Anukam, K. U.1 and Adumanya, O.C. (2021). Evaluation of the Nutritional Potentials of Three Species of *Canavalia* for Use in Livestock Diets. *Journal of Sustainable Technology*, 11(2): 86-92.
35. NRC-NAS. (1989) Recommended dietary allowances. National Academy Press, Washington DC.
36. Akpoussan RA, Digbeu YD, Koffi DM, Kouadio JPEN, DuéEA, Kouame PL. 2015. Protein Fractions and Functional Properties of Dried *Imbrasia oyemensis* Larvae Full-Fat and Defatted Flours. *Inter J Biochem Res Rev*. 5(2):116-126.
37. Assiélou B, Dué EA, Koffi MD, Dabonné S, Kouamé PL. 2015. *Oryctesowariensis* Larvae as Good Alternative Protein Source: Nutritional and Functional Properties. *An Res Rev Biol*. 8(3): 1-9.
38. Keskin SO, Ali TM, Ahmed J, Shaikh M, Siddiq M, Uebersax MA. Physico-chemical and functional properties of legume protein, starch, and dietary fiber—A review. *Leg. Sci*. 2022; 4: 1-15.
39. Okomoda VT, Tihamiyu LO, Uma SG. Effects of hydrothermal processing on nutritional value of *Canavalia ensiformis* and its utilization by *Clarias gariepinus* (Burchell, 1822) fingerlings. *Aquac. Rep*. 2016;3: 214-219.
40. Akande KE, Doma UD, Agu HO, Adamu HM. Major Antinutrients Found in Plant Protein Sources: Their Effect on Nutrition. *Pak. J. Nut*. 2010; 9: 827-832.
41. Cheng JZ, Hardy RW, Usry JL. Plant protein ingredients with lysine supplementation reduce dietary protein level in rainbow trout (*Oncorhynchus mykiss*) diets, and reduce ammonia nitrogen and soluble phosphorus excretion. *Aquaculture*. 2003; 218: 553-565.
42. Millar KA, Gallagher E, Burke R, McCarthy S, Barry-Ryan C. Proximate composition and anti-nutritional factors of fava-bean (*Vicia faba*), green-pea and yellow-pea (*Pisum sativum*) flour. *J. Food Comp. Anal*. 2019;82(), 103233–. doi:10.1016/j.jfca.2019.103233
43. WHO/FAO/UNU. Protein and amino acid requirements in human nutrition: report of a joint FAO/WHO/UNU expert consultation. WHO Technical Report Series. Place Published: World Health Organization. 2019.