

1
2
3
4
5
6
7
8
9
10
11
12

Form and Function in Biological Macromolecules: Kinetic Stability is Key, with Oblique Roles for Intramolecularity and Hydrophobicity in Enzyme Catalysis

ABSTRACT

Certain structure-reactivity aspects of biological macromolecules, with particular emphasis on protein folding and enzyme catalysis, are discussed herein. Furthermore, the role played by the hydrophobic effect and intramolecularity in enzymic reactivity are evaluated afresh, with new insights of much importance in chemical biology.

Thus, the sum of the energies of the hydrogen bonds constituting the tertiary structures of proteins, determines the overall Gibbs energy of activation for loss of conformational integrity. As protein molecules of even modest size consist of a relatively large number of intramolecular hydrogen bonding interactions, the activation barrier to even partial unfolding of the α -helices and β -sheets forming the tertiary structure would be prohibitively high under normal conditions.

The resulting kinetic stability conserves the natural conformation of a protein molecule established at the ribosomal site of synthesis, carrying the molecule through the thick-and-thin of a range of metabolic pathways during its 'journey of life'. However, protein molecules also acquire flexibility via 'strain delocalization' (Ramachandran plots being relevant), thus enabling stabilization of multiple transition states along a pathway (particularly in case of covalent enzyme-substrate complexes).

Two mechanistic features of enzyme catalysis that have been exhaustively studied are intramolecularity and the hydrophobic effect. Although intramolecularity has for long been touted as the origin of enzymic reactivity, this can be challenged on fundamental physical-organic grounds. Intriguingly, however, the collapse of the classical Michaelis-Menten mechanism for enzyme catalysis leads to a reconsideration of the role of intramolecularity, although not as hitherto envisaged. Thus, a majority of enzymes apparently form covalent enzyme-substrate complexes—possibly also exergonically—so the subsequent reactions at the active site may well benefit from the traditional propinquity effect: The critical caveat would be the highly exergonic formation of final products.

It is argued that the hydrophobic effect—although intuitively reasonable—is difficult to pin down quantitatively, model systems (including micelles) leading to inconsistent and debatable results. However, the hydrophobic effect likely contributes to enzymic reactivity along with charge-relay via the proteinic backbone.

13 *Keywords: Hydrogen bonding, Michaelis-Menten, micelles, propinquity effect, protein folding,*
14 *Ramachandran plots*
15

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68

1. INTRODUCTION

1.1 General background

A fundamental understanding of the structure and function of enzymes is of critical importance in the evolving field of chemical biology [1-5]. Thus, enzymes represent a key nodal point in the manifestation of genetic information as the myriad of metabolic processes that constitute life. This derives essentially from the one gene-one enzyme hypothesis, which intriguingly delegates to the enzymes an awesome importance as the key to the efflorescence of life itself. It is thus hardly surprising that enzyme function and reactivity have served as a focus of fundamental enquiry and debate, with chemistry—the ‘central science’—leading the way in this greatest of reductionist adventures.

Progress in mechanistic enzymology has been predicated on advances in protein science—in theory and practice—and understandably so. Pioneering work, by Buchner on yeast-mediated fermentations (1897) and kinetic studies by Michaelis and Menten (1913), represented the first stirrings of the nascent science of enzymology [4]. However, advances in protein isolation, purification and characterization, not to mention the founding of molecular biology in the ensuing decades, subsequently laid the foundations of modern enzymology.

The near-miraculous catalytic capabilities of enzymes also began to engage the attention of chemical science, which was itself increasing in sophistication in parallel with advances in biology. A synergistic approach between chemical and biological science thus emerged and has continued to evolve since the de facto founding of ‘chemical biology’ ~ 1950. The realization that enzyme catalysis was in a class of its own elicited the Pauling theory of transition state stabilization (1948) [5], followed by the determination of the structure of DNA (1953) and the founding of molecular biology [6]. The elucidation of the genetic code in terms of transcription (to mRNA) and translation (to protein) indicated that the proteinic enzymes are destined to hold the key to the mystery of dynamic life itself.

However, although the urgent need to reach a fundamental understanding of enzyme catalysis appears self-evident, the complex nature of enzymes and theoretical uncertainties of chemical science have conspired to envelop enzymology in controversy (if not shrouding it in more mystery)! Thus, the classical Michaelis-Menten equation that served as a bed-rock of enzyme kinetics for a century has been challenged [4]; the Pauling hypothesis remains to be elucidated in mechanistic detail; and physical-organic models particularly based on intramolecularity [7] and hydrophobic effects [8]—however ingenious—have apparently floundered on fundamental grounds.

This paper briefly reviews this essential background and argues that enzymology is indeed emerging from the state of flux, although a bold and frank assessment of previous inadequacies is needed to connect the dots and forge ahead to a new era of chemical biology.

1.2 Thermodynamic considerations

Modern theories of enzyme catalysis and action are based on the assumption that, despite their biological origins and often forbidding molecular complexity, enzymes are subject to the same laws as apply to in vitro catalysis by small inorganic and organic molecules. These theories essentially derive from the classical law of mass action that is the basis of chemical equilibrium and reaction rate. The thermodynamic versions of these theories relate to the

69 Boltzmann equation and its formulation in terms of the Gibbs energy, itself composed of
70 enthalpy and entropy contributions [9]. The key idea that a catalyst is regenerated during the
71 reaction—and does not change the equilibrium constant either—is also retained in the case
72 of the enzyme reaction.

73
74 These approaches—shorn of all traces of vitalism—thus hold the key to the reduction of
75 sentient life to the laws governing the properties of inanimate matter.

76 77 **2. DISCUSSION**

78 79 **2.1 Uniqueness of enzyme catalysis. Recalcitrant problems**

80 81 **2.1.1 The dual role of enzymes: as ‘evolutionary fulcrum’ and as metabolic workhorse**

82
83 Enzymes are generally medium-sized proteins that are direct products of genetic expression,
84 representing the first stage in the conversion of the genetic code to a non-nucleic acid
85 information system. Thus, enzymes serve as an ‘evolutionary fulcrum’ that, however, has
86 apparently obfuscated the fact that their proteinic nature is also critical to their catalytic
87 power! In other words, smaller non-protein catalysts are unlikely to rival the catalytic powers
88 of natural enzymes, as is becoming increasingly apparent.

89
90 However, the full significance of enzymes can only come to light if the origins of their
91 catalytic powers are fully understood. This is much more than the mere cataloguing of the
92 various catalytic groups and their mechanistic interplay, which have indeed been replicated
93 in model systems [10]. Yet, even the most ingeniously designed models have failed to
94 approach the catalytic powers routinely observed in natural enzymes: As it stands, the whole
95 remains greater than the sum of its parts!

96
97 Thus, enzyme catalysis is characterized by enormous rate enhancements (typically $> 10^{10}$)
98 that are brought about under the mildest of aqueous conditions [1-5]. Hence, they serve as
99 the metabolic work-horses of life. Their remarkable features have elicited various
100 mechanistic theories as explanations, noting that an important practical goal is the design of
101 artificial catalysts that can rival the catalytic powers of enzymes. Indeed, this remains the
102 holy grail of physical organic chemistry, in the service of chemical biology and science itself.

103 104 **2.1.2 Physical organic models: Intramolecularity and the hydrophobic effect**

105
106 Two important mechanistic approaches to enzyme catalysis invoke intramolecularity and the
107 hydrophobic effect [11]. These have been implicated in enzyme catalysis essentially
108 because they have led to enhanced rates in model systems, but also because they
109 apparently possess some of the characteristics of enzyme catalysis. Thus, enzyme catalysis
110 occurs via an enzyme-substrate complex, its further ‘turnover’ to product being necessarily
111 intramolecular; and enzyme active sites are hydrophobic pockets that are apparently
112 complementary to the rate-determining transition state of the reaction.

113 114 ***2.1.2.1 Intramolecularity***

115
116 However, the intramolecularity model has been challenged as enzyme catalysis is
117 bimolecular (considering only enzyme and substrate) [7], whereas intramolecular reactions
118 are unimolecular. Thus, the enhanced rates of intramolecular reactions are due to a raised
119 ground state (often entropic in origin but also enthalpic in certain cases), whereas enzyme
120 reactions must involve transition state stabilization in some manner.

121

122 It is also possible that catalysis involves the energy of the catalyst itself in some **form**,
123 although this is not so apparent as the catalyst is regenerated. Thus, the catalyst could be a
124 source of potential energy, but problems remain with the exact interpretation of transition
125 state theory as to whether the Gibbs energy or the potential energy is to be employed [12].
126

127 In fact, a raised ground state ostensibly justifies the intramolecularity model, the conundrum
128 itself possibly indicating the limitations of chemical theory in explaining complex biological
129 phenomena! All the same, the intramolecularity model is intriguingly resuscitated in light of
130 the recent reinterpretation of enzyme kinetics, as discussed further below.
131

132 **2.1.2.2 The hydrophobic effect**

133

134 The view that the hydrophobic effect plays a major role in enzyme catalysis has also been
135 debated [8], the results of model studies being difficult to interpret. Although the studies
136 demonstrate enhanced rates of certain reactions (particularly cycloadditions) in water, the
137 results are stymied by the low solubility of the hydrophobic reactants in water! The results
138 are thus valid in select cases, apparently, with water catalysis being a viable option that
139 cannot be easily ruled out. Generally, in fact, a hydrophobic medium is non-polar and thus
140 not conducive to enhanced reactivity: Unsurprisingly, enhancements have been observed in
141 the case of concerted reactions, polar reactions likely requiring supplementary charge relay
142 effects as may occur in enzymes (and possibly micellar systems too, *vide infra*).
143

144 Thus, the hydrophobic effect observed in certain model systems is possibly misleading. The
145 results evidence the intuitively reasonable idea that a reaction in which the transition state is
146 less hydrophobic than the ground state is accelerated in water (relative to an organic
147 medium). However, the greater hydrophobicity of the substrate implies a correspondingly
148 lower solubility, so the rate enhancements are not practically significant. In fact, substantial
149 accelerations are also apparently observed in organic-water emulsions, which likely implies
150 interfacial catalysis by water that cannot be separated from the hydrophobic effect *per se*.
151 Also, the hydrophobic effect is not a general accelerating effect applying to all reactions in
152 aqueous media, as is often assumed!
153

154 **2.2 Enzymes as proteins**

155

156 **2.2.1 Macromolecular nature of enzymes: Protein folding and activity**

157

158 Biological macromolecules manifest overall three-dimensional shape as biological function
159 [13-16], thus apparently representing a twilight zone: between synthetic high polymers that
160 are characterized by their mechanical properties (e.g., polyethylene [17]), and smaller
161 oligomeric fragments of no defined form or function. In fact, a complex and intimate
162 relationship between form and function, particularly among the ubiquitous proteins
163 (enzymes, wool, skin, etc.), distinguishes biological macromolecules from their smaller and
164 larger congeners.
165

166 Furthermore, among the three major classes of biological macromolecule—nucleic acids,
167 proteins and carbohydrates—proteins are the warp and woof of the central dogma of
168 biological information flow [6]: DNA – RNA – protein – organism. However, whilst the
169 proteinic enzymes control and regulate every aspect of life's complex machinery, they also
170 possess a myriad of conformational states. Clearly, only one among these states is
171 apparently enforced at the ribosomal site of biosynthesis [18]: but what helps retain the
172 overall shape thus conferred? The answer holds the key to the mystery of life's sustenance,
173 based in the unerring consistency of an organism's biochemical complexity [19].
174

175 And intriguingly, the overall conformation conferred need not be the thermodynamically most
176 stable one, else both function and stability would need to be optimized by evolution! Indeed,
177 the degeneracy of the genetic code could imply—via codon-specific t-RNA-synthetases—
178 changes in the ribosomal micro-environment, which lead to different tertiary structures being
179 adopted by the same polypeptide primary sequence! Furthermore, a subtle balance between
180 flexibility and rigidity is of critical importance to macromolecular function and enzyme action
181 in particular, as argued below.

182 **2.2.2 Protein dynamics and enzyme action: Ramachandran plots**

183 **2.2.2.1 Polypeptide flexibility: mechanisms and catalytic role**

184
185 As noted above, the mystery of enzyme catalysis is predicated on an understanding of
186 protein structure and dynamics. Of particular importance to enzyme catalysis is the
187 possibility of conformational changes that would lead to a flexible active site, which can
188 stabilize several different transition states along a reaction pathway. The possibility of charge
189 relay along the protein backbone has also been proposed as a way of stabilizing polar
190 transition states in a hydrophobic environment [4]. (Indeed, the presence of the catalytic triad
191 in proteases indicates the importance of charge-relay in a hydrophobic environment [4].)
192 That proteins strike a balance between rigidity and flexibility via 'strain delocalization' has
193 been proposed as being critical to the reactivity of catalytic groups around the active site
194 (vide infra).

195 **2.2.2.2 The concept of strain delocalization**


196
197 An intriguing and novel mechanism by which a polypeptide backbone can acquire flexibility
198 is the possibility of strain delocalization [4]. This is possible as bond angle strain increases
199 exponentially with decreasing angle, so the distribution of angle strain over several angles is
200 thermodynamically favored over the concentration of the same angle strain in a single or
201 fewer angles. The consequent 'delocalization of strain' is indeed feasible in macrocyclic and
202 pseudo-macrocyclic systems, e.g., in a relatively long polypeptide chain. Strain
203 delocalization, in fact, would enable precise reaction trajectories to be attained by interacting
204 catalytic groups in the active site. Indeed, this possibly explains the macromolecular nature
205 of enzymes, as smaller molecules fail to show similar catalytic powers.

206 **2.2.2.3 Ramachandran plots: infinite thermodynamic degeneracy**

207
208 Intriguingly, in the absence of any kinetic barriers, protein molecules are infinitely flexible, as
209 is apparent in the well-known Ramachandran plots [15,16,18]. These contour diagrams
210 essentially imply that an infinite range of dihedral angles (ϕ , ψ) around the peptide bonds in
211 a polypeptide are acceptable. Thus, to the extent that the contours are not discontinuous,
212 they indicate that an amino acid residue can adopt an infinite range of conformations—within
213 an allowed subset—that are thermodynamically degenerate. These arguments can be
214 extended to whole molecules with due caveats, leading to 'impressionistic' diagrams
215 displaying smudged distributions of nearly overlapping data points, indicating a high level of
216 thermodynamic degeneracy.

217 **2.2.2.4 Kinetic stability of polypeptide conformers: additivity of hydrogen bond strengths**

218
219
220
221
222
223
224



225
226

227 **Figure 1. Gibbs energy (G) changes during the unfolding of a polypeptide fragment**

228
229

230 Whilst Ramachandran plots imply infinite thermodynamic flexibility, they also indicate the
231 importance of kinetic barriers to the maintenance of the tertiary structures of proteins! A
232 major part of the kinetic stability of a polypeptide would undoubtedly be provided by the
233 network of intramolecular hydrogen bonds—literally the molecular scaffolding—that forms
234 the α -helices and β -sheets, but also arranges them in molecular space [13-16].

235
236
237
238

Indeed, notwithstanding the mildness of biological conditions—and the strength of the
peptide bond—non-covalent forces control protein folding, its retention through thick and thin
during a cell's life cycle thus representing a remarkable feat of molecular endurance!

239
240
241
242
243
244
245

The key to this “conundrum of protein folding”, in fact, lies not just in the network of hydrogen
bonds that maintains a polypeptide conformation, but also in its cumulated thermodynamic
effect (cf. Fig. 1). Thus, a network of n hydrogen bonds is stabilized by the strength of the
first $(n-1)$ hydrogen bonds, and by the barrier to the cleavage of the last hydrogen bond: The
sum of all these represents the activation. barrier to the dismantling of the overall
polypeptide conformation, i.e., its kinetic stability [20]. (In Fig. 1, $n = 4$, $E_1 - E_4$ being the
stated hydrogen bond energies.)

246
247
248
249
250
251

Thus, a single turn of the α -helix would be stabilized by a sizable barrier $> 20 \text{ kcal mol}^{-1}$
(based on 7 hydrogen bonds worth 3 kcal mol^{-1} each [13-16]). Clearly, even moderately
sized proteins composed of several α -helix and β -sheet units would possess forbidding
barriers to unfolding. Little wonder that the denaturation of proteins requires particularly
harsh conditions and that controlled unfolding of proteins is catalysed by ATP-dependent
enzymes [21]!

252 **2.3 Enzymes as catalysts**

253

254 **2.3.1 Enzyme kinetics: collapse and renewal**

255

256 ***2.3.1.1 The Michaelis-Menten equation and its aftermath***

257

258 The pioneering studies of enzyme kinetics by Michaelis and Menten (1913) represent an
259 intrepid foray into the then evolving fields of biological catalysis and chemical kinetics [11].
260 Early observations indicated that enzyme catalysis followed 'saturation kinetics', wherein the
261 reaction rate (ν) initially increased linearly with the substrate concentration ($[S]$), but levelled
262 off asymptotically at high $[S]$. This led to the proposal that the reaction occurred via the rapid
263 pre-equilibrium formation of an enzyme-substrate complex (ES), which 'turned over'
264 relatively slowly to products, enshrined in the Michaelis-Menten rate law (Eq. 1), $[E_o]$ being
265 the (constant) initial enzyme concentration, k_{cat} the turnover number and K_M the Michaelis
266 constant.

267

268
$$\nu = k_{cat}[E_o][S]/(K_M + [S]) \quad (1)$$

269

270
$$\nu = k_{cat}[E_o] \quad (2)$$

271

272
$$K = k_{cat}/k_{cat}' \quad (3)$$

273

274
$$K = [P]/[S] \quad (4)$$

275

276
$$(k_{cat}/k_{cat}') \neq [P]/[S] \quad (5)$$

277

278

279 ***2.3.1.2 Inadequacies of the Michaelis-Menten equation***

280

281 Interestingly, Eq. 1 reduces to Eq. 2 at high $[S]$, apparently reproducing the experimentally
282 observed 'saturation kinetics'. This ostensibly indicated the gradual saturation of the pre-
283 equilibrium that was nearly complete at high $[S]$.

284

285 However, a serious problem with Eq. 2 is that it leads to Eq. 3 under conditions of overall
286 equilibrium between substrate and product (P) (at high $[S]$ and $[P]$, K is the overall
287 equilibrium constant and k_{cat}' the turnover number for the reverse reaction).

288

289 Clearly, Eq. 3 is in conflict with the natural equilibrium constant of the reaction (K , Eq. 4).
290 Indeed, this led to the proposal of 'one-way' enzymes although this idea contravened a
291 fundamental tenet of thermodynamics that the equilibrium constant depends only on the
292 Gibbs energy change in the reaction (at constant temperature, Eq. 5 representing the
293 general case).

294

295 **2.3.2 Alternatives to the Michaelis-Menten scheme: Inhibition at high $[S]$ via**
296 **secondary binding**

297

298 In view of the above anomalies, the Michaelis-Menten formulation of enzyme kinetics is
299 fundamentally invalid and needs to be abandoned. In fact, a viable alternative that has been
300 proposed is based on the idea of a gradual inhibition of the enzyme at high substrate
301 concentrations that leads to the observed levelling of the rate [4]. Such inhibition is almost
302 certainly likely to occur via the binding of a second molecule of substrate at the active site,
303 which would prevent the release of product formed and regeneration of enzyme.
304 Accordingly, Eq. 6 was proposed as an alternative to the Michaelis-Menten equation (k is the

305 overall reaction rate constant, K_S the equilibrium constant for the secondary binding of
306 substrate and E_F the free enzyme).

307

$$308 \quad v = k[E_o][S] - kK_S[E_F][S]^3 \quad (6)$$

309

310 In Eq. 6, the first term on the right-hand side refers to the linear increase in rate with
311 increasing [S], whereas the second term refers to the inhibition via secondary binding. Thus,
312 K_S is low as secondary binding is weak, hence the second term becomes significant at high
313 [S], increasing exponentially to 'rein in' the first term. An analogous equation can be written
314 for the reverse reaction, leading to the thermodynamic equilibrium constant, noting that the
315 forward and reverse reactions are catalyzed only by E_F .

316

317 Thus Eq. 6 captures the observed levelling effect of the rate at high [S] and also leads to the
318 thermodynamically valid equilibrium constant. It is also noteworthy that, although Eq. 6 was
319 originally derived for a pre-equilibrium formation of the enzyme-substrate complex, this
320 assumption is strictly not necessary as the rate constant k is per se not based on the pre-
321 equilibrium assumption. Therefore, Eq. 6 is generally valid regardless of the relative stability
322 of the enzyme-substrate complex, thus leading to various mechanistic possibilities for further
323 consideration.

324

325 **2.3.3 Exergonic formation of the enzyme-substrate complex and intramolecularity**

326

327 ***2.3.3.1 Endergonic and exergonic enzyme-substrate binding; covalent bond formation***

328

329 As noted above, the Michaelis-Menten equation was derived on the basis of a pre-
330 equilibrium mechanism, in which a weakly-bound enzyme-substrate complex was formed
331 endergonically. (The observed levelling of rate was believed to arise from the putative
332 saturation of the pre-equilibrium.) However, in view of the collapse of the Michaelis-Menten
333 regime in toto, alternative mechanisms can now be considered.

334

335 An interesting possibility is the exergonic formation of the enzyme-substrate complex from
336 enzyme and substrate. In fact, this mechanism acquires particular significance in light of
337 findings that a substantial number of enzymes form covalent enzyme-substrate complexes
338 [5]. An interesting advantage of covalent enzyme-substrate binding is that it allows for
339 conformational changes in the enzyme, so the active site would be flexible enough to
340 stabilize all possible transition states in the reaction pathway.

341


342 **The covalent complex, however,** may also be formed endergonically. Whilst this indicates
343 that the strength of the covalent bond linking enzyme and substrate is carried over to the
344 transition state, this can contribute to the catalysis only if the binding is stronger in the
345 transition state relative to the ground state. This implies that the ground state complex is
346 strained in some manner, the strain being relieved on reaching the transition state.

347

348 However, a more likely possibility is that the covalently linked complex is formed
349 exergonically, considering the strength of a covalent bond. In the earlier Michaelis-Menten
350 scheme, however, such exergonic formation of the enzyme-substrate complex was ruled
351 out, essentially because the turnover step would be slower rather than faster (as would also
352 be release of enzyme). (This was the 'thermodynamic pit' problem that apparently negates
353 catalysis [11].) However, this mechanism now merits serious consideration for several
354 reasons, not least of which is the intriguing involvement of intramolecularity (vide infra).


355

356 The exergonic formation of an enzyme-substrate complex, however, would be predicated on
357 the overall reaction itself being highly exergonic, in order to avoid a buildup of the said



358
359
360
361
362
363
364
365

Fig. 2. Gibbs energy (G) profile for the rate-limiting exergonic formation of an enzyme-substrate complex (ES)



366
367
368
369
370

Fig. 3. Gibbs energy (G) profile for the rapid and exergonic formation of an enzyme-substrate complex (ES)

371 complex. All the same, this may not be a serious limitation as—for obvious practical
372 reasons—the majority of enzyme catalyzed reactions are perforce exergonic. (Like any
373 catalyst, an enzyme cannot alter the equilibrium constant of a reaction; however, even an
374 endergonic equilibrium can be displaced by further reactions of the product.)
375

376 An interesting problem with the exergonic formation of the enzyme-substrate complex,
377 however, is noteworthy. This is the possibility that the initial exergonic pre-equilibrium would
378 be partially reversed with an increase in temperature, leading to a subdued (or even
379 negative) temperature dependence of the overall rate [3,20]. Apparently, however, the
380 positive temperature dependence of the subsequent sub-reactions generally overcomes the
381 reversal of the initial pre-equilibrium. Also, the reversal may not occur if the by-product (not
382 shown) in the formation of ES_1 (Figs 2 and 3, vide infra) is concomitantly expelled from the
383 active site. (Any reversal, however, would lead to an observed Gibbs energy of activation
384 that would be lower than the real one.)
385

386 **2.3.3.2 Intramolecularity makes a comeback: the ES complex is the ground state!**

387

388 The energy profile for exergonic enzyme-substrate binding—whether covalent or not—is
389 shown in Fig. 2. In a fascinating mechanistic flip, it is now observed that the enzyme-
390 substrate complex (ES) represents the ground state of the reaction! An intriguing
391 consequence is that the subsequent clutch of catalytic reactions would possess all the
392 virtues of intramolecularity, so the traditional propinquity effect can be invoked as
393 contributing to the overall rate enhancement. The mechanistic reasoning is of considerable
394 subtlety, and only becomes apparent upon comparison with a putative profile in which—
395 hypothetically—intramolecularity does not play a role (dashed lines in Fig. 2).
396

397 An important caveat to these arguments, however, is that the formation of the covalent
398 enzyme-substrate complex, as also the final release of product, have become rate-
399 determining. The corresponding transition states, of course, do not benefit from
400 intramolecularity, and are thus subject to the Pauling hypothesis. In other words, the enzyme
401 now needs to stabilize only two extreme transition states in the overall profile, the propinquity
402 effect dealing with the key catalytic reactions occurring within the confines of the active site.
403 (Thus, the enzyme need not necessarily stabilize the several intermediate transition states!)

404
405 An interesting variation is the case wherein formation of the enzyme-substrate complex is
406 exergonic, but fast (possibly diffusion-controlled) relative to the intermediate intramolecular
407 steps (Fig. 3). (The release of product, likewise, may also be fast.) The intramolecular steps
408 would then be rate-determining with the overall rate being dependent on the stability of the
409 enzyme-substrate complex. Thus, in the (presumed) absence of any stabilization of the
410 intermediate transition states, an excessively stable enzyme-substrate complex would
411 detract from the overall rate enhancement. It is noteworthy that intramolecular reactions
412 show an enormous range of effective molarity (EM) values that reflect—inter alia—
413 correspondingly raised ground states.
414

415 It is particularly noteworthy that these arguments were impossible in the framework of the
416 Michaelis-Menten scheme, as it was based on the endergonic formation of the enzyme-
417 substrate complex, with the ground state being represented by free enzyme and substrate.
418 Intramolecularity was hence indefensible in that context, with transition state stabilization
419 being sine qua non for all steps.
420

421 Furthermore, the alternative kinetic proposal (Eq. 6) is broadly applicable to both cases, i.e.,
422 endergonic and exergonic formation of the enzyme-substrate complex (ES). This is apparent
423 from the presence of the unitary rate constant k in Eq. 6: In the case of exergonic formation

424 of ES, k would refer to the formation of ES itself; in the case of endergonic formation of ES, k
425 would refer to the breakdown of ES to products. Both cases would be valid without prejudice
426 to the observed levelling of rate, as inhibition via secondary binding would apply equally.

427

428 **3. CONCLUSIONS**

429

430 The above discussion has briefly reviewed an extensive range of topics of fundamental
431 significance to the origins of enzyme catalysis. The inevitable conclusion, apparently, is that
432 mechanistic enzymology has entered into a state of flux, from which it is indeed beginning to
433 emerge. The major conclusion is that the classical theory based on the Michaelis-Menten
434 equation stands discredited and needs to be replaced by a credible alternative.

435

436 Currently, a viable mechanism based on secondary binding of substrate at high
437 concentrations appears reasonable. This is in accord with the essential theory of chemical
438 equilibrium and kinetics, and can also explain the levelling of the rate at high substrate
439 concentrations. Although the Pauling theory of transition state stabilization remains
440 paramount, alternatives also may now be considered, in particular the intramolecularity
441 concept. The hydrophobic effect also plays a role, although along with a charge-relay
442 mechanism possibly involving the protein backbone.

443

444 The proteinic nature of enzymes also needs to be accorded more serious consideration
445 henceforth. In particular, the kinetic stabilization of the tertiary structure afforded by the
446 network of hydrogen bonds plays foil to the inherent flexibility of a polypeptide chain. The
447 classical theory of protein structure, particularly based on the Ramachandran plots, may be
448 augmented by newer ideas (e.g., strain delocalization). These lead to novel insights into the
449 dynamics of polypeptide molecules likely of key significance to enzyme catalysis.

450

451 In summary, an integrated approach involving ideas from both mechanistic chemistry and
452 polypeptide dynamics is indicated for a comprehensive understanding of enzyme catalysis to
453 be reached. (Indeed, a palette of mechanistic options is now apparent, with each enzyme
454 employing the most appropriate mechanism to fulfill its metabolic purpose.)

455

456 **CONFLICT OF INTEREST**

457

458 The contents of this article bear no conflicts of interests.

459

460 **REFERENCES**

461

462 1. Agarwal PK. A biophysical perspective on enzyme catalysis. *Biochemistry*
463 2019;58(6):438–449. (DOI: 10.1021/acs.biochem.8b01004)

464 2. Kirby AJ, Hollfelder F. *From enzyme models to model enzymes*. Cambridge (UK): Royal
465 Society of Chemistry; 2009.

466 3. Chandrasekhar S. The origins of enzyme catalysis and reactivity: Further assessments.
467 *Asian Journal of Chemical Sciences*. 2021;9(3):38-47. (DOI:
468 10.9734/AJOCS/2021/v9i319075)

469 4. Chandrasekhar S. Understanding enzymic reactivity – new directions and
470 approaches. *Asian Journal of Research in Biochemistry*. 2020; 7(2):1-13. (DOI:
471 10.9734/ajrb/2020/v7i230133)

- 472 5. Zhang X, Houk KN. Why enzymes are proficient catalysts: beyond the Pauling paradigm.
473 Acc Chem Res. 2005;38(5):379-385. (DOI: 10.1021/ar040257s)
- 474 6. Watson, J.D.; Baker, T.A.; Bell, S.P.; Gann, A.; Levine, M.; Losick, R. Molecular Biology of
475 the Gene. 7th ed. Boston: Pearson; 2013.
- 476
- 477 7. Chandrasekhar S. Intramolecularity and enzyme modelling: a critique. Res Chem
478 Intermed. 2003;29(1):107-123. (DOI: 10.1163/156856703321328451)
- 479
- 480 8. Chandrasekhar S. The hydrophobic effect in chemistry, biology and medicine: An update.
481 Asian Journal of Chemical Sciences. 2021;9(4):22-36. (DOI:
482 10.9734/AJOCS/2021/v9i419078)
- 483 9. Atkins P, de Paula J. Atkins' physical chemistry. 8th ed. New York: W. H. Freeman; 2006.
- 484 10. Silverman RB. The organic chemistry of enzyme-catalyzed reactions. San Diego:
485 Academic Press; 2002.
- 486 11. Fersht A. Structure and mechanism in protein science. 2nd ed. Cambridge (UK): Kaissa
487 Publications; 2017.
- 488 12. Chandrasekhar S. The Gibbs Energy–Potential Energy Conundrum and Chemical
489 Reactivity: Implications for Catalysis and Enzyme Action. Asian Journal of Chemical
490 Sciences 9(4): 1-7, 2021. (DOI: 10.9734/AJOCS/2021/v9i41907)
- 491 13. Dill KA, MacCallum JL. The protein folding problem, 50 years on. Science. 2012;
492 338(6110):1042-1046. (DOI: 10.1126/science.1219021)
- 493
- 494 14. Kuriyan J, Konforti B, Wemmer D. The Molecules of Life: Physical and Chemical
495 Principles. New York: Garland Science; 2013.
- 496
- 497 15. Kessel A, Ben-Tal N. Introduction to Proteins Structure, Function and Motion. 2nd ed.
498 Boca Raton: CRC Press; 2018.
- 499
- 500 16. Buxbaum E. Fundamentals of Protein Structure and Function. New York: Springer
501 Science; 2017.
- 502
- 503 17. Koltzenburg S, Maskos M, Nuyken O, Hughes K. Polymer Chemistry. Berlin-Heidelberg:
504 Springer-Verlag; 2017.
- 505
- 506 18. Rosenberg AA, Marx A, Bronstein AM. Codon-specific Ramachandran plots show amino
507 acid backbone conformation depends on identity of the translated codon. Nat Commun.
508 2022;13:2815. (DOI: 10.1038/s41467-022-30390-9)
- 509
- 510 19. Nelson DL, Cox MM. Lehninger principles of biochemistry. 7th ed. New York: Freeman;
511 2017.
- 512
- 513 20. Anslyn EV, Dougherty DA. Modern physical organic chemistry. Sausalito CA: University
514 Science Books; 2006.
- 515
- 516 21. Olivares AO, Baker TA, Sauer RT. Mechanical protein unfolding and degradation. Annu
517 Rev Physiol. 2018;80: 413-429. (DOI: 10.1146/annurev-physiol-021317-121303)
- 518
- 519