

Original Research Article

Studies on genetic variability, divergence and characters associated with yield components in rice**Abstract**

The present investigation was conducted to estimate genetic parameters such as genotypic coefficient of variation (GCV), phenotypic coefficient of variation (PCV), heritability and genetic advance (GA) along with correlations, path coefficients and Genetic divergence from data collected on 22 rice genotypes. The results revealed highly significant mean squares due to genotypes for all traits studied, indicating the existence of sufficient variation among the genotypes and therefore an ample scope for effective selection. High phenotypic coefficient of variation (PCV) were observed for only seed yield per plant. Moderate phenotypic coefficient of variation and genotypic coefficient of variation (GCV) observed for number of tillers/plant, number of spikelets/panicle, flag leaf length, flag leaf width, number of filled grains/panicle, 1000-grain weight and biological yield. High heritability and genetic advance as per cent of mean was observed for number of tillers/plant, number of spikelets/panicle, flag leaf length, flag leaf width, number of filled grains/panicle, 1000-grain weight, biological yield and seed yield per plant, indicating the effectiveness of direct selection for improvement of these traits. Panicle length, 1000-grain weight and biological yield had recorded positive and significant association with seed yield per plant. The results on path analysis also revealed high and positive direct effect for biological yield followed by harvest index, days to 50% flowering, plant height, panicle length and number of spikelets/panicle and hence, these traits were identified as the most effective selection criteria for improvement of seed yield per plant in rice genotypes. The results on divergence analysis revealed that the genotypes were grouped into five clusters. Cluster I constituted maximum number of (10) genotypes. Maximum differences among the genotypes within the same cluster (intra-cluster) were shown by cluster IV (18.33) followed by cluster I, II, III. Cluster V showed zero intra-cluster distance. Cluster IV and V (20.20) showed maximum inter cluster distance, suggesting that the genotypes constituted in these clusters may be used as parents for future hybridization programme.

1. Introduction

Rice (*Oryza sativa* L.) is a plant belonging to the family of grasses (Gramineae) Poaceae. It is one of the most important cereal crops in the world and forms the staple diet of about 50% of the world's population. Asia Pacific Region is the leader in rice production accounting for about 90% of the world's production. Over 75% of the world supply is consumed by people in Asian countries and thus rice is of immense importance to the food security of Asia. The demand for rice is expected to increase with the continuous increase in the global population.

India is having a vast and long history of rice cultivation. Globally, India stands first in rice cultivation area and second in rice production, after China. It contributes 30% of global rice production. Within the country, rice occupies one-quarter of the total cropped area, which contributes about 40 to 43% of total food grain production and continues to play a vital role in the national food and livelihood security system. In India, rice is cultivated over an area of about 45.0 million hectares. About 40% of the area in India is rainfed which 23% of the rainfed rice area is upland and 77% is lowland. The Rice is grown in a vast diverse geographical and climatic

conditions ranging from below sea level areas like Kuttanad in Kerala to ^{the high} high altitude valley of Kashmir.

In India Rice is grown in 43.78 million ha, the production level is 118.43 million tonnes, and the productivity is about 20705 kg/ha during 2019-2020 (Agriculture statistics at a glance-2020). India occupies the first position in rice under the area of cultivation and second position in production which almost tripled its production during the second half of the last century. Global demand for rice is projected to grow at least equal to population growth, thus requiring a 70% (765 mt) increase in the supply of rice by the year 2025. To meet the demand for rice as the population increases the future demand for rice also increases, we have to work on the increasement of production to match the rate of increase in population growth. India is one of the leading exporters of rice, particularly basmati rice. *Oryza sativa* has many ecotypes or cultivars adapted to various environmental conditions.

A critical analysis of the genetic variability is a prerequisite for initiating any crop improvement programme and for adopting appropriate selection techniques. Correlation is the measure of the mutual relationship between two variables. Further studies on correlations help the plant breeder to know what exactly and how one character brings improvement in another character. Path coefficient analysis is a standard regression coefficient and measures the direct influence of one variable upon the other. Direct selection for yield is not a reliable approach since it is influenced by the environment. Therefore, it is essential to identify the component characters through which yield can be improved. The use of the correlation coefficient is to establish the extent of association between yield and yield components and some other character, which are having a decisive role in influencing the yield.

Earlier workers considered distances in place of origin as an index of genetic diversity and used it for the selection of parents for hybridization programmes. However, the genetic diversity of the selected parents has not been found to be based on factors such as geographic diversity/place of release or ploidy level (Singh et al., 2018).

The present study was carried out with the objectives of estimating the genetic variability, correlation, path analysis and genetic divergence.

2. Materials and methods

The experimental material for the present study comprised of 22 genotypes of rice laid in randomized block design (RBD) with three replications at Research farm, Department of Genetics and Plant Breeding, School of Agriculture, Lovely Professional University, Phagwara, Punjab., during kharif season, 2021. Standard agronomic practices and plant protection measures were taken as per schedule. Observations were recorded on five randomly selected plants per replication for days to 50% flowering, plant height (cm), panicle length (cm), number of tillers per plant, number of spikelets/panicle, flag leaf length (cm), flag leaf width (cm), number of filled grains/panicle, days to maturity, 1000-grain weight (gm), biological yield (gm), seed yield per plant (gm), harvest index (%) on plot basis. The data ^{were} subjected to Analysis of variance Panse and Sukhatme 1978, Heritability Hanson et. al., 1956, Genetic advance Johnson et. al., 1955, Correlation coefficient analysis Miller et al., 1958, Path coefficient analysis Dewey and Lu 1959, Genetic diversity using D2 analysis Mahalanobis 1936.

3. Results and discussion

Analysis of variance (ANOVA) indicated that the mean sum of square (MSS) due to genotypes were highly significant for all the characters at 1% and 5% level of significance and it indicated the presence of ^{the consisted} amount of genetic variability for all the traits (Table 1). As a result, there is a lot of scope for effective selection. Table 2 shows the results for yield and yield component characteristics in terms of mean, range, phenotypic coefficient of variation (PCV), genotypic coefficient of variation (GCV), heritability, and genetic advance as a percentage of ^{the mean} mean. A perusal of these results revealed ^{a maximum} maximum range of variability for the trait number of spikelets/panicle (112.53-242.03) while ^{a minimum} minimum range (0.9-1.56) was recorded for flag leaf width. In the present study, the phenotypic coefficient of variation was greater than genotypic coefficient of variation for all the traits. It means ^{the environment} the observed variation is not only due to the genotypes, but also due to suitable influence of environment. PCV and GCV were estimates classified into low (below 10%); medium (10-20%) and high (above 20%) in order to draw conclusions about these parameters. In the present study, there is only high PCV for and moderate GCV for seed yield per plant, indicating the existence of ^{a high} high amount of variability for effective selection towards ^{the improvement} the improvement of these traits. However, the magnitude of PCV and GCV were moderate for the number of tillers per plant, number spikelets per panicle, 1000 grain weight, biological yield, flag leaf length, flag leaf width, plant height and harvest index. Similar kind of results were found by Sumanth *et al.*, 2017 for number of tillers/plant, number of spikelets/panicle and flag leaf length, Keerthiraj & Biju 2020 flag leaf width and 1000-grain weight and Srujana *et al.*, 2017 for biological yield. Low PCV and low GCV were observed for traits viz., days to 50% flowering, panicle length and days to maturity, however indicating little scope for improvement of these traits through selection. Similar kind of results found by Singh *et al.*, 2021 for days to 50% flowering and panicle length, Lipi *et al.*, 2020 for days to maturity. Heritability estimates for the various traits studied ranged from 60.25 (harvest index) to 98.09 (flag leaf length). High estimates of heritability (> 60%) were recorded for all the 13 traits studied. Similar kind of results found by ^{were found} Sumanth *et al.*, 2017, Srujana *et al.*, 2017. A perusal of the results on genetic advance as percent of mean revealed high values (>20%) number of tillers/plant, number of spikelets/panicle, flag leaf length, flag leaf width, number of filled grains/panicle, 1000-grain weight, biological yield and seed yield per plant. It indicates that ^{transferred} maximum genetic gain can be received by using these characters and these characters are transfer from parent to their offspring. Similar kind of results found by Sandeep *et al.*, 2018 for number of tillers/plant and 1000-grain weight, Singh & Verma 2018 for number of spikelets/panicle, biological yield, Keerthiraj & Biju 2020 for flag leaf length and flag leaf width. Moderate genetic advance as percentage of (10-20%) mean was recorded for days to 50% flowering, plant height and harvest index. It indicates partially genetic gain by selection. Similar kind of results were found by Sandeep *et al.*, 2018 for days to 50% flowering, Kishore *et al.*, 2015 for plant height.

High heritability coupled with high genetic advance as per cent of mean was recorded for number of tillers/plant, number of spikelets/panicle, flag leaf length, flag leaf width, number of filled grains/panicle, 1000-grain weight, biological yield and seed yield per plant. Indicating that heritability observed was due to additive gene effects and therefore selection would be effective in ^{the improvement} improvement of these traits. High heritability coupled with moderate genetic advance observed in traits viz., days to 50% flowering and plant height. while moderate heritability with moderate genetic advance was observed only for harvest index. Low heritability with low genetic advance was not observed in any character. In the present study traits like number of tillers/plant, number of spikelets/panicle, flag leaf length, flag leaf width, number of filled grains/panicle, 1000-grain weight, biological yield and seed yield per plant ^{indicate} indicating the preponderance of additive gene action and therefore, ^{the scope} scope for effective improvement of these traits through selection.

The correlation analysis among the thirteen characters studied is presented in Table 3. In the

present findings, the magnitudes of genotypic correlation coefficient were greater than corresponding phenotypic correlation coefficient in general. In genotypic correlation coefficient Seed yield per plant exhibited positive and high significant correlation with the characters namely panicle length, flag leaf length, 1000-grain weight, and biological yield. While harvest index had showed positive and significance correlation for seed yield per plant. For days to 50% flowering, plant height, number of tillers/plant, number of spikelets/panicle, flag leaf width, number of filled grains/panicle and days to maturity where significance had not shown for the negative correlation with seed yield per plant. Similar findings were found with seed yield per plant exhibit for 1000-grain weight Shrestha *et al.*, 2018; Rashid *et al.*, 2014, similar findings for panicle length by Rashid *et al.*, 2014 and for harvest index by Moosavi *et al.*, 2015; Idris *et al.*, 2012. In phenotypic correlation coefficient seed yield per plant exhibited positive and high significant correlation with the characters namely 1000-grain weight, panicle length and biological yield. while harvesting index are shown positive and significant only at 5 % level and Flag leaf length had shown negative and significant correlation with the seed yield. Days to 50% flowering, plant height, no of tillers per plant, no of spikelets per panicle , flag leaf width , number of grain per panicle , days to maturity. Similar findings were found by Idris *et al.*, 2012. The results revealed that there is scope for simultaneous improvement of these traits through selection.

With the help of path coefficient analysis, the correlation coefficient is partitioned into direct and indirect effects (Table 4). The highest positive direct effect on seed yield per plant was exhibited by biological yield followed by harvest index, days to 50% flowering, plant height, panicle length and number of spikelets/panicle. Negative direct effect on seed yield per plant was exhibited by panicle length, flag leaf length followed by 1000-grain weight, number of filled grains per panicle, and days to maturity. The highest positive indirect effect on seed yield per plant was exhibited by biological yield, harvest index, 1000-grain weight, panicle length, followed by flag leaf width had shown indirect negative effect on seed yield, number of filled grains/panicle, days to 50% flowering, flag leaf length, number of spikelets/panicle, plant height, number of tillers per plant , days to maturity. Similar findings were found with Days to 50% flowering Ekka *et al.*, 2011, Plant height Gour *et al.*, 2017, Panicle length and number of filled grains/panicle Moosavi *et al.*, 2015; Idris *et al.*, 2012; Ekka *et al.*, number of spikelets/panicle and flag leaf width Jeke *et al.*, 2021, days to maturity and harvest index Gour *et al.*, 2017; Moosavi *et al.*, 2015, number of tillers/plant Shrestha *et al.*, 2018, Laxmi *et al.*, 2014, flag leaf length Reddy *et al.*, 2013, 1000-grain weight Nirmaladevi *et al.*, 2015; Rashid *et al.*, 2014, biological yield Reddy *et al.*, 2013. Therefore, harvest index, days to 50% flowering, plant height, panicle length and number of spikelets/panicle are the traits which can be directed in selection programmes to obtain higher grain yield.

The twenty-two genotypes under study were grouped into five clusters using Mahalanobis D2 analysis (Table 5). Among five clusters, cluster I was the biggest with 10 genotypes followed by cluster II (4 genotype), cluster III (3 genotype), IV (4 genotype) and V (1 genotype). The discrimination of genotypes into discrete clusters suggested presence of high degree of genetic diversity in the material evaluated. Presence of substantial genetic diversity among the parental material screened in the present study indicated that this material may serve as good source for selecting the diverse parents for hybridization programme. In conformity to the present investigation, similar findings were found by Koli *et al.*, 2022; Ranjith *et al.*, 2018.

The inter and intra average distances among nine clusters were computed and have been given in Table 6. Divergence analysis exhibited were shown by cluster IV (12.68) followed by cluster II (11.28), followed by cluster I (10.10), followed by cluster III (7.91), followed by cluster V showed zero intracluster distance. Cluster IV and V (20.45) showed maximum inter cluster distance followed by that between cluster III and V (20.20), cluster III and IV (18.29), cluster III and II (16.64), cluster II and V (16.60), cluster I and V (15.88), cluster II and IV (14.66), cluster I and IV (14.58), cluster I and II (14.48) and cluster I and III (13.32), cluster I and III (13.32). To

realize much variability and high heterotic effect, Mishra et al. (2003) and Chaturvedi and Maurya (2005) recommended that parents should be selected from two clusters having wider inter cluster distance.

The cluster mean values showed a wide range of variations for all the characters undertaken in the study (Table 7). The maximum cluster mean for days to 50% flowering in cluster V, plant height in cluster III, panicle length in cluster IV, number of tillers/plant in cluster III, number of spikelets/panicle in cluster III, flag leaf length in cluster III, flag leaf width in cluster III, number of filled grains/panicle in cluster I, days to maturity in cluster V, 1000-grain weight in cluster IV, biological yield in cluster II, seed yield per plant in cluster IV, harvest index in cluster III. The above observation confirms wide variation from one cluster to another in respect of cluster mean, which indicated that genotypes having distinct different mean performance for various characters were separated into different clusters. The five clusters in the aforesaid divergence analysis contained frequently the genotypes of heterogeneous origin. Although the genotype originated in ^{the same} same place or geographic region were also found to be grouped together in same cluster. The instances of grouping of genotypes of different origin or geographical region also in same cluster were observed in case of all the five clusters. This suggests ^{a lack} lack of parallelism between genetic and geographic diversity. Therefore, the selection of parental material for hybridization programme simply based on geographic diversity may not be a successful exercise for the choice of suitable divergent parents. Selection based on genetic divergence analysis would be more rewarding than the choice made based on geographic diversity.

The contribution of each trait to total divergence is presented in table 8. Among the traits studied, flag leaf length contributed maximum divergence (29.00%) followed by ^{the number} number of spikelets/panicle (25.97%), flag leaf width (11.26%) and biological yield (12.55%). The minimum percentage of contribution was observed in 1000-grain weight (7.36%), number of filled grains/panicle (6.06%), panicle length (5.63%) followed by days to 50% flowering (0.43%), days to maturity (1.43%), days to maturity (0.87%), seed yield per plant (0.87%) and plant height, number of tillers/plant, harvest index contributed zero percent divergence. The traits *viz* flag leaf length contributed maximum divergence followed by number of spikelets/panicle, flag leaf width and biological yield contributed 78.78 per cent towards total divergence. Hence, these characters should be given importance during hybridization and selection in the segregating population.

4. Conclusion

The findings indicate that the genotypes studied have adequate genetic variability. Characters *viz.*, panicle length, flag leaf length, 1000-grain weight, biological yield and harvest index. Path analysis revealed ^{a high} high and positive direct effect for biological yield followed by harvest index, days to 50% flowering, plant height, panicle length and ^{the number} number of spikelets/panicle and hence, these traits were identified as the most effective selection criteria for improvement of seed yield per plant in rice genotypes. Genetic diversity analysis was carried out through D² analysis in order to assess the genetic divergence among genotypes under study. Based on D² ^{the maximum} analysis twenty-two rice genotypes were grouped into five clusters. Cluster I constituted maximum number of genotypes. Further, the results ^{of} on divergence analysis revealed that Cluster I constituted maximum number of genotypes. Maximum differences among the genotypes within the same cluster (intra-cluster) were shown by cluster I followed by ^{clusters} cluster II, III, IV. Where cluster V showed zero intra-cluster distance. ^{Clusters} Cluster III and V showed maximum inter cluster distance, suggesting that the genotypes constituted in these clusters may be used as parents for future hybridization programme.

Table 1: Analysis of variance for seed yield and yield components in rice

S. No.	Characters	Source of variation			CV
		Replications	Treatments	Error	
		2	20	40	
1	Days to 50% flowering	19.69	108.71	13.34	3.77
2	Plant height (cm)	485.54	511.49	107.40	8.71
3	Panicle length (cm)	2.58	6.20	1.32	6.79
4	Number of tillers/plant	0.31	16.70	0.72	4.91
5	Number of spikelets/panicle	114.56	3539.45	68.20	4.37
6	Flag leaf length (cm)	0.54	75.74	1.44	3.75
7	Flag leaf width (cm)	0.001	0.09	0.004	5.50
8	Number of filled grains/panicle	72.57	2254.78	155.64	6.89
9	Days to maturity	4.80	100.48	8.23	2.20
10	1000 Grain weight (gm)	1.06	28.53	1.18	8.81
11	Biological yield (gm)	33.98	236.13	21.23	6.17
12	Seed yield per plant	19.37	76.25	4.63	8.86
13	Harvest index	51.15	71.75	28.519	11.31

Table 2.: Genetic parameters of variability for yield and yield attributing characters in Rice

S. No.	Characters	Range		Mean	Coefficient of variation		Heritability in broad sense (%)	Genetic advance in per cent of mean
		min	max		GCV	PCV		
1	Days to 50% flowering	87.33	106.33	96.87	5.81	6.21	87.72	11.22
2	Plant height (cm)	95.62	147.43	118.95	9.75	10.97	79.00	17.86
3	Panicle length (cm)	8.30	16.36	12.55	5.43	6.13	78.58	9.92
4	Number of tillers/plants	20.44	26.16	23.46	18.37	18.37	95.65	37.02
5	Number of spikelets/panicles	112.53	242.03	188.83	18.01	18.19	98.07	36.74
6	Flag leaf length (cm)	22.52	41.99	32.03	15.53	15.68	98.09	31.68
7	Flag leaf width (cm)	0.9	1.56	1.17	14.62	14.96	0.9549	29.44
8	Number of filled grains/panicle	131.95	233.50	180.83	14.62	15.16	93.10	29.07
9	Days to maturity	119.86	141.06	130.18	4.25	4.44	91.80	8.40
10	1000 Grain weight (gm)	16.47	34.45	24.43	17.13	17.50	95.86	34.56
11	Biological yield (gm)	12.62	24.69	17.61	16.27	17.06	91.01	31.99
12	Seed yield per plant	31.86	64.86	51.98	19.99	20.63	93.92	39.91
13	Harvest index	36.03	56.72	47.17	8.04	10.36	60.25	12.86

Table.3 Genotypic and Phenotypic correlation among 13 different characters in 21 rice genotypes

S. N	Characters	Days to 50% flowering	Plant height (cm)	Panicle length (cm)	Number of tillers/plants	Number of spikelets/panicle	Flag leaf length (cm)	Flag leaf width (cm)	Number of filled grains/panicle	Days to maturity	1000 grain weight	Biological yield	Harvest index	Seed yield per plant
1	Days to 50% flowering	1.0000	-0.1809	-0.2387	-0.2976	-0.1058	-0.0973	0.1639	0.0084	0.6019**	-0.0148	-0.0668	-0.2460	-0.1788
		1.0000	-0.1972	-0.2748	-0.3633	-0.1197	-0.0929	0.1824	0.0172	0.6601**	-0.0328	-0.0875	-0.2864	-0.1782
2	Plant height (cm)		1.0000	-0.1481	0.4831*	0.2676	0.7574**	-0.1497	-0.2170	0.3499	-0.1380	-0.3290	0.1681	-0.2276
			1.0000	-0.1621	0.5362**	0.3172	0.8528**	-0.1863	-0.2312	0.3879	-0.1749	-0.3701	0.2879	-0.2464
3	Panicle length (cm)			1.0000	-0.5596**	-0.0321	-0.4409*	-0.3224	-0.4688*	-0.4540*	0.3297	0.5978**	0.1267	0.5785**
				1.0000	-0.6476**	-0.0343	-0.4551*	-0.3325	-0.4855*	-0.4834*	0.3446	0.6323**	0.2124	0.6223**
4	Number of tillers/plant				1.0000	0.0900	0.6335**	0.1672	0.1434	0.1229	0.1065	-0.4133	0.2367	-0.2479
					1.0000	0.1119	0.7253**	0.1719	0.1757	0.1474	0.1152	-0.4904*	0.2496	-0.3199
5	Number of spikelets/panicle					1.0000	0.2769	0.1652	0.2302	0.1341	-0.2926	-0.1153	0.0411	-0.0779
						1.0000	0.2835	0.1797	0.2475	0.1377	-0.3000	-0.1225	0.0692	-0.0755
6	Flag leaf length(cm)						1.0000	0.0634	-0.0174	0.4176	-0.1169	-0.6565**	0.1875	-0.5082*
							1.0000	0.0691	-0.0203	0.4345*	-0.1135	-0.6907**	0.2446	-0.5263**
7	Flag leaf width (cm)							1.0000	0.0588	0.1973	0.0405	-0.3359	0.2871	-0.1504
								1.0000	0.0671	0.1987	0.0407	-0.3470	0.3324	-0.1667
8	Number of filled grains/panicle								1.0000	0.0714	-0.1444	-0.1143	0.0534	-0.0423
									1.0000	0.0994	-0.1576	-0.1451	0.0702	-0.0618

9	Days to maturity										1.0000	0.0436	-0.2795	-0.2131	-0.3639
											1.0000	0.0313	-0.3170	-0.2494	-0.3887
10	1000 grain weight										1.0000	0.4457*	0.3810	0.5738**	
											1.0000	0.4710*	0.5074*	0.6034**	
11	Biological Yield										1.0000	-0.0215	0.8603**		
											1.0000	0.1144	0.9135**		
12	Harvesting index										1.0000	0.4853*			
											1.0000	0.5089*			

Table 4 Genotypic Direct and indirect effect of different characters on seed yield per plant in Rice

R SQUARE = 1.0024 RESIDUAL EFFECT =SQRT(1- 1.0024)

S. N	Characters	Days to 50% flowering	Plant height (cm)	Panicle length (cm)	Number of tillers/plant	Number of spikelets/panicle	Flag leaf length (cm)	Flag leaf width (cm)	Number of filled grains/panicle	Days to maturity	1000 grain weight	Biological yield	Harvest index	Seed yield per plant
1	Days to 50% flowering	0.0461	-0.0091	-0.0127	-0.0167	-0.0055	-0.0043	0.0084	0.0008	0.0304	-0.0015	-0.0040	-0.0132	-0.1782
2	Plant height (cm)	-0.0290	0.1472	-0.0239	0.0790	0.0467	0.1256	-0.0274	-0.0340	0.0571	-0.0257	-0.0545	0.0424	-0.2464
3	Panicle length (cm)	0.0403	0.0238	-0.1468	0.0951	0.0050	0.0668	0.0488	0.0713	0.0710	-0.0506	-0.0928	-0.0312	0.6223
4	Number of tillers/plant	0.0221	-0.0327	0.0394	-0.0609	-0.0068	-0.0442	-0.0105	-0.0107	-0.0090	-0.0070	0.0299	-0.0152	-0.3199
5	Number of spikelets/panicle	-0.0097	0.0256	-0.0028	0.0090	0.0807	0.0229	0.0145	0.0200	0.0111	-0.0242	-0.0099	0.0056	-0.0755
6	Flag leaf length(cm)	0.0219	-0.2012	0.1074	-0.1712	-0.0669	-0.2360	-0.0163	0.0048	-0.1025	0.0268	0.1630	-0.0577	-0.5263
7	Flag leaf width (cm)	-0.0109	0.0112	0.0199	-0.0103	-0.0108	-0.0041	-0.0599	-0.0040	-0.0119	-0.0024	0.0208	-0.0199	-0.1667
8	Number of filled grains/panicle	0.0000	0.0004	0.0009	-0.0003	-0.0005	0.0000	-0.0001	-0.0019	-0.0002	0.0003	0.0003	-0.0001	-0.0618

9	Days to maturity	-0.0733	-0.0431	0.0537	-0.0164	-0.0153	-0.0482	-0.0221	-0.0110	-0.1110	-0.0035	0.0352	0.0277	-0.3887
10	1000 grain weight	-0.0048	-0.0257	0.0506	0.0169	-0.0440	-0.0167	0.0060	-0.0231	0.0046	0.1467	0.0691	0.0745	0.6034
11	Biological Yield	-0.0620	-0.2624	0.4483	-0.3477	-0.0868	-0.4897	-0.2460	-0.1029	-0.2248	0.3340	0.7090	0.0811	0.9135
12	Harvesting index	-0.1189	0.1195	0.0882	0.1036	0.0287	0.1015	0.1380	0.0291	-0.1035	0.2106	0.0475	0.4150	0.5089

Table 5: Clustering pattern of 21 Rice genotypes & based on Mahalanobis's D^2 statistic

Cluster No.	No. of Genotypes	Genotypes
Cluster 1	10	RNR2458, RNR15048, JYOTHI, JGL11727, EXP092, NLR34449, NLR3049, CHECK -PR126, WGL32100, BPT5204
Cluster 2	4	KPS2879, AMAN, MTU1010, MTU1153
Cluster 3	3	DAMINI, PLA1100, JAYSRI
Cluster 4	4	JGL24423, ADT39, PLA1100, ADT37
Cluster 5	1	RNR18833

Table 6: Estimates of average intra and inter-cluster distances for the five clusters in rice.

Clusters	Cluster 1	Cluster 2	Cluster 3	Cluster 4	Cluster 5
Cluster 1	10.10	14.48	13.32	14.58	15.88
Cluster 2		11.28	16.64	14.66	16.60
Cluster 3			7.91	18.29	20.20
Cluster 4				13.32	20.45
Cluster 5					0.00

Table 7: Cluster means for different characters in Rice

Characters	Days to 50% flowering	Plant height (cm)	Panicle length (cm)	Number of tillers/plant	Number of spikelet/p anicle	Flag leaf length (cm)	Flag leaf width (cm)	Number of filled grains/pani cle	Days to maturity	1000 Grain weight (gm)	Biologica l Yield (gm)	seed yield per plant	Harvestin g index
Cluster 1	98.27	118.83	11.92	23.20	201.41	31.66	1.13	191.55	130.78	22.09	15.60	50.50	43.89
Cluster 2	92.00	120.42	13.66	24.26	159.44	33.28	1.00	168.76	127.45	27.42	21.17	58.16	47.52
Cluster 3	93.22	134.26	11.24	24.96	209.60	40.10	1.40	177.67	132.40	20.06	16.98	38.17	52.61
Cluster 4	99.00	106.74	15.07	22.21	190.26	25.76	1.29	169.45	127.05	31.70	19.85	61.79	51.48
Cluster 5	105.00	117.33	8.33	23.50	112.57	31.80	1.23	177.10	141.07	20.03	16.53	44.50	45.23

Table 8: Contribution of 13 traits of Rice towards divergence

S.No.	Source	Times Ranked 1st	Contribution %
1	Days to 50% Flowering	1	0.43 %
2	Plant height (cm)	0	0.00 %
3	Panicle length (cm)	13	5.63 %
4	Number of tillers/plants	0	0.00 %
5	Number of spikelets/panicles	60	25.97 %
6	Flag leaf length (cm)	67	29.00 %
7	Flag leaf width (cm)	26	11.26 %
8	Number of filled grains/panicle	14	6.06 %
9	Days to maturity	2	0.87 %
10	1000 Grain weight (gm)	17	7.36 %
11	Biological Yield (gm)	29	12.55 %
12	Seed yield per plant	2	0.87 %
13	Harvesting index	0	0.00 %

References

1. Agriculture statistics.2020. Pocket book on Agricultural statistics, ministry of Agricultural Government of India.
2. Dewey, D. R. and Lu. K. H. (1959). Correlation and path coefficient analysis of crested wheat grass seed production, *Agrion. J.*, 51: 515-518.
3. Ekka, R. E., Sarawgi, A. K., & Kanwar, R. R. (2011). Correlation and path analysis in traditional rice accessions of Chhattisgarh. *Journal of rice research*, 4(1), 11-18.
4. Gour, L., Maurya, S. B., Koutu, G. K., Singh, S. K., Shukla, S. S., & Mishra, D. K. (2017). Characterization of rice (*Oryza sativa* L.) genotypes using principal component analysis including scree plot & rotated component matrix. *International Journal of Chemical Studies*, 5(4), 975-83.
5. Hanson, C.H., H.F. Robinson and R.E. Comstock. 1956. Biometrical studies on yield in segregating population of Korean lespedesa. *Agron. J.* 48:268-272.
6. Idris, A. E., Justin, F. J., Dagash, Y. M. I., & Abuali, A. I. (2012). Genetic variability and inter relationship between yield and yield components in some rice genotypes. *American Journal of Experimental Agriculture*, 2(2), 233.
7. Johnson HW, HF Robinson and RE Comstock. 1955. Estimates of genetic and environmental variability in soybeans. *Agronomy Journal* 47: 314-318
8. Jeke, E., Mzengeza, T., Kyung-HO, K., & Imani, C. (2021). Correlation and Path Coefficient Analysis of Yield and Component Traits of KAFACI doubled haploid Rice (*Oryza sativa* L) Genotypes in Malawi. *Int J Agriculture Technology*, 1(2), 1-9.
9. Keerthiraj, B., & Biju, S. (2020). Genetic variability, heritability and genetic advance of yield and lodging-related traits in rice (*Oryza sativa* L.). *Electronic Journal of Plant Breeding*, 11(04), 1093-1098.
10. Kishore, N. S., Srinivas, T., Nagabhushanam, U., Pallavi, M., & Sameera, S. K. (2015). Genetic variability, correlation and path analysis for yield and yield components in promising rice (*Oryza sativa* L.) genotypes. *SAARC Journal of Agriculture*, 13(1), 99-108.
11. Koli, G. K., Upadhyay, S., Rathi, S. R., Gupta, S., Kumawat, G., Singh, P. K., & Sinha, B. (2022). Study of genetic divergence for yield and yield attributing traits of wild germplasm in rice. *The Pharma Innovation Journal*; 11(2): 472-474
12. Lipi, L. F., Hasan, M. J., Akter, A., Quddus, M. R., Biswas, P. L., Ansari, A., & Akter, S. (2020). Genetic variation, heritability and genetic advance in some promising rice hybrids. *SAARC Journal of Agriculture*, 18(2), 39-49.
13. Lakshmi, V. I., Sreedhar, M., Vanisri, S., Anantha, M. S., Rao, L. S., & Gireesh, C. (2019). Multivariate analysis and selection criteria for identification of African rice (*Oryza glaberrima*) for genetic improvement of indica rice cultivars. *Plant Genetic Resources*, 17(6), 499-505.
14. Miller PJ, Willianis JC, Robinson HF and Comostock RE 1958. Estimates of genotypic and environmental variance and covariance in upland cotton and their implication in selection. *Agron. J.* 50: 126-131
15. Mahalanobis, P. C. (1936). On the generalized distance in statistics", proceeding of the national institute of sciences of India. 2 (1), 49-55.
16. Moosavi, M., Ranjbar, G., Zarrini, H. N., & Gilani, A. (2015, January). Correlation between morphological and physiological traits and path analysis of grain yield in rice genotypes under Khuzestan conditions. *In Biological Forum* (Vol. 7, No. 1, p. 43). Research Trend.
17. Nirmaladevi, G., Padmavathi, G., Kota, S., & Babu, V. R. (2015). Genetic variability, heritability and correlation coefficients of grain quality characters in rice (*Oryza sativa* L.). *SABRAO journal of Breeding and Genetics*, 47(4), 424-433.

It does not appear in the text.
Lakshmi

18. Panse VG, Sukhatme PV. *Statistical Methods for Agricultural Workers*. ICAR, New Delhi 1978, 103-108
19. Rashid, K. A. S. H. I. F., Kahliq, I., Farooq, M. O., & Ahsan, M. Z. (2014). Correlation and cluster analysis of some yield and yield related traits in Rice (*Oryza sativa*). *Journal of Recent Advances in Agriculture*, 2(6), 271-276.
20. Reddy, G. E., Suresh, B. G., Sravan, T., & Reddy, P. A. (2013). Interrelationship and cause-effect analysis of rice genotypes in north east plain zone. *The Bioscan*, 8(4), 1141-1144.
21. Ranjith, P., Sahu, S., Dash, S. K., Bastia, D. N., & Pradhan, B. D. (2018). Genetic diversity studies in Rice (*Oryza sativa* L.). *Journal of Pharmacognosy and Phytochemistry*, 7(2), 2529-2531.)
22. Singh, Y., Singh, P. K., Pathak, V. N., Misra, V., & Mall, A. K. (2018) genetic divergence among rice genotypes studied through non-hierarchical euclidean cluster analysis.
23. Sumanth, V., Suresh, B. G., Ram, B. J., & Srujana, G. (2017). Estimation of genetic variability, heritability and genetic advance for grain yield components in rice (*Oryza sativa* L.). *Journal of Pharmacognosy and Phytochemistry*, 6(4), 1437-1439.
24. Srujana, G., Suresh, B. G., Lavanya, G. R., Ram, B. J., & Sumanth, V. (2017). Studies on genetic variability, heritability and genetic advance for yield and quality components in rice (*Oryza sativa* L.). *Journal of pharmacognosy and phytochemistry*, 6(4), 564-566.
25. Singh, B., Gauraha, D., Sao, A., & Nair, S. K. (2021). Assessment of genetic variability, heritability and genetic advance for yield and quality traits in advanced breeding lines of rice (*Oryza sativa* L.). *The Pharma Innovation*, 10(8), 1627-1630.
26. Sandeep, S., Sujatha, M., Subbarao, L. V., & Neeraja, C. N. (2018). Genetic variability, heritability and genetic advance studies in rice (*Oryza sativa* L.). *International Journal of Current Microbiology and Applied Science*, 7(12), 3719-3727.
27. Sumanth, V., Suresh, B. G., Ram, B. J., & Srujana, G. (2017). Estimation of genetic variability, heritability and genetic advance for grain yield components in rice (*Oryza sativa* L.). *Journal of Pharmacognosy and Phytochemistry*, 6(4), 1437-1439.
28. Singh, N., & Verma, O. (2018). Genetic variability, heritability and genetic advance in rice (*Oryza sativa* L.) under salt stressed soil. *Journal of Pharmacognosy and Phytochemistry*, 7(3), 3114-3117.
29. Shrestha, N., Poudel, A., Sharma, S., Parajuli, A., Budhathoki, S., & Shrestha, K. (2018). Correlation Coefficient and Path Analysis of Advance Rice Genotypes in Central Mid-hills of Nepal. *International Journal of Research in Agricultural Sciences*, 5(3), 2348-3997.