

Assessment of Genetic Variability, Heritability and Genetic Advance in Garden pea (*Pisum sativum* L.)

Comment [WU1]: Under what agroclimatic condition???

Comment [WU2R1]: Name of country??

Comment [WU3R2]:

Abstract

Genetic variability estimation was conducted in ten genotypes of peas during 2021-22. Analysis variance was highly significant for all characters studied except plant spread. The highest genotypic and phenotypic co-efficient of variation was recorded for pod yield per plant (15.3 ,16.97) followed by seed yield per plot (13.91 ,18) and lowest was recorded for days to 1st flowering (3.56,3.63) high heritability with genetic advance as per cent of mean was recorded as yield per plant (81,28.4) and pod yield per plant (59,22.1) and lowest was recorded for plant height (0.6,1.72). this study indicates that emphasis should be given to these characters for yield improvement.

Comment [WU4]: In what geo agri condition???

Keywords: Genetic variability, GCV, PCV, Heritability, Genetic Advance, Pea

Comment [WU5]: Concluding remarks

Comment [WU6R5]: Recommendation???

Introduction

Field pea (*Pisum sativum* L.), a legume crop, belongs to the Leguminosae family and contains a high amount of protein including amino acids, especially lysine. Beans are considered to be the most nutritious part of the human diet because they contain 42.65% carbohydrates, 27.8% protein and iron (Fe), sodium (Na) phosphorus (P), potassium (K), and some other important elements. Field pea has multifaceted uses as food, feed and fodder. The residues provide nutritious food for cattle and dairy cows and thus provide an additional benefit to poor farming families. Field peas are cultivated mainly for green beans and seeds are used as vegetables. Mature seeds can be used to make 'dal' or 'chapati' and other value-added goods such as 'fried dal'. As a legume, the field has a unique ability to fix atmospheric nitrogen through symbiosis. It may have originated in South Asia but it is one of the most important herbaceous vegetables in Bangladesh. Field peas with an annual production of 1,35,34,166 tons are one of the most produced pulses crops in the world. Major pea producing countries include Canada, the Russian Federation, China, Ukraine, India, USA, France, Australia, Ethiopia and Germany. The success of the breeding program depends on the amount of genetic variability available for absorption and the extent to which the desirable characters are heritable.

Collection, preservation and evaluation of germplasm are key information for any breeder to initiate a breeding program for crop improvement. Heritability acts as a predictive tool in expressing the reliability of phenotypic traits and thus high heritability traits can aid in the effective selection of specific characters and create future breeding programs. The analysis of the relationship helps in evaluating the existing relationship between the yield and its components. The study of genetic variability and interrelationships is of great value in the selection of the preferred characters of field pea genotypes to increase seed yield and quality. Genetic diversity is available in multivariate analysis, multiple statistical tools such as Euclidean clustering and principal component analysis (PCA). Multivariate analysis is an important tool for estimating genetic divergence in the populations of several workers. The principal component analysis is an adaptive data analysis technique that is used to effectively visualize similarities and differences between genotypes and to identify the quantitative

characters that contribute the highest towards genetic divergence. To meet the present demand, there is an essential germplasm evaluation for the genetic improvement of pea to develop desired high yielding genotypes. Thus, the present study was conducted to identify higher genotypes based on genetic variability and association between yield and yield-related traits and to identify genetically different genotypes using PCA for further different breeding programs in the development of high yielding field pea varieties.

Materials and methods

The experiment was carried out at Crop Research Centre, School of Agriculture, ITM University, Gwalior, Madhya Pradesh. The experimental material comprised of ten genotypes which were collected from different source. The experiment was laid out in randomized block design (RBD) with three replications. The whole investigation was done under the scientific management practices. During study, observations for plant height (cm), plant spread, number of branches per plant, days to first flowering, days to 50 % flowering, days to first picking, number of pods per plant, number of seeds per pod, pod length (cm), pod width (cm), weight of 100 seeds, pod yield per plant (g), pod yield per plot, T.S.S and protein (Ali khan, 1973) were recorded on five randomly selected plants from each treatment. The genotypic and phenotypic coefficients of variation were calculated as per the method was suggested by Burton and Devane (1953). Heritability in broad sense and expected genetic advance on the basis of percent of mean were worked out according to the method advocated by Robinson *et al.*, (1953) and Johnson *et al.*, (1955), respectively.

Results and discussion

The Analysis of Variance (ANOVA) for different traits showed that mean sum of squares due to genotypes exhibit significant differences for most of the traits under observation at 1% probability level except for pod length which was significant at 5% probability level and plant height which was non significant at both levels of (Table 1). It shows large amount of genetic variability among pea genotypes.

Table: 1 Analysis of variance of 13 characters in garden pea genotypes

S. No.	Characters	Mean of sum square		
		Replication DF=2	Treatment DF=2	Error DF=13
1	Plant height	130.86	108.2	90.63
2	Plant spread	204.25	124.62	85.99
3	Days to first flowering	0.73	13.6**	0.18
4	Days to 50% flowering	0.16	21.192**	0.04
5	Number of branches per plant	0.6	0.35	0.45
6	Number of pods per plant	1.68	5.04	2.05
7	Number of seeds per pod	0.08	1.3**	0.13
8	Pod length	0.36	0.14	0.35
9	Pod width	0.25	0.695**	0.14
10	Days to 1 st picking	1.23	7.88	4.04
11	Weight of 100 seeds	0.3	23.8**	3.28
12	Pod Yield per plant	655.84	1044.55**	74.11
13	Pod yield per plot	0.181	0.269**	0.05

** Level of significance at 5%

Comment [WU7]: Aims & %objective of research??

Comment [WU8R7]: Background of research

Comment [WU9R8]: Problem statement???

Comment [WU10]: LSD test for comparison of means???

Comment [WU11]: Name of country???

Comment [WU12]: Very old reference

Comment [WU13]: Old reference

Comment [WU14]: Old reference

Comment [WU15]: need more discussion with comparison with past result

Mean of sum squares in ANOVA revealed high variability among 10 genotypes for all the characters (except number of branches per plant) at 5% level of probability (table1).

The mean performance of various genotypes has also showed good range of variability for various characters, which were studied in present investigation (table1) the range recorded for plant height (93.6-52.8), plant spread (100-59), days to 1st flowering (64.2-55.4), days to 50% flowering (70.2-60.2), number of branches per plant (7.6-5) number of pods per plant (23.6-16.6), number of seeds per pod (8.6-6), pod length (5.2-3.6), pod width (7.7-5.3) weight of 100 seeds (34-19), days to 1st harvest (79-70), yield per plant (160-72) and pod yield per plot (2.6-1.3). The characters under investigation were analysed for genotypic coefficient of variation (GCV), phenotypic coefficient of variation (PCV), heritability (broad sense) and genetic advance as percent of mean (Table 2). It was observed that magnitude of phenotypic coefficient of variation (PCV) were higher than genotypic coefficient of variation (GCV) for all the characters under study which is an indicator of additive effect of the environment on the expression of the trait. It is also observed that the low range between PCV and GCV so it reveals that these traits have low sensitivity to environmental effects and it is reducible. Similar findings were also reported by Katoch *et al.*, (2016), Siddika *et al.*, (2013), Kumar *et al.*, (2013) and Ahmad *et al.*, (2013).

Genotypic coefficient of variation (GCV) ranged from (1.54-15.3), higher magnitude of GCV was recorded pod yield per plant (15.3), pod yield per plot (13.91), weight of 100 seeds (8.95). similar findings were recorded by (Naveen Kumar Jaiswal, G. Roopa Lavanya) (Archi Gupta 2020). (D. Satish, Abdul Karim 2019) plant height (3.4), plant spread (4.65), days to 1st flowering (3.56), days to 50% flowering (4.2), number of branches per plant (2.94), number of pods per plant (4.8), number of seeds per pod (8.87), days to 1st harvesting (1.54), weight of 100 seeds (8.95) pod length (5.89), pod width (6.58), pod yield per plant (15.3), pod yield per plot (13.91). High values of GCV is an indication of high genetic variability among the genotypes and thus the scope for improvement of these characters through simple selection would be better.

Phenotypic coefficient of variation (PCV) ranged from (3.63-18) higher magnitude of PCV was recorded pod yield per plot (18), pod yield per plant (16.97), plant height (13.81) similar findings were recorded by (Naveen Kumar Jaiswal, G. Roopa Lavanya) (Archi Gupta 2020). (D. Satish, Abdul Karim 2019) plant height (13.81), plant spread (12.89), days to 1st flowering (3.63), days to 50% flowering (4.21), number of branches per plant (10.23) number of pods per plant (8.42), number of seeds per pod (10.28), pod length (11.78), pod width (8.8), weight of 100 seeds (10.93), days to 1st harvest (3.14), pod yield per plant (16.97), pod yield per plot (18).

Heritability (bs) ranges from (99-0.6) higher magnitude of heritability was recorded in days to 50% flowering (99) similar findings were recorded by (Hafiz Bashir Ahmad, Salsabeel Rauf, Ch. Muhammad Rafiq) estimates of heritability of characters plant height (60), plant spread (13), days to 1st flowering (96), days to 50% flowering (99), number of branches per plant (0.6), number of pods per plant (32), number of seeds per pod (74), pod length (25), pod width (55), days to 1st harvesting (24), weight of 100 seeds (67), pod yield per plant (81), pod yield per plot (59). Similar findings were recorded by hafiz Bashir Ahmad (2014) and Naveen Kumar Jaiswal (2014).

Genetic advance ranged from (33.4 to 0.43) estimates of genetic advance was recorded as plant height (1.22), plant spread (2.66), number of branches per plant (0.12) , days to 1st flowering (4.27), days to 50% flowering (5.45) , number of pods per plant (1.17),number of seeds per plant (1.11) , pod length (0.07), pod width (0.66) , days to 1st harvesting (1.14) , weight of 100 seeds (4.36),pod yield per plant (33.41) pod yield per plot (0.43). Similar findings were recorded by Archi Gupta (2013) and Suraj Lathura (2020)

Genetic advance as percentage of mean ranged from 1.72 to 28.4 for plant height, yield per plot the estimates of genetic advance as percentage of mean was recorded as plant height (1.72), plant spread (3.46), number of branches per plant (3.02), days to 1st flowering (7.19), days to 50%flowering (8.63), number of pods per plant (5.65), number of seeds per pod (15.76) weight of 100seeds (15.1), days to 1st picking (1.55), pod length (36.8), width (10.14), yield per plant (28.4), yield per plot (22.1). Similar findings were recorded by Archi Gupta (2013), AsfakunsiddikaA.K.M.Aminulislam (2013) and Suraj Lathura (2020).

Table 3: Estimates of genetic parameters for growth, earliness, yield and quality parameters in garden pea genotypes

Characters	Grand mean	Range min	Range max	GCV	PCV	Heritability %	Genetic advance	Genetic advance as % of mean
Plant height	71.12	93.6	52.8	3.4	13.81	60	1.22	1.72
Plant spread	77.09	100	59	4.65	12.89	13	2.66	3.46
No. of branches/plant	6.36	7.6	5	2.94	10.23	0.6	.012	3.02
Days to first flowering	59.38	55.4	64.2	3.56	3.63	96	4.27	7.19
Days to 50% flowering	63.14	70.2	60.2	4.2	4.21	99	5.45	8.63
Days to first picking	73.46	79	70	1.54	3.14	24	1.14	1.55
Weight of 100 seeds	28.9	34	19	8.95	10.93	67	4.36	15.11
No. of pods per plant	20.75	23.6	16.6	4.8	8.42	32	1.17	5.65
No. of seeds per pod	7.04	8.6	6	8.87	10.28	74	1.11	15.76
Pod length	4.52	5.2	3.6	5.89	11.78	0.6	0.07	36.8
Pod width	6.5	7.78	5.35	6.58	8.8	55	0.66	10.14
Pod yield per plant	117.49	160	72	15.3	16.97	81	33.41	28.44
Pod yield per plot	1.94	2.6	1.3	13.91	18.05	59	0.43	22.1




Fig. 1 Estimates of GCV and PCV for 13 characters in pea.




Fig. 2 Estimates of Heritability, Genetic Advance and Genetic Advance (%) mean for 13 characters in pea.

References

- A. Bhargava, F. Fuentes, S. Shukla, S. Srivastava, S. Dubey, y D. Ohri (2019) Genetic Variability in vegetable *Chenopodium* for morphological and quality traits over different cuttings. *J. Cien. Agr.* 46(2): 179-186, DOI: 10.7764/rcia.v46i2.2145.
- Afreen S, Singh AK, Moharana DP, Singh V, Singh P, Singh B. (2017) Genetic evaluation for yield and yield attributes in garden pea (*Pisum sativum* var. *hortense* L.) under North Indian gangetic plain conditions. *Int. J. Curr. Microbiol. App. Sci.* 2017; 6 (2):1399-1404.
- Ahmad HB, Rauf S, Rafiq M, Ullah Mohsin A, Iqbal A. (2014) Estimation of genetic variability in pea (*Pisum sativum* L.). *J. Glob. Innov. Agric. Soc. Sci.* 2014;2 (2):62-64.
- Ajay Kumar Verma (Author) (2014) Genetic variability, heritability and advances in Okra, Munich, *GRIN Verlag*, <https://www.grin.com/document/308061>.
- Archi Gupta, Bijendra Singh, Mukesh Kumar, Pooran Chand, Vaishali and Sharma, V. R. (2020) Studies on Genetic Variability, Heritability and Genetic Advance in Table Pea (*Pisum sativum* var. *hotense* L.). *Int.J.Curr.Microbiol.App.Sci.* 9(09): 3449-3454.
- Asfakun siddika A.K.M. Animul Islam, M.Golam Rasul, M.Abdul Kahleque Mian, Jalal Uddin Ahmed (2013) Genetic Variability in advanced generations of vegetable pea (*Pisum sativum* L.) *International journal of Plant Breeding* 2013. 7(2), 124-128.
- Azam MG, Iqbal MS, Hossain MA, Hossain J, Hossain MF (2020) Evaluation of Field pea (*Pisum sativum* L.) Genotypes based on Genetic Variation and association among Yield and

Comment [WU16]: Cofnclusiofn % & recomendatiofn are missing

Comment [WU17]: Refrefnces formate is fnot accordifng to the journal

Comment [WU18R17]: Please tally eacgh refrefnce witgh text

Yield Related Traits under High Ganges River Floodplain. *Int J Plant Biol Res* 8(2): 1120

Barkat Ali, Sujon Kumar, Waleed Ahmed (2021) Genetic Variability, Heritability and Correlation Coefficient in Production Traits of Pea (*Pisum sativum* L.) Genotypes. *International Journal of Genetics and Genomics*. Vol. 9, No. 4, 2021, pp. 78-88. doi: 10.11648/j.ijgg.20210904.13

Baskaran, V., Jayanthi, R., Janakiram, T., & Abirami, K. (2009). Studies on Genetic Variability, Heritability and Genetic Advance in Chrysanthemum. *Journal of Horticultural Sciences*, 4(2), 174-176. Retrieved from <https://jhs.ihr.res.in/index.php/jhs/article/view/540>

Borate, V. V., Dalvi, V. V. and Jadhav, B. B. (2010). Estimates of genetic variability and heritability. *Journal Maharashtra Agricultural University*, 35(1): 47-49.

Burton, G. W., and Devane, D. E. (1953). Estimating heritability in tall fescue (*Festuca arundinacea*) from replicated clonal material. *Agronomy Journal*, 45(10): 478-481.

Fisher, R. A., and Yates, F. (1938). Statistical tables: For biological, agricultural and medical research. Oliver and Boy

Hafiz Bashir ahmad (2014). Estimation of genetic variability in pea (*Pisum sativum* L.) *J.Glob.Innov.agric.soc.sci*.2014.2(2):62-64.

Kavya Eppakayala, Saidaiah Pidigam, Sivaraj Natarajan, Geetha Amarapalli and Ravinder Reddy Komati reddy (2020). Study of genetic variability and heritability and genetic advance and yield parameters in tomato (*Solanum lycopersicum* L.) germplasm. *Journal of Pharmacognosy and Phytochemistry* 2020. 10(1): 768-771.

Kumar B. (2018) Variability, heritability and genetic advance in pea (*Pisum sativum* L.). *International Journal of Plant Sciences*. 2018; 3(1):L211-212.

M.L. Meena, N. Kumar, J.K. Meena and T. Rai (2016) Genetic variability, Heritability and Genetic Advances in chilli, *capsicum Annuum*. Department of Applied Plant Sciences (Horticulture), Babasaheb Bhimrao Ambedkar Central University, Vidya Vihar Raebareli Road, Lucknow-226 025. *An open access international journal* P-ISSN: 0974-6455 E-ISSN: 2321-4007.

M.Shahbaz Akhtar, Yoko Oki, Tadashi Adachi and Md. H.R. Khan (2007) Analyses of the genetic parameters (variability, Heritability, Genetic Advances, Relationship of Yield and Yield Contributing characters) for some plant traits among *Brassica* Cultivars under Phosphorous starved environmental cues. *Journal of the faculty of environmental science and technology*, Okayama University Vol.12, No.1, pp.91-98, March 2007

Melaku AB, Mohamed W, Kumar V. (2020) Variability, heritability and genetic advance in indigenous and exotic okra [*Abelmoschus esculentus* (L.) *moench*] genotypes for yield and yield related traits at Dire Dawa, Eastern Ethiopia. *MOJ Eco Environ Sci*. 2020;5(4):164–169. DOI: 10.15406/mojes.2020.05.0018.

Melaku AB, Mohamed W, Kumar V. Variability, heritability and genetic advance in indigenous and exotic okra [*Abelmoschus esculentus* (L.) *moench*] genotypes for yield and yield related traits at Dire Dawa, Eastern Ethiopia. *MOJ Eco Environ Sci*.

2020;5(4):164–169. DOI: 10.15406/mojes.2020.05.00189.

Naveen kumarjaiswal, (2013) Genetic variability analysis in field pea (*Pisum sativum* L.) “*International journal of science and research (ISJR) ISSN(online):2319-7064.*

Netra Hari Ghimire, Paras Mani Mahat. (2019). Variability, Heritability and Genetic Advance of Advanced Breeding Lines of Barley (*Hordeum vulgare* L.) Under Mountain Environment of Nepal. *Int. J. Adv. Res. Biol. Sci.* 6(11): 34-42. DOI: <http://dx.doi.org/10.22192/ijarbs.2019.06.11.006>.

Pushpa Gudadinni, Vijay Bahadur, Priyanka Ligade, S.E. Topno and Prasad, V.M. (2017) Study on Genetic Variability, Heritability and Genetic Advance in Garden Pea (*Pisum sativum* var. *hortense* L.). *Int.J.Curr.Microbiol.App.Sci.*6(8):2384-2391. <https://doi.org/10.20546/ijcmas.2017.608.282>.

Suraj lathura (2020) study on genetic variability, heritability, and genetic advance in garden pea (*pisum sativum* var. *hortense* L.) *Journal of Pharmacognosy and Phytochemistry* 2020; 9(4): 2036-2039.

T.N.Vijaya Kumar T. B. Alloli, H. P. Hadimani, P. S. Ajjappalavar, D. Satish, Abdul Kareem and Hanchinamani, C. N. (2019) Studies on Genetic Variability, Heritability and Genetic Advance in Garden Pea (*Pisum sativum* L.) varieties. *Int.J.Curr.Microbiol.App.Sci.*8(12):3032-3038

Tejaswini.N, K. Ravinder Reddy, P. Saidaiah and T. Ramesh. (2017) Studies on variability, Heritability and Genetic Advance in Vegetable Amaranth (*Amaranthus tricolor* L.) genotypes. *Green Farming Vol. 8 (5) : 1071-1075 ; September-October*

Thakur, S., Thakur, R. and Mehta, K, D (2016). Genetic variability and association studies for green pod yield and component horticultural traits in garden pea under high hill dry temperate conditions of Tabo valley of Spiti district of Himachal Pradesh *International Journal of Science, Environment and Technology* 5(4):1987- 1992.

Tiwari, Garima, and G. Roopa Lavanya. (2012) Genetic variability, character association and component analysis in F4 generation of field pea (*Pisum sativum* var. *arvense* L.). *Karnataka Journal of Agricultural Sciences* 25.2 (2012).

Vinod Kumar (2021) Genetic variability, heritability and genetic advance for seed yield and related traits in diverse genotypes of pea (*Pisum sativum* L.) *The pharma innovation journal* 2021;10 (12)2972-2977.

Yücel, D. (2020). Genotypic and Phenotypic Variability for Yield and its Components in Normal and Late Sown Chickpea (*Cicer arietinum* L.). *Legume Research*, 43(1): 18-24.

Zerfu, A., Hailu, F. and Adal, M. (2021). Phenotypic Variability and Association (among yield components and) Yield Related Trait in desi Type Chickpea (*Cicer arietinum* L.) in Raya Kobo District, Northern Ethiopia. *J.Agrotechnology*, 10(1): 203.

UNDER PEER REVIEW

