

# Effect of moisture regimes and gypsum levels on growth and yield of *rab*Groundnut (*Arachis hypogea* L.)

## ABSTRACT

A Field experiment was conducted in groundnut at Krishivignan Kendra, Palem, Nagarkurnool during rab2021 to study the effect of moisture regimes and gypsum levels on growth and yield of *rab*groundnut (*Arachis hypogea* L.). Among three moisture regimes, I<sub>3</sub> (1.0 IW/CPE) recorded significantly higher values of growth parameters- Plant height (27.87cm), number of branches plant<sup>-1</sup> (14.20), leaf area 174.06cm<sup>2</sup> plant<sup>-1</sup>) and dry matter production (2711.67 kg ha<sup>-1</sup>) which is on par with I<sub>2</sub> (0.8 IW/CPE) of plant height, number of branches plant<sup>-1</sup>, leaf area at harvest and dry matter production at 90 DAS and at harvest and yield attributes pod yield (2337.68 kg ha<sup>-1</sup>), halum yield (1659.06 kg ha<sup>-1</sup>) compared to other treatments. In case of gypsum levels, gypsum @ 600 kg ha<sup>-1</sup> showed significantly higher values of growth parameters- plant height (28.93 cm), number of branches plant<sup>-1</sup> (15.85), leaf area (194.66 cm<sup>2</sup> plant<sup>-1</sup>) and dry matter production (3038.22 kg ha<sup>-1</sup>) which is on par with I<sub>2</sub> (0.8 IW/CPE) of plant height, number of branches plant<sup>-1</sup> at 60 DAS and yield attributes pod yield (2829.78 kg ha<sup>-1</sup>) and haulm yield ( 1752.22 kg ha<sup>-1</sup>) which is followed by I<sub>2</sub> (0.8 IW/CPE) and I<sub>1</sub>(0.6 IW/CPE). The interaction of moisture regimes and gypsum level I<sub>3</sub> (1.0 IW/CPE) and G<sub>4</sub> (600 kg ha<sup>-1</sup>) showed significantly higher values of dry matter production (1456.0 cm<sup>2</sup> plant<sup>-1</sup>) at 60 DAS, pod yield (3056.4 kg ha<sup>-1</sup>) and halum yield (1956.0 kg ha<sup>-1</sup>) over other treatments.

**Keywords:** Moisture regimes, Gypsum levels, groundnut

## 1. INTRODUCTION

Groundnut is an important oil and protein source to a large portion of the population in India. It is an annual, herbaceous legume and considered as king of vegetable oilseed crops in India and occupies a pre-eminent position in national edible oil economy. Groundnut seed contains 47-53 % oil, 26 % protein and 11.5 % starch. It is currently grown in an area of 21.8 Million hectares over the globe (2021). India, China and the United States of America are the leading producers since last 25 years and account for 70% of the area in the world. In India it is grown on an area of about 4.82 m ha with a total production of 9.95 m tonnes and productivity of 2.06 tonnes ha<sup>-1</sup> during 2019-20. In Telangana it occupies an area of 0.11 m ha with total production of 0.26 m tonnes and productivity of 2.39 tonnes ha<sup>-1</sup> in 2019-20 [1].

Comment [K1]: field

Comment [K2]: on

Comment [K3]: groundnuts

Comment [K4]: the

Comment [K5]: growing

Comment [K6]: million

Comment [K7]: throughout

Comment [K8]: have been

Comment [K9]: for the

Cultivation of groundnut under rainfed conditions and imbalanced nutrient management are the main reasons for low productivity of groundnut. Irrigation water, a crucial input in crop production is scarce and expensive. Resource efficient use of this input is essential which can be achieved through judicious water management practices. Adequate and timely supply of water is essential for higher yields. Keeping the total quantity of irrigation water constant, increasing the frequency of irrigation would maximize the yields in several crops. There is possibility to double the groundnut yields during *rabi* season over *kharif* with limited number of irrigations. Because of high productivity under assured irrigation, a climatological approach based on IW/CPE ratio (IW-irrigation water, CPE - cumulative pan evaporation) in irrigation scheduling has been found most appropriate as it integrates most of the weather parameters which determine the water requirement of a crop and increase production by at least 15 to 20 %. Scheduling of irrigation based on IW/CPE depends on both available water holding capacity of soil and climatic parameters.

Comment [K10]: groundnuts

Comment [K11]: groundnuts

Comment [K12]: ,

Comment [K13]: ,

Comment [K14]: of

Comment [K15]: a

Comment [K16]: the

Comment [K17]: a limited amount

Comment [K18]: irrigation

Comment [K19]: increases

To ensure increased yields of *rabi* groundnut in traditional areas of telanganait is necessary to have a thorough understanding of the changes in the soil-plant-water relations and various morpho-physiological processes in relation to scheduling of irrigation water. Studies on various aspects of groundnut nutrition are limited particularly under varied soil moisture regimes, hence efforts are needed to quantify the crop response *vis-a-vis* at different nutrient levels.

Comment [K20]: Telangana,

Comment [K21]: the

Comment [K22]: ,

Comment [K23]: quantify

Sulphur is essential to plant nutrition, and the fourth major plant nutrient next to nitrogen (N), phosphorus (P) and potassium (K) especially for oilseed crops. It is essential for synthesis of oil, in addition to protein production and activation of enzymes. It is a constituent of the aminoacids like cysteine, cystine and methionine and also involved in the formation of glucosides or glucosinolates, which on hydrolysis increase the oil content. It also plays an important role in chlorophyll and vitamin formation [2].

Comment [K24]: ,

Calcium is required by groundnut plants from the time when pegs begins to appear fruit formation, until the pods are mature. Calcium deficiency leads to high percentage of aborted seeds (empty pods), improperly filled pods [3].

Comment [K25]: begin

Comment [K26]: in

Comment [K27]: a

Comment [K28]: and

Gypsum is a moderately soluble source of the essential plant nutrients *viz.*, calcium and sulphur, and improve overall plant growth. Gypsum amendments can also improves soil physical properties, prevents crust formation, promote seedling emergence, increase water infiltration rates and movement through the soil profile.

Comment [K29]: improves

Comment [K30]: improve

Comment [K31]: prevent

Keeping this points in view, an experiment was designed in groundnut to study the effect of growth and yield of *rabi* groundnut for a profitable approach to realize the maximum yield potential.

Comment [K32]: point

Comment [K33]: on

Comment [K34]: realizing

## 2. Materials and Methods

The experiment was conducted during *rabi* season 2021 at Krishivignan Kendra, Palem, Nagarkurnool, India. The present research work is framed with an objective to study the effect of moisture regimes and gypsum levels on growth and yield of *rabi* groundnut. The experiment was laid out in split plot design. The soil was sandy loam in texture with pH of 7.1, Medium in available nitrogen, phosphorus and Potassium. The experiment consisted of 15 treatment combinations viz., three moisture regimes (0.6IW/CPE, 0.8 IW/CPE and 1.0 IW/CPE), and five gypsum levels ( $G_0$ - control,  $G_1$ - 300 kg ha<sup>-1</sup>,  $G_2$ - 400 kg ha<sup>-1</sup>,  $G_3$ - 500 kg ha<sup>-1</sup> and  $G_4$ - 600 kg ha<sup>-1</sup>) applied as basal at peg formation stage. Fertilizers urea and DAP (for N and P), MOP for K<sub>2</sub>O were applied to supply 30 kg N, 40 kg P<sub>2</sub>O<sub>5</sub> and 50 kg ha<sup>-1</sup>. Entire phosphorus and potash, half of the nitrogen (urea) were uniformly broadcasted before sowing and incorporated into the soil. Remaining half dose of nitrogen was top-dressed by band placement between 25-30 days after sowing depending on irrigation dates as per schedule of treatments. Crop was sown at a row distance of 30cm and plant to plant of 10cm. The crop was sown during 4<sup>th</sup> week of December and harvested at 4<sup>th</sup> week of April. Five plants were randomly select in each net plot area for taking observations on growth and yield attributing parameters. The samples were first dried under shade and then in electric oven at temperature of 60°C till attaining constant weight, on the basis of weight of these samples, then converted into dry matter production (kg ha<sup>-1</sup>). Data of growth attributes that is plant height (cm), number of branches per plant<sup>-1</sup>, leaf area (cm<sup>2</sup> plant<sup>-1</sup>), dry matter production (kg ha<sup>-1</sup>). Data obtained were statistically analyzed as suggested by [4].

### 3. Results and Discussion

#### 3.1 Growth parameters

##### 3.1.1 Moisture Regimes

The plant height varied among different moisture regimes. At before gypsum application stage the plant height showed non-significant with different moisture regimes. Higher values of plant height at 60, 90 DAS and at harvest (20.20, 23.99 and 27.87 cm) was recorded with I<sub>3</sub> (1.0 IW/CPE) which was significantly higher and on par with the treatment I<sub>2</sub> (0.8 IW/CPE) (18.16, 23.60 and 26.69 cm), this was due to availability of more number of irrigations which favoured the plant height. The results are in line with pawaret *al.* [5].

Higher number of branches plant<sup>-1</sup> (13.07 and 14.20) were recorded at 60 and 90 DAS with I<sub>3</sub> (1.0 IW/CPE) which was on par (12.42 and 13.17) with I<sub>2</sub> (0.8 IW/CPE) followed by I<sub>1</sub> (0.6 IW/CPE), whereas at before gypsum application plant height found to be non-significant. This is due to availability of more moisture at root zone depth due to frequent irrigations. These results were similar with Desai *et al.* [6].

Leaf area (934.86, 706.93 and 174.06 cm<sup>2</sup> plant<sup>-1</sup>) at 60, 90 DAS and at harvest, which was significantly higher with I<sub>3</sub> (1.0 IW/CPE) over other treatments except at harvest which is on par with I<sub>2</sub> (0.8 IW/CPE), dry matter plant<sup>-1</sup> (1113.76, 1996.78 and 2711.67 kg ha<sup>-1</sup>) at 60, 90 DAS and at harvest was significantly

Comment [K35]: the

Comment [K36]: 2021 at

Comment [K37]: studying

Comment [K38]: a

Comment [K39]: medium

Comment [K40]: potassium

Comment [K41]: broadcast

Comment [K42]: The remaining

Comment [K43]: ,

Comment [K44]: The crop

Comment [K45]: the

Comment [K46]: in

Comment [K47]: April

Comment [K48]: selected

Comment [K49]: an

Comment [K50]: a temperature

Comment [K51]: they attaining a constant

Comment [K52]: the

Comment [K53]: was

Comment [K54]: ,

Comment [K55]: significance

Comment [K56]: were

Comment [K57]: ,

Comment [K58]: the

Comment [K59]: irrigation

Comment [K60]: ,

Comment [K61]: ,

higher with I<sub>3</sub> (1.0 IW/CPE) which is on par with I<sub>2</sub> (0.8 IW/CPE) at 90 DAS and at harvest. Whereas, before gypsum application stage was found to be non-significant. Frequent irrigations provide more moisture, more nutrients to the crop and production of more biomass through the process of plant metabolism produced more leaf area. This findings were in close conformity with Laxmi [7] and with Karunasagar and Narsareddy [8].

### 3.1.2 Gypsum levels

Among the gypsum levels, G<sub>4</sub> @ 600 kg ha<sup>-1</sup> showed significantly higher values of plant height (20.84, 25.48 and 28.93 cm) which was on par with G<sub>3</sub> @ 500 kg ha<sup>-1</sup> (19.35, 24.74 and 28.07 cm) compared to other treatments. The increase in plant height might be attributed due to which might be due to the availability of calcium and sulphur through gypsum might have favoured for put forth and more plant height. Results were similar with Pasalaramya & Rajesh Singh [9] and Kannan [10].

Branches plant<sup>-1</sup> at before gypsum application was found to be non-significant with varied gypsum levels. At 60 and 90 DAS branches plant<sup>-1</sup> showed significantly higher values (13.80 and 15.85) with G<sub>4</sub> @ 600 kg ha<sup>-1</sup> which is on par (13.36) with G<sub>3</sub> @ 500 kg ha<sup>-1</sup> at 60 DAS. These findings were similar with findings of Rao & Shakawat [11].

Higher values of leaf area (cm<sup>2</sup> plant<sup>-1</sup>) (1351.3, 843.6 and 194.6 cm<sup>2</sup> plant<sup>-1</sup>) at 60, 90 DAS and at harvest respectively were observed with G<sub>4</sub> @ 600 kg ha<sup>-1</sup> which is significantly higher compared to other treatments. Dry matter accumulation (kg ha<sup>-1</sup>) (1566.7, 2201.5 and 3038.2 kg ha<sup>-1</sup>) showed significantly higher values with (G<sub>4</sub> @ 600 kg ha<sup>-1</sup>) compared to other treatments, while lowest values recorded with (G<sub>0</sub>- control) (650.4, 1770.5 and 2375.0 kg ha<sup>-1</sup>). Similar results were documented by Longkumer & Gohain [14].

The leaf area (cm<sup>2</sup> plant<sup>-1</sup>) interaction at 60 DAS with moisture regimes and gypsum levels at I<sub>3</sub>G<sub>4</sub> (1.0 IW/CPE) and (G<sub>4</sub> @ 600 kg ha<sup>-1</sup>) showed significantly higher value (1456.0 cm<sup>2</sup> plant<sup>-1</sup>) whereas lowest leaf area (463.0 cm<sup>2</sup> plant<sup>-1</sup>) recorded with I<sub>1</sub>G<sub>0</sub> (0.6 IW/CPE) and (G<sub>0</sub>- control). These findings were similar with findings of Nareshaef *et al.* [15].

## 3.2 Yield attributes

### 3.2.1 Moisture Regimes

Pod yield and halum yield (kg ha<sup>-1</sup>) (2337.6 and 1659.0 kg ha<sup>-1</sup>) is significantly higher with I<sub>3</sub> (1.0 IW/CPE) followed by I<sub>2</sub> (0.8 IW/CPE) (2254.0 and 1465.8 kg ha<sup>-1</sup>) while least value (2113.6 and 1309.4 kg ha<sup>-1</sup>) with I<sub>1</sub> (0.6 IW/CPE). Pod yield increase is due to increase in irrigation which ascribed to adequate

Comment [K62]: the

Comment [K63]: non-significant

Comment [K64]: irrigation provides

Comment [K65]: produces

Comment [K66]: areas

Comment [K67]: these

Comment [K68]: ,

Comment [K69]: ,

Comment [K70]: ,

Comment [K71]: ,

Comment [K72]: the

Comment [K73]: to the

Comment [K74]: The pod

Comment [K75]: increases

Comment [K76]: ,

moisture availability in turns have favored congenial conditions for the luxuriant growth of crop and consequently increased the values of the yield attributes with I<sub>3</sub> compare to I<sub>2</sub> and I<sub>1</sub> treatments. These findings were in close conformity with Naresha *et al.* [15], Dash *et al.* [16] and Kumaran *et al.* [17].

Comment [K77]: ,

Comment [K78]: turn has

### 3.2.2 Gypsum levels

Among varied levels of gypsum G<sub>4</sub> @ 600 kg ha<sup>-1</sup> showed significantly higher values (2829.7 and 1752.2 kg ha<sup>-1</sup>) of pod and haulm yield (kg ha<sup>-1</sup>) compared to other treatments. Least value of pod and haulm yield (1815.6 and 1070.0 kg ha<sup>-1</sup>) with G<sub>0</sub>- control. Gypsum application might have ensured adequate supply of calcium and sulphur, have favoured not only in pod formation but also in better filling of the pods thus would have increased number of filled pods. Salkeet *et al.* [18]

Comment [K79]: an

Comment [K80]: and

Comment [K81]: ,

Comment [K82]: the

Data regarding interaction between moisture regimes and gypsum levels on pod yield and haulm yield (kg ha<sup>-1</sup>) showed that significantly higher values with I<sub>3</sub>G<sub>4</sub> (1.0 IW/CPE and G<sub>4</sub> @ 600 kg ha<sup>-1</sup>) (3056.4 kg ha<sup>-1</sup>) and compared to other treatments, while least value (1717.8 kg ha<sup>-1</sup>) with I<sub>1</sub>G<sub>0</sub> (0.6 IW/CPE and G<sub>0</sub>- control). Higher amount of gypsum level increased the calcium and sulphur levels have favored not only pod formation but also in better filling of the pods and more irrigations favoured the pod yield in groundnut. These findings were similar with Naresh *et al.* [15].

Comment [K83]: amounts

Comment [K84]: levels

Comment [K85]: to

## 4. CONCLUSION

This field experiment inferred that moisture regime I<sub>3</sub> (1.0 IW/CPE) and Gypsum level (G<sub>4</sub> @ 600 kg ha<sup>-1</sup>) showed highest values of growth parameters and yield attributes which can be cultivated for realizing higher pod yield under *rabi* conditions on sandy loam soils of Telangana region.

Comment [K86]: the

Comment [K87]: yields

Comment [K88]: the

Table: 1 Plant height (cm) of *rabi* groundnut as influenced by moisture regimes and gypsum levels at different crop growth stages

Treatments	Plant height (cm)			
	Before gypsum	60 DAS	90 DAS	At harvest
<b>MOISTURE REGIMES (I)</b>				
I <sub>1</sub> - 0.6 IW/CPE	12.07	16.74	23.20	24.36
I <sub>2</sub> - 0.8 IW/CPE	12.78	18.16	23.60	26.69
I <sub>3</sub> - 1.0 IW/CPE	13.16	20.20	23.99	27.87
SEm ±	0.47	0.54	0.04	0.55
CD (P=0.05)	NS	2.20	0.19	2.24
<b>GYPSUM LEVELS (GP)</b>				
G <sub>0</sub> - Control	12.01	16.01	21.44	23.53

G <sub>1</sub> - Gypsum @ 300 kg ha <sup>-1</sup>	12.48	17.26	22.59	24.34
G <sub>2</sub> - Gypsum @ 400 kg ha <sup>-1</sup>	12.33	18.36	23.73	26.66
G <sub>3</sub> - Gypsum @ 500 kg ha <sup>-1</sup>	13.27	19.35	24.73	28.07
G <sub>4</sub> - Gypsum @ 600 kg ha <sup>-1</sup>	13.24	20.84	25.48	28.93
SEm±	0.34	0.61	0.17	0.81
CD (P=0.05)	NS	1.80	0.51	2.38
INTERACTION (I X GP)				
Sub treatment at same level of main treatment				
SEm±	1.06	1.22	0.10	1.24
CD (P= 0.05)	NS	NS	NS	NS
Main treatment at same or different level of sub treatment				
SEm±	0.71	1.09	0.27	1.37
CD (P=0.05)	NS	NS	NS	NS

Table: 2 Number of branches plant<sup>-1</sup> of *rabigroundnut* as influenced by moisture regimes and gypsum levels at different crop growth stages

Treatments	Number of branches plant <sup>-1</sup>		
	Before gypsum	60 DAS	90 DAS
MOISTURE REGIMES (I)			
I <sub>1</sub> - 0.6 IW/CPE	6.90	11.36	12.19
I <sub>2</sub> - 0.8 IW/CPE	7.13	12.42	13.17
I <sub>3</sub> - 1.0 IW/CPE	7.24	13.07	14.20
SEm ±	0.32	0.18	0.35
CD (P=0.05)	NS	0.73	1.41
GYPSUM LEVELS (GP)			
G <sub>0</sub> - Control	6.37	10.83	11.22
G <sub>1</sub> - Gypsum @ 300 kg ha <sup>-1</sup>	6.92	11.22	11.81
G <sub>2</sub> - Gypsum @ 400 kg ha <sup>-1</sup>	6.87	12.20	12.37

G <sub>3</sub> - Gypsum @ 500 kg ha <sup>-1</sup>	7.42	13.36	14.67
G <sub>4</sub> - Gypsum @ 600 kg ha <sup>-1</sup>	7.86	13.80	15.85
SEm±	0.37	0.29	0.39
CD (P=0.05)	NS	0.87	1.15
INTERACTION (I X GP)			
Sub treatment at same level of main treatment			
SEm±	0.72	0.40	0.78
CD (P= 0.05)	NS	NS	NS
Main treatment at same or different level of sub treatment			
SEm±	0.65	0.49	0.70
CD (P=0.05)	NS	NS	NS

Table: 3 Leaf area (cm<sup>2</sup> plant<sup>-1</sup>) of *rabigrroundnut* as influenced by moisture regimes and gypsum levels at different crop growth stages

Treatments	Leaf area cm <sup>2</sup> plant <sup>-1</sup>			
	Before gypsum	60 DAS	90 DAS	At harvest
MOISTURE REGIMES (I)				
I <sub>1</sub> - 0.6 IW/CPE	432.40	777.06	650.60	165.20
I <sub>2</sub> - 0.8 IW/CPE	433.80	852.06	682.33	170.40
I <sub>3</sub> - 1.0 IW/CPE	434.20	934.86	706.93	174.06
SEm ±	4.03	4.25	3.98	1.18
CD (P=0.05)	NS	17.16	16.06	4.79
GYPSUM LEVELS (GP)				
G <sub>0</sub> - Control	425.88	485.00	525.00	146.66
G <sub>1</sub> - Gypsum @ 300 kg ha <sup>-1</sup>	429.22	624.11	603.88	156.66
G <sub>2</sub> - Gypsum @ 400 kg ha <sup>-1</sup>	435.22	799.33	672.44	168.00
G <sub>3</sub> - Gypsum @ 500 kg ha <sup>-1</sup>	437.55	1013.55	754.77	183.44
G <sub>4</sub> - Gypsum @ 600 kg ha <sup>-1</sup>	439.44	1351.33	843.66	194.66
SEm±	3.74	8.44	5.99	1.20
CD (P=0.05)	NS	24.79	17.60	3.53
INTERACTION (I X GP)				
Sub treatment at same level of main treatment				
SEm±	9.02	9.52	8.91	2.66
CD (P= 0.05)	NS	44.47	NS	NS
Main treatment at same or different level of sub treatment				
SEm±	7.06	13.75	10.10	2.21
CD (P=0.05)	NS	41.83	NS	NS

Table: 4 Drymatter production (kg ha<sup>-1</sup>) of *rabigroundnut* as influenced by moisture regimes and gypsum levels at different crop growth stages

Treatments	Drymatter production (kg ha <sup>-1</sup> )			
	Before gypsum	60 DAS	90 DAS	At harvest
<b>MOISTURE REGIMES (I)</b>				
I <sub>1</sub> - 0.6 IW/CPE	442.45	946.82	1933.84	2618.42
I <sub>2</sub> - 0.8 IW/CPE	442.52	1017.67	1973.47	2663.44
I <sub>3</sub> - 1.0 IW/CPE	443.04	1113.76	1996.78	2711.67
SEm ±	2.48	10.48	7.29	13.91
CD (P=0.05)	NS	41.26	29.39	56.11
<b>GYPSUM LEVELS (GP)</b>				
G <sub>0</sub> - Control	442.11	650.48	1770.56	2375.07
G <sub>1</sub> - Gypsum @ 300 kg ha <sup>-1</sup>	442.26	812.96	1870.45	2538.41
G <sub>2</sub> - Gypsum @ 400 kg ha <sup>-1</sup>	442.43	929.59	1941.89	2622.37
G <sub>3</sub> - Gypsum @ 500 kg ha <sup>-1</sup>	442.67	1170.59	2055.67	2748.49
G <sub>4</sub> - Gypsum @ 600 kg ha <sup>-1</sup>	443.89	1566.78	2201.59	3038.22
SEm±	3.73	10.71	18.34	21.91
CD (P=0.05)	NS	31.45	53.86	64.34
<b>INTERACTION (I X GP)</b>				
Sub treatment at same level of main treatment				
SEm±	5.54	23.43	16.30	31.12
CD (P= 0.05)	NS	60.28	NS	NS
Main treatment at same or different level of sub treatment				
SEm±	6.29	19.63	29.33	36.69
CD (P=0.05)	NS	63.77	NS	NS

Table : 4.a Dry matter production (kg ha<sup>-1</sup>) of *rabigroundnut* at 60 DAS as influenced by interaction between moisture regimes and gypsum levels

GYPSUM LEVELS	MOISTURE REGIMES (I)			
	I <sub>1</sub> - 0.6 IW/CPE	I <sub>2</sub> - 0.8 IW/CPE	I <sub>3</sub> -1.0 IW/CPE	MEAN
G <sub>0</sub> - Control	590.2	650.6	710.5	650.4
G <sub>1</sub> - Gypsum @ 300 kg ha <sup>-1</sup>	775.7	812.5	850.5	812.9
G <sub>2</sub> - Gypsum @ 400 kg ha <sup>-1</sup>	890.6	901.4	996.6	929.5
G <sub>3</sub> - Gypsum @ 500 kg ha <sup>-1</sup>	1201.0	1145.2	1345.5	1170.5
G <sub>4</sub> - Gypsum @ 600 kg ha <sup>-1</sup>	1456.4	1578.4	1665.4	1566.7

MEAN	946.8	1017.6	1113.7	
Sub treatment at same level of main treatment				
SEm±				23.4
CD (P=0.05)				60.2
Main treatment at same or different level of sub treatment				
SEm±				19.6
CD (P=0.05)				63.7

Table: 5 Yield attributes of *ragi* groundnut as influenced by moisture regimes and gypsum levels at different crop growth stages

Treatments	Yield	
	Pod yield (kg ha <sup>-1</sup> )	Halum yield (kg ha <sup>-1</sup> )
MOISTURE REGIMES (I)		
I <sub>1</sub> - 0.6 IW/CPE	2113.60	1309.40
I <sub>2</sub> - 0.8 IW/CPE	2254.00	1465.86
I <sub>3</sub> - 1.0 IW/CPE	2337.68	1659.06
SEm ±	11.01	9.18
CD (P=0.05)	44.39	37.02
GYPSUM LEVELS (GP)		
G <sub>0</sub> - Control	1815.62	1070.00
G <sub>1</sub> - Gypsum @ 300 kg ha <sup>-1</sup>	1964.53	1410.00
G <sub>2</sub> - Gypsum @ 400 kg ha <sup>-1</sup>	2145.76	1532.22
G <sub>3</sub> - Gypsum @ 500 kg ha <sup>-1</sup>	2419.73	1626.11
G <sub>4</sub> - Gypsum @ 600 kg ha <sup>-1</sup>	2829.78	1752.22
SEm±	22.28	10.299
CD (P=0.05)	65.43	30.24
INTERACTION (I X GP)		
Sub treatment at same level of main treatment		
SEm±		20.53
CD (P= 0.05)		57.24
Main treatment at same or different level of sub treatment		
SEm±		18.41

CD (P=0.05)	110.07	59.07
-------------	--------	-------

Table: 5.a Pod yield (kg ha<sup>-1</sup>) of *rabi* groundnut as influenced by interaction between moisture regimes and gypsum levels

GYPSUM LEVELS	MOISTURE REGIMES (I)			
	I <sub>1</sub> - 0.6 IW/CPE	I <sub>2</sub> - 0.8 IW/CPE	I <sub>3</sub> -1.0 IW/CPE	MEAN
G <sub>0</sub> - Control	1717.8	1834.5	1894.4	1815.6
G <sub>1</sub> - Gypsum @ 300 kg ha <sup>-1</sup>	1902.3	1967.4	2023.7	1964.5
G <sub>2</sub> - Gypsum @ 400 kg ha <sup>-1</sup>	2023.5	2167.9	2245.7	2145.7
G <sub>3</sub> - Gypsum @ 500 kg ha <sup>-1</sup>	2356.7	2434.5	2467.9	2419.7
G <sub>4</sub> - Gypsum @ 600 kg ha <sup>-1</sup>	2567.4	2865.4	3056.4	2829.7
MEAN	2113.6	2254.0	2337.6	
Sub treatment at same level of main treatment				
SEm±				24.6
CD (P=0.05)				117.2
Main treatment at same or different level of sub treatment				
SEm±				36.2
CD (P=0.05)				110.0

Table: 5.b Haulm yield (kg ha<sup>-1</sup>) of *rabi* groundnut as influenced by interaction between moisture regimes and gypsum levels

GYPSUM LEVELS	MOISTURE REGIMES (I)			
	I <sub>1</sub> - 0.6 IW/CPE	I <sub>2</sub> - 0.8 IW/CPE	I <sub>3</sub> -1.0 IW/CPE	MEAN
G <sub>0</sub> - Control	1011.3	1098.0	1100.6	1070.0
G <sub>1</sub> - Gypsum @ 300 kg ha <sup>-1</sup>	1234.0	1398.0	1598.0	1410.0
G <sub>2</sub> - Gypsum @ 400 kg ha <sup>-1</sup>	1301.0	1498.0	1797.6	1532.2
G <sub>3</sub> - Gypsum @ 500 kg ha <sup>-1</sup>	1434.0	1601.3	1843.0	1626.1
G <sub>4</sub> - Gypsum @ 600 kg ha <sup>-1</sup>	1566.6	1734.0	1956.0	1752.2
MEAN	1309.4	1465.8	1659.0	
Sub treatment at same level of main treatment				

SEm±	20.5
CD (P=0.05)	57.2
Main treatment at same or different level of sub treatment	
SEm±	18.4
CD (P=0.05)	59.0

#### Reference:

1. Indiatat. Area, production and productivity of groundnut 2019-20. <https://www.indistat.com>
2. Gosh,P.K.,Hati,K.M.,Mandal,K.G.,Misra,A.K.,Chaudhary,A.SandBandyopadhyay, K. K., 2000. Sulphur nutrition in oilseeds and oilseed-basedcroppingsystems.*Fertiliser News*.45(8): 27–40.
3. Islam, A.M. 2013. Effect of phosphorus, calcium and boron on the growth and yield of groundnut (*Arachis hypogea*L.). *International Journal of Bio-science and Bio- technology*. 5 (3).
4. Panse,V.GandSukhatme,P.V.1985.Statisticalmethodsforagriculturalworkers. ICAR publication.
5. Pawar, D. D., Dingre, S. K and Nanaware, D. M. 2013. Yield and quality of *summer*groundnut under different irrigation scheduling through micro sprinkler in clayloamsoilsofwesternMaharashtra.*JournalofAgriculturalResearchandTechnology*.38 (1): 102-106.
6. Desai, N.D., Joshi, R.S., and Patel K. R. 1985. Effect of irrigation on growth andyield attributes of *summer* groundnut.*AgriculturalScienceDigest*.5:63-66 .
7. Lakshmi, M. B. 1990. Studies on irrigation and phosphorus management in groundnut-maize cropping sequence. PhD Thesis andhra Pradesh Agricultural University,Hyderabad.
8. Karunasagar, G and Narsareddy, S. 2003. Effect of irrigation regimes on growth andyieldofintercroppedgroundnut.*Journal ofOilseedsResearch*. 20(1):124-125.
9. Ramya, P. and Singh, R., 2022. Effect of gypsum and boron on growth and yield of groundnut [*Arachis hypogea* L.]
10. Kannan, P., Swaminathan, C. and Ponmani, S., 2017. Sulfur nutrition for enhancing rainfed groundnut productivity in typical alfisol of semi-arid regions in India. *Journal of Plant Nutrition*, 40(6), pp.828-840.
11. Rao, S.S. and Shaktawat, M.S., 2001. Effect of organic manure, phosphorus and gypsum on growth, yield and quality of groundnut (*Arachishypogaea* L.). *Indian journal of plant physiology*, 6(3), pp.306-311.
12. Longkumer, I.T. and Gohain, T., 2012. Effect of sulphur and calcium on growth and yield of sesame under rainfed condition of Nagaland. *Annals of Plant and Soil Research*, 14(1), pp.58-60.
13. Naresha, R., Laxminarayana, P., Devi, K.S. and Sailaja, V., 2018. Effect of Irrigation Scheduling and Phosphogypsum Levels on Yield Attributes, Yield and Available Nutrients in Soil after Harvest of Rabi Groundnut. *Int. J. Pure App. Biosci*, 6(2), pp.1300-1308.

14. Dash, A. K., Nayak, B. R., Panigrahy, N., Mohapatra, S and Samant, P. K. 2013. Performance of groundnut (*Arachis hypogaea* L.) under different levels of sulphur and irrigation. *Indian Journal of Agronomy*, 58(4): 578-582.
15. Kumaran, S., Solaimalai, A., Ravisankar, N. and Chandrasekaran, B., 2000. Role of organic, inorganic fertilizers and gypsum application in groundnut: A Review. *Agricultural reviews-agricultural research communications centre India*, 21(3), pp.193-198.
16. Salke, S.R., Shaikh, A.A. and Dalavi, N.D., 2011. Effect of phosphatic fertilizers, gypsum and sulphur on groundnut (*Arachis hypogaea* L.). *BIOINFOLET-A Quarterly Journal of Life Sciences*, 8(3), pp.276-277.

UNDER PEER REVIEW