

Spatio-temporal dynamics of land use in lake ecosystems in the Sudano-Guinean zone([A case study in Adamaoua-Cameroon](#))

Abstract

This study aims to characterise the spatio-temporal mutations experienced by four lake ecosystems in the Vina Department, Adamaoua Region, Cameroon, over 31 years between 1990 and 2021, due to strong anthropic activities caused by the increase in demography. Landsat satellite images from 1990, 2006 and 2021 were exploited using remote sensing and GIS. Unsupervised classification was used to obtain nine land cover and land use classes (agricultural space, forest gallery, tree savannah, shrub savannah, grassy savannah, bare soil, water surface, hydromorphic zone and housing zone). From 1990 to 2021, the evolutionary typology of the plant formations is essentially regressive for the shrub savannahs (16.21 ha at Lake Bini, 30.56 ha at Lake Dang, 118.13 at Lake Mballang and 9.67 ha at Lake Ngaoundaba). On the other hand, the dynamics is progressive for the agricultural space (24.40 ha at Lake Bini, 24.38 ha at Lake Dang, 16.72 at Lake Mballang). The water surface, being the particular specificity of the lacustrine wetlands, has decreased in the lacustrine ecosystem of Bini (0.27ha) and Mballang (0.04). On the other hand, there was an increase in the lake ecosystems of Lake Dang (0.40 ha) and Ngaoundaba (0.21 ha). The banks of Lake Ngaoundaba are devoid of remarkable agricultural space. ~~The factors of this degradation are mainly human (agriculture, overgrazing and galloping demography) and result in the fragmentation and fragmentation of natural habitats to the benefit of cultivated areas and purely anthropogenic housing zones.~~ [\(Recase please\) As well make some recommendation too at the end of the abstract.](#)

Key words: lake ecosystems; spatio-temporal mutations; occupation units, land use; Vina Department; Adamaoua-Cameroon.

Introduction

~~In the Sahelian countries, the combined effects of changing rainfall patterns, population growth and land degradation have led to a deterioration of soil and water resources.~~ [\(source please acknowledge the author\)](#) Since the 19th century, the Sahel has experienced both droughts and wetter periods [\(As above reference pls\)](#). In addition to this natural variability, it is now necessary to take into account the impact of climate change on the rainfall regime [\(De France et al., 2017\)](#). ~~if the author is same as above than pls use linking word to link the 1st and 2nd sentence e.g moreover, as a result etc~~ These uncertain factors influencing rainfall have a major impact on the evolution of Sahelian lake systems. ~~Indeed, a large~~

Formatted: Font color: Auto

Formatted: Font color: Red

Formatted: Font color: Red, Strikethrough

Formatted: Font color: Red

Formatted: Font color: Red

Formatted: Font color: Auto

Formatted: Font color: Red

Formatted: Superscript

Formatted: Font color: Red, Strikethrough

Formatted: Font color: Red

Formatted: Superscript

Formatted: Superscript

Formatted: Font color: Red

proportion of these lakes are endoreic systems that form reservoirs during the dry season. (reference) Their vulnerability to climatic constraints, combined with their great societal importance, makes it essential to understand their hydrological functioning. In the context where ecosystem landscapes have experienced major disturbances due to both natural processes and anthropogenic activities (Recase) (Vitteket *al.*, 2014). Lake ecosystems are undergoing many changes related to more intense human activities to meet the socio-economic needs of the populations. This is materialised by the fragmentation of the landscape and the loss of habitats of the flora and fauna of the banks and the hydrosystem (Bahroun, 2011).

Formatted: Font color: Red

In Cameroon, the pressure exerted on lake ecosystems by populations in search of new fertile land for market gardening or even off-season crops and fishery resources is at the origin of the transformation and fragmentation of natural habitats (Touchard, 1996). These transformations result in the conversion of natural formations into fields, ranches, effluent outfalls, and the erosion of biological diversity and the shortening of the life of lakes. (Reference)

Formatted: Font color: Red

Formatted: Font color: Auto

The ecological manifestations resulting from this state of affairs are worrying for the populations that depend directly or indirectly on the natural resources that are being lost. Understanding the succession dynamics of the constituent elements of lake ecosystems is therefore a necessity for sustainable management of natural resources for the benefit of the population's well-being (Messager *et al.*, 2016). In addition, degradation and deforestation caused by human activities are at the origin of the emission of a significant amount of carbon. Anthropogenic carbon leads to an increase in GHG Green House gases (GHG) write in full than subsequently abbreviate concentration, which can contribute to climate change (Chaplin-Kramer *et al.*, 2015). Quantifying degradation or deforestation processes is therefore a necessity for the implementation of a policy to increase ecosystem carbon stocks in the framework of REDD+ write in full as above pls. Therefore, Remote sensing is an important tool for providing quantitative information over time and space on land use and occupation.

Formatted: Font color: Red

Formatted: Font color: Red

Formatted: Strikethrough

The study of land cover dynamics in time and space, using satellite images and remote sensing techniques, contributes effectively to the sustainable management of natural resources (Leimgruber *et al.*, 2005). Indeed, the analysis of satellite images through remote sensing provides data on land cover and its changes over large inaccessible areas repeatedly for a period of over 40 years (Vitteket *al.*, 2014).

In Cameroon, land use studies have reported degradation of wetland lake ecosystems without quantifying the factors that have changed (Reference). Few studies have focused on the dynamics of lake complexes over time, and analysed quantitatively and qualitatively the changes that have occurred within these units. In

Formatted: Font color: Red

Formatted: Font color: Red, Strikethrough

addition, spatially explicit knowledge of all these parameters is crucial for understanding and modelling a wide variety of Earth system processes and interactions with the environment, including water budgets (Muller *et al.*, 2014); carbon and methane exchange rates (Bastviken *et al.*, 2011); sediment trapping (Downing *et al.*, 2008); heat flux and coupled effects of weather and climate (Balsamo *et al.*, 2012); the cycling of pollutants and nutrients (Norges *et al.*, 2009); as well as associated ecological processes such as lake productivity (Stæhr *et al.*, 2012); species richness (Dodson *et al.*, 2000); food chain dynamics (Post *et al.*, 2010); and inland fisheries yields (De Graaf *et al.*, 2012).

The overall objective of this work is to evaluate the spatio-temporal dynamics of the land use units of the lake complexes (Lake Bini, Lake Dang, Lake Mballang and Lake Ngaoundaba) (where location pls), by coupling remote sensing tools with geographic information systems.

Formatted: Font color: Red

1. Data and methods

1.1. Description of the study area

I will like if you have started with the study area is Located between 6° and 8° North latitude and between 11° and 15° (reference as well) East longitude, the Adamaoua Cameroon region extends over 63701 km². It is bordered to the North by the North region, to the South by the Centre and East region, to the East by the Central African Republics, to the West by the Federal Republic of Nigeria and to the South-West by the West and North West regions. (reference) The climate is Sudano-Guinean, mild and cool, characterised by two seasons, a rainy season (April to October) and a dry season (November to March) (MINEF, 1994). The average annual rainfall is 1500mm per year spread over seven months. It is considered the "water tower" of the country because of its highly diversified water network, which originates in the same region and many of the country's rivers have their source there (reference). The investigations took place in the Vina department (Fig. 1), more precisely in the lake ecosystems of the districts of Ngaoundéré III (Lake Bini and Lake Dang), Nymbaka (Lake Ngaoundaba) and Ngan-ha (Mballang).

Formatted: Font: Not Bold

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font: Not Bold, Font color: Red

Formatted: Font: Not Bold, Font color: Red

Formatted: Font: Not Bold, Font color: Red

The paper will be reach if you have highlighted the profile of the study area with reference to : location, brief history, climate (specifically rainfall data for those period of study), topography, vegetation and soil as well as demography.

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font: (Default) +Body, 14 pt, Not Bold

Formatted: Font: (Default) +Body, 14 pt, Not Bold

Refer to your satellite images they are termed as plates not figures.

Maps and charts are figures while image or pictures are plates pls reconcile




Figure 1: Location map of the Adamaoua region (A), Vina Department (B) and lake ecosystems (C)

1.2 Planimetric data

The data used in the study of land cover dynamics here are essentially satellite images, all captured at the beginning of the dry season. Indeed, the very low cloud cover of the dry season allows the satellite sensor to have good views (Inoussa *et al.*, 2011). Therefore, Landsat_5 from February 1990, Landsat_7 from February 2006 and Landsat_8 from February 2021, all with a resolution of 30 m, were uploaded to the website <http://earthexplorer.usgs.gov> in GEOTIFF format. Topographic maps and base maps from the National Institute of Cartography were also used in this study. A GARMIN GPS (Global Positioning System) was used to locate the position of the various field control points. In addition to these planimetric data, some data (density, sown areas, pedology, etc.) were collected during a light survey of the populations to better profile the analysis of the dynamics of land use (Agbanou *et al.*, 2018).

1.3 Methods

After downloading these images (1990; 2006 and 2021) with less than 10% cloud cover, we imported them into Envi 4.5. It is in this software that the supervised classification operations took place. Thus we applied the supervised classification algorithm on the maximum likelihood parameter. Beforehand, it was necessary to import the shapefile vector layer of the boundary of the different lake ecosystems in order to create the metadata on which a coloured composition is applied in order to create the mask. The creation of the mask is essential since it

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Formatted: Default Paragraph Font, Font: (Default) +Body, French (France)

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

allows us to delimit only the area in which we want to work. Once the mask has been created, it must be applied to the metadata and then a coloured composition (RGB) must be made to leave only the area to be studied. Here the radiometric bands that can provide us with information on land use and land cover units were overlaid by additive synthesis of primary colours (Agbanouet *al.*, 2018). Thus, the bands were combined through a false colour composition with the combination of bands 2, 3 and 4 for the ETM+ image and 3, 4 and 5 for the OLI-TIR image. This operation therefore consisted in making a false colour composition in RGB (Red-Green-Blue). In order to facilitate the overlay of images for diachronic analysis, the 1990, 2006 and 2021 images were co-registered using the image-to-image coregistration method, which ensures a good alignment of the pixels of the different images. The land use points were painted in different colours characteristic of the land uses.

Then, the study area was cut to the part of the image corresponding to the area of interest from the polygon representing the boundaries of the Boroughs. The supervised classification method according to the maximum likelihood algorithm was used. The classification established and the colours used to represent each land use category were adapted from the Corine Land Cover (CLC) nomenclature. The accuracy and Kappa index were used to validate the classifications. The land cover maps produced provided the basic data for the analysis and quantification of the landscape. The areas of the different land-use categories were calculated and diagrams were drawn to assess the speed and intensity of change. Both descriptive and diachronic approaches were used for the analysis.

Quantum Gis 2.14 (Geographic Information System) was used to design the maps and calculate the areas of each land use class via the "groupstats" extension, which can be downloaded directly from the list of Qgis extensions. The following landscape indices were calculated to assess the observed dynamics:

❖ Kappa coefficient or index

The coefficient or Kappa index, which corresponds to the ratio of the number of well ranked pixels to the total number of pixels surveyed, is retained as it better indicates the agreement between the predicted models and reality (Tilahunet *al.*, 2015). It is expressed as:

$$K = (P_o - P_a) / 1 - P_a$$

With P_o = actual percentage of land cover elements classified, P_a = estimate of the probability of obtaining a correct classification.

❖ Normalized Differential Vegetation Index (NDVI)

This is the most widely used vegetation index and is calculated from the visible and nearinfrared light reflected by the vegetation. Healthy vegetation

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

absorbs most of the visible light that reaches it and reflects most of the near infrared light (Halima and Djamel, 2020). It is calculated using this formula:

$$\text{NDVI} = (\text{NIR} - \text{Red}) / (\text{NIR} + \text{Red})$$

Formatted: No underline, Font color: Auto

Knowing that NIR is the Near Infrared band, represented by band 4 in Landsat 5TM and 7ETM+ images and band 5 in Landsat 8 OLI/TIRS images; Red is the Red Band, represented by band 3 in Landsat 5TM and 7ETM+ images and band 4 in Landsat 8 OLI/TIRS images. Knowing that NDVI values are theoretically between -1 and +1, negative values correspond to non-vegetated surfaces such as snow, water or clouds. For bare soil, the NDVI has values close to 0. Vegetation formations have positive NDVI values, generally between 0.1 and 0.7, with the highest values corresponding to the densest cover. The NDVI index offers us the possibility to analyse the change in vegetation cover during the study years (Halima and Djamel, 2020).

Formatted: No underline, Font color: Auto

❖ Typology of changes (Δ)

The typology of changes was characterised based on the nomenclature described by FAO (2011). To do so, a subtraction is performed between the digital values of the two images so that the change detection consisted in identifying the change of codes for the homologous vectors (Δi). After the land use maps of 1990; 2006 and 2021 were drawn up, a two-by-two comparative analysis (1990-2006; 1990-2006 and 1990-2021) was made. The analysis of the evolution of the different land use units was done using the

Formatted: No underline, Font color: Auto

$$\Delta = \text{Sit} - \text{St1}$$

Formatted: No underline, Font color: Auto

Sit the area of land occupation by a unit at an initial date, St1 the area of land occupation of the same unit at a given date and Δ the variation of this inter-periodic area. If: $\Delta = 0$ then there is stability (S); $\Delta > 0$ then there is progressive evolution (P); $\Delta < 0$ then there is regression (R).

Formatted: No underline, Font color: Auto

❖ Average annual rates of spatial expansion

The areas of land use and land cover units between the two dates were used to calculate the annual expansion rate according to the following formula from Bernier (1992):

Formatted: No underline, Font color: Auto

$$T = (\ln s_1 - \ln s_0) / \ln e (t_1 - t_0) * 100$$

Formatted: Font color: Auto

S0 and S1 represent the areas of a landscape unit at date t0 and t1 respectively; ln is the natural logarithm, and e represents the base of the natural logarithm ($e = 2.71828$). This rate expresses the annual proportion of change in each unit of land cover and land use.

Formatted: Font color: Auto

Formatted: Font color: Auto

❖ Conversion rate

Formatted: Font color: Auto

In order to quantify the change between two dates, the conversion rate of each discriminated land-use class was calculated. The conversion rate measures the degree of conversion of a given unit into other landscape units between two dates t_0 and t_1 . It is obtained from the transition matrix (Arouna, 2012), according to the formula :

$$Tc = (S_{it} - S_{is}) / S_{it} * 100$$

S_{it} : Area of the landscape unit at the initial date t ; S_{is} : Area of the same unit remaining stable at date t_1

❖ Speed of change of land use categories

In order to know the rate of change of the identified land use categories (Agbanou *et al.*, 2018), the following formula was used:

$$\Delta s = \frac{St_1 - Sit}{t_1 - t_i}$$

Where: Δs = Rate of change (extension or regression in ha/year); S_{it} = Area occupied by the occupation category considered in year 1 (ha); S_{t1} = Area occupied by the occupation category considered in year 2 (ha); t_i = year 1; t_1 = year 2.

2. Results

2.1. Discriminated units

Table 1 presents the quality evaluation indices of the classified images. It can be seen that the overall statistical accuracies of the classified images are between 90.23 and 97.14% with kappa indices ranging from 86.93 to 95.65% (Table 1). These indices are respectively better in increasing chronological order. The lake ecosystems of Lake Dang, Bini, Mballang and Ngaoundaba recorded the best overall accuracy and Kappa indices respectively

Table 1: Quality assessment scores for classified image


	Lake ecosystems	Overall accuracy (%)	Kappa (%)
1990	lake Bini	96,78	95,30
	lake Dang	97,14	95,65
	lake Mballang	95,27	93,81
	lake Ngaoundaba	93,52	92,09
	lake Bini	96,11	94,64

	lake Dang	97,05	95,56
	lakeMballang	92,53	91,11
	lakeNgaoundaba	90,23	88,85
	lake Bini	95,84	94,37
2021	lake Dang	96,47	94,99
	lakeMballang	94,31	92,86
	lakeNgaoundaba	92,64	91,22

- Formatted: Font color: Auto
- Formatted: Font color: Auto
- Formatted: Font color: Auto
- Formatted: Font color: Auto
- Formatted: Font color: Auto
- Formatted: Font:
- Formatted: Font color: Auto
- Formatted: Font color: Auto
- Formatted: Font:
- Formatted: Font color: Auto
- Formatted: Font color: Auto
- Formatted: Font:
- Formatted: Font:
- Formatted: Font color: Auto
- Formatted: Font color: Auto
- Formatted: Font color: Auto
- Formatted: Font:
- Formatted: Font color: Auto
- Formatted: Font color: Auto
- Formatted: Font:
- Formatted: Font color: Auto
- Formatted: Font color: Auto
- Formatted: Font color: Auto
- Formatted: Font color: Auto

2.2. Spatial distribution of land use and land cover units in 1990, 2006 and 2021

Figures 2, 3, 4 and 5 show the different land use changes in the four lake ecosystems between 1990 and 2021. It is apparent that nine common land-use unit classes have been discriminated in the lake ecosystems of Mballng (Fig. 4) and Dang (Fig.3). These are agricultural space, forest gallery, tree savannah, shrub savannah, grassy savannah, bare soil, water surface, hydromorphic zone, and housing area. On the other hand, the Lake Bini ecosystem (Fig. 2) has eight land use and occupancy unit classes, including agricultural space, forest gallery, tree savannah, shrub savannah, bare soil, water surface, hydromorphic zone and housing area. The Ngaoundaba lake ecosystem (Fig. 5) presents the same land use and occupancy unit classes as the Bini lake ecosystem (Fig. 2), except for the agricultural area.






Figure 5: Spatial distribution of land use units in the Lake Ngaoundaba ecosystem between 1990 and 2021

2.3 Estimation of the Normalized Differential Vegetation Index (NDVI)

The NDVI maps (Fig. 6; 7; 8; and 8) show a remarkable difference between the different years. The recorded values of the NDVI of lake ecosystems varied from 0 to 0.214; from 0 to 0.214; from -0.116 to 0.326 and from -0.06 to 0.282 respectively for the lake ecosystems of Bini (Fig 6), Dang (Fig 7), Mballang (Fig 8) and Ngaoundaba (Fig 9). The low NDVI values are recorded at the level of the water bodies of the different lakes while the maximum values are recorded at the level of the riparian strips.

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto




Figure 6: Spatial distribution of land use units in the Lake Bini ecosystem between 1990 and 2021

Formatted: Font color: Auto




Figure 7: Spatial distribution of land use units in the Lake Dang ecosystem between 1990 and 2021.

Formatted: Font color: Auto




Figure 8: Spatial distribution of land use units in the Lake Mballang ecosystem between 1990 and 2021

Formatted: Font color: Auto




Figure 9: Spatial distribution of land use units in the Lake Ngaoundaba ecosystem between 1990 and 2021

Formatted: Font color: Auto

Formatted: Font color: Auto


2.4. Dynamics of land use and land cover units

Formatted: Font color: Auto

Formatted: Font color: Auto

Figure 10 shows the different land use units of the Bini lake ecosystem. Examination of this figure shows that in 1990, the shrub savannah (26.43%) followed by the hydromorphic zone (25.68 %) and tree savannah (16.29 %) recorded the largest areas compared to the habitation area (0.26 %). In 2006, the hydromorphic zone (44.13 %) followed by the agricultural area (29.28 %) and the

shrub savannah came in first place to the detriment of the forest gallery (0.98%), which was the least represented. In contrast, in 2021, the agricultural area (41.05 ha) is followed by the hydromorphic zone (29.30%) and the residential area (10.11%). The forest gallery (0.85%) is the land use unit with the smallest surface area in 2021.




AA: Agricultural area, FG: Forest gallery, TS: Tree savannah, SS: Shrub savannah, GS: Grassy savannah, BS: Bare soil, WS: Water surface, HYA: Hydromorphic area, HOA: Housing area

Figure 10: Areas of different land use units in the Bini lake ecosystem

For the Dang lake ecosystem (Fig. 11), it can be seen that in 1990 the grassy savannah (40.23%) followed by the shrub savannah (21.29%) and the water area (9.65%) occupied more surface area compared to the residential area (0.59) which occupied the least surface area. In 2006, the grassy savannah (40.23%) followed by the residential area (21.09%) and the agricultural area (14.93%) occupies more surface area than the forest gallery (1.24%), which is poorly represented. On the other hand, in 2021 the residential area (27.75%), the agricultural area (22.70%) followed by the grassy savannah occupied more land to the detriment of the forest gallery (0.47%) which remained the least represented.

Formatted: Font color: Auto


Formatted: Font color: Auto



AA: Agricultural area, FG: Forest gallery, TS: Tree savannah, SS: Shrub savannah, GS: Grassy savannah, BS: Bare soil, WS: Water surface, HyA: Hydromorphic area, HoA: Housing area

Figure 11: Areas of different land use units in the Dang Lake ecosystem


The Mballang lake ecosystem (Fig. 12) shows in 1990 a higher extent of land use units of the shrub savannah type (33.05%) followed by the grassy savannah (30.05%) and the agricultural area (17.27%). The smallest area in 1990 was noted in the hydromorphic zone (0.08%). The land use units in 2006 are dominated by grassy savannah (32.51 ha) followed by shrub savannah (32.05%) and agricultural area (16.57%). The forest gallery (0.64%) had the smallest area in 2006. In 2021, the grassy savannah (56.49%) followed by the agricultural area (21.65%) and the water area (11.89%). The hydromorphic zone (0.13%) had the smallest area in 2021.



AA: Agricultural area, FG: Forest gallery, TS: Tree savannah, SS: Shrub savannah, GS: Grassy savannah, BS: Bare soil, WS: Water surface, HyA: Hydromorphic area, HoA: Housing area

Figure 12: Areas of the different land-use units in the Mballang lake ecosystem

The most important land use unit at Lake Ngaoundaba (Fig 13) in 1990 was shrubsavannah (22.50%) followed by tree savannah (21.53%) and the hydromorphic zone (16.31%). In 2006, the grassy savannah (21.17%) followed by the shrub savannah (20.12%) and the treesavannah (19.37%) occupied the largest areas respectively. In 2021, the hydromorphic zone (28.13%) followed by shrub savannah (19.04%) and tree savannah (19.37%). In addition, the water surface occupied the smallest areas for the years 1990, 2006 and 2021, i.e. 2.35%;3.50% and 3.93% respectively.



AA: Agricultural area, FG: Forest gallery, TS: Tree savannah, SS: Shrub savannah, GS: Grassy savannah, BS: Bare soil, WS: Water surface, HyA: Hydromorphic area, HoA: Housing area

Figure 13: Areas of the different land use units of the Ngaoundaba lake ecosystem

2.5. Typology of the dynamics of change 1990 to 2021

Table 2 shows the diachronic analysis of the typology of the land use units of the lake ecosystems of Bini, Dang, Mballang and Ngaoundaba during the periods from 2021 to 1990; from 2021 to 2006 and from 2006 to 1990.

The analysis of this table 2 shows that in the Bini lake ecosystem, the agricultural space, the bare soil and the hydromorphic zone have evolved

progressively to the detriment of the forest gallery, the tree savannah, the shrub savannah and the water surface, which have gone into decline.

However, in the Dang lake ecosystem, the agricultural space, the tree savannah, the water surface and the housing area have progressively gone to the detriment of the forest gallery, the shrub savannah, the grassy savannah and the hydromorphic area. However, in the Mballang lake ecosystem, the agricultural area, the shrub savannah, the grassy savannah and the hydromorphic zone have progressively increased at the expense of the shrub savannah and the soil. On the other hand, in the Mballang lake ecosystem, the forest gallery and the water surface described a progressive dynamic exclusively during the periods 2021 to 2006 and regressive from 2006 to 1990. On the other hand, the Ngaoundaba lake ecosystem shows that the water surface and the hydromorphic zone record a progressive dynamic to the detriment of the tree and shrub savannah and the bare soil which are regressive. On the other hand, the forest gallery showed a progressive dynamic during the period 2021 to 2006.

Table 2: Change in the area of land use units of the lake ecosystems studied

	Land use unit	Evolution 2021-1990 (Ha)		Superficie 2021-2006 (Ha)		Superficie 2006-1990 (Ha)	
		Value	Trend	Value	Trend	Value	Trend
Lake de Bini	Agricultural space	24,40	Progressive	10,06	Progressive	14,34	Progressive
	Forestygallery	-3,53	Regressive	-0,11	Regressive	-3,42	Regressive
	Woodedsavannah	-11,20	Regressive	0,22	Progressive	11,42	Regressive
	Shrubbsavannah	-16,21	Regressive	-2,98	Regressive	13,22	Regressive
	Bareground	0,63	Progressive	1,92	Progressive	1,29	Progressive
	Water surface	-5,60	Regressive	-4,22	Regressive	-1,38	Regressive
	Hydromorphic zone	3,09	Progressive	-12,67	Regressive	15,75	Progressive
	Residential area	8,42	Progressive	7,78	Progressive	0,65	Progressive
Lake de Dang	Agricultural space	24,38	Progressive	12	Progressive	12,38	Progressive
	Forestygallery	-7,53	Regressive	-1,19	Regressive	-6,34	Regressive
	Woodedsavannah	3,49	Progressive	1,5	Progressive	1,99	Progressive
	Shrubbsavannah	-30,56	Regressive	-0,7	Regressive	29,86	Regressive
	Grasslandsavannah	-35,34	Regressive	-28,31	Regressive	-7,03	Regressive
	Bareground	-2,17	Regressive	0,8	Progressive	-2,97	Regressive
	Water body	8,48	Progressive	2	Progressive	6,48	Progressive
	Hydromorphic zone	-2,70	Regressive	-1,60	Regressive	-1,09	Regressive
Residential area	41,95	Progressive	15,51	Progressive	26,44	Progressive	
5 n	Agricultural space	16,72	Progressive	19,41	Progressive	-2,68	Regressive

Formatted	[... [1]
Formatted	[... [2]
Formatted	[... [4]
Formatted	[... [5]
Formatted	[... [6]
Formatted	[... [3]
Formatted	[... [7]
Formatted	[... [8]
Formatted	[... [9]
Formatted	[... [10]
Formatted	[... [11]
Formatted	[... [12]
Formatted	[... [13]
Formatted	[... [14]
Formatted	[... [15]
Formatted	[... [16]
Formatted	[... [17]
Formatted	[... [18]
Formatted	[... [19]
Formatted	[... [24]
Formatted	[... [25]
Formatted	[... [20]
Formatted	[... [21]
Formatted	[... [22]
Formatted	[... [23]
Formatted	[... [30]
Formatted	[... [31]
Formatted	[... [26]
Formatted	[... [27]
Formatted	[... [28]
Formatted	[... [29]
Formatted	[... [32]
Formatted	[... [33]
Formatted	[... [34]
Formatted	[... [35]
Formatted	[... [36]
Formatted	[... [37]
Formatted	[... [38]
Formatted	[... [39]
Formatted	[... [40]
Formatted	[... [41]
Formatted	[... [42]
Formatted	[... [43]
Formatted	[... [49]
Formatted	[... [44]
Formatted	[... [44]
Formatted	[... [45]
Formatted	[... [46]
Formatted	[... [47]
Formatted	[... [48]
Formatted	[... [50]
Formatted	[... [51]
Formatted	[... [52]
Formatted	[... [53]
Formatted	[... [54]
Formatted	[... [55]
Formatted	[... [61]
Formatted	[... [56]
Formatted	[... [57]
Formatted	[... [58]
Formatted	[... [59]

	Forestygallery	-0,12	Regressive	0,31	Progressive	-0,43	Regressive
	Woodedsavannah	10,17	Progressive	6,42	Progressive	3,75	Progressive
	Shrubbsavannah	-118,13	Regressive	114,80	Regressive	-3,33	Regressive
	Grasslandsavannah	97,78	Progressive	91,55	Progressive	6,23	Progressive
	Bareground	-6,62	Regressive	-4,84	Regressive	-1,78	Regressive
	Water surface	-0,88	Regressive	3,01	Progressive	-3,89	Regressive
	Hydromorphic zone	0,21	Progressive	0,09	Progressive	0,12	Progressive
	Residential area	0,87	Progressive	-1,13	Regressive	2,00	Progressive
Lake Ngaoundaba	Forestygallery	-5,24	Regressive	1,18	Progressive	-6,42	Regressive
	Woodedsavannah	-8,25	Regressive	-2,19	Regressive	-6,05	Regressive
	Shrubbsavannah	-9,67	Regressive	-3,01	Regressive	-6,65	Regressive
	Grasslandsavannah	-9,86	Regressive	-20	Regressive	10,14	Progressive
	Bareground	-4,38	Regressive	-1,03	Regressive	-3,35	Regressive
	Water surface	4,40	Progressive	1,21	Progressive	3,19	Progressive
	Hydromorphic zone	33,00	Progressive	23,84	Progressive	9,15	Progressive

2.6. Conversion and expansion rates of occupancy units

Table 3 presents the rate of conversion and expansion of land use units of the studied lake ecosystems during the periods 2021 to 1990; 2021 to 2006 and 2006 to 1990.

The rate of conversion and expansion at the Bini lake ecosystem (Table 3) is highest at the settlement area followed by the agricultural area and bare ground. However, the rate of conversion and expansion is lowest during the period 2021 to 1990 at the gallery forest unit followed by the shrub and tree savannah. The rate of conversion and expansion of land use units in the Dang Lake ecosystem on the other hand shows that during the periods 2021 to 1990; 2021 to 2006 and 2006 to 1990, the rate of conversion and expansion is highest at the settlement area followed by the agricultural area. These rates are lower in the forest gallery unit followed by the tree and shrub savannah.

The rate of conversion and expansion of land use units in the Mballang lake ecosystem (Table 3) shows that during the periods 2021 to 1990; 2021 to 2006 and 2006 to 1990, the rate of conversion and expansion is highest in the shrub savannah, the habitation area. In contrast, these rates are lower in the shrub savannah and bare ground units. The rate of conversion and expansion of the land use units of the Ngaoundaba lake ecosystem (Table 3) shows that during the periods 2021 to 1990; 2021 to 2006 and 2006 to 1990, the rate of conversion and

Formatted	... [118]
Formatted	... [119]
Formatted	... [120]
Formatted	... [121]
Formatted	... [122]
Formatted	... [123]
Formatted	... [124]
Formatted	... [125]
Formatted	... [126]
Formatted	... [127]
Formatted	... [128]
Formatted	... [129]
Formatted	... [132]
Formatted	... [130]
Formatted	... [131]
Formatted	... [133]
Formatted	... [134]
Formatted	... [135]
Formatted	... [136]
Formatted	... [137]
Formatted	... [138]
Formatted	... [139]
Formatted	... [140]
Formatted	... [141]
Formatted	... [142]
Formatted	... [143]
Formatted	... [144]
Formatted	... [145]
Formatted	... [146]
Formatted	... [147]
Formatted	... [148]
Formatted	... [149]
Formatted	... [150]
Formatted	... [151]
Formatted	... [152]
Formatted	... [153]
Formatted	... [154]
Formatted	... [155]
Formatted	... [156]
Formatted	... [157]
Formatted	... [158]
Formatted	... [159]
Formatted	... [160]
Formatted	... [161]
Formatted	... [162]
Formatted	... [163]
Formatted	... [164]
Formatted	... [165]
Formatted	... [166]
Formatted	... [167]
Formatted	... [168]
Formatted	... [169]
Formatted	... [170]
Formatted	... [171]
Formatted	... [172]
Formatted	... [173]
Formatted	... [174]
Formatted	... [175]
Formatted	... [176]

expansion is highest in the hydromorphic zone and the water surface. However, these rates are lower in the bare soil unit followed by the grassy savannah.

Table 3: Conversion and expansion rates of land use and land cover units in the Bini lake ecosystem

	Land use units	Conversion rate			Expansion rate		
		2021-1990	2021-2006	2006-1990	2021-1990	2021-2006	2006-1990
Lake Bini	Agricultural space	227,98	40,23	133,89	3,83	2,25	5,31
	Forestygallery	-82,92	-13,33	-80,29	-5,70	-0,95	-10,15
	Woodedsavannah	-80,38	8,80	-81,97	-5,25	0,56	-10,71
	Shrubbysavannah	-71,78	-31,89	-58,57	-4,08	-2,56	-5,51
	Bareground	36,03	146,84	-44,89	0,99	6,02	-3,72
	Water surface	-62,00	-55,38	-14,83	-3,12	-5,38	-1,00
	Hydromorphic zone	33,49	-22,99	73,34	0,93	-1,74	3,44
	Residential area	3863,30	900,00	296,33	11,87	15,35	8,61
Lake Dang	Agricultural space	227,98	52,02	115,75	3,83	2,79	4,81
	Forestygallery	-91,19	-62,10	-76,74	-7,84	-6,47	-9,12
	Woodedsavannah	54,85	11,28	39,15	1,41	0,71	2,06
	Shrubbysavannah	-93,80	-25,74	-91,65	-8,97	-1,98	-15,22
	Grasslandsavannah	-56,86	-51,36	-11,31	-2,71	-4,80	-0,75
	Bareground	-31,49	21,45	-43,59	-1,22	1,30	-3,88
	Water surface	56,94	9,35	43,52	1,45	0,60	2,46
	Hydromorphiczone	-22,92	-15,05	-9,26	-0,84	-1,09	-0,11
Residential area	4559,13	56,68	2873,59	12,39	2,99	21,10	
Lake Mbatlang	Agricultural space	25,37	41,90	-11,65	0,73	2,33	-0,77
	Forestygallery	-4,10	12,56	-14,80	-0,13	0,79	-1,00
	Woodedsavannah	242,08	80,72	89,29	3,97	3,95	3,99
	Shrubbysavannah	-94,37	-94,22	-2,64	-9,28	-19,00	-0,77
	Grasslandsavannah	82,95	73,78	5,28	1,95	3,68	0,22
	Bareground	-35,88	-29,98	-8,42	-1,43	-2,38	-0,53
	Water surface	-1,89	7,10	-8,40	-0,06	0,46	-0,55
	Hydromorphic zone	70,00	-90,59	1706,67	1,71	-15,76	18,99
Residential area	178,13	-45,48	410,17	3,30	-4,04	10,88	
a Forestygallery	-14,38	3,94	-17,62	-0,50	0,26	-1,11	

Formatted	...	[219]
Formatted	...	[220]
Formatted	...	[221]
Formatted	...	[222]
Formatted	...	[223]
Formatted	...	[224]
Formatted	...	[225]
Formatted	...	[226]
Formatted	...	[227]
Formatted	...	[228]
Formatted	...	[229]
Formatted	...	[230]
Formatted	...	[231]
Formatted	...	[232]
Formatted	...	[233]
Formatted	...	[234]
Formatted	...	[235]
Formatted	...	[236]
Formatted	...	[237]
Formatted	...	[238]
Formatted	...	[239]
Formatted	...	[240]
Formatted	...	[241]
Formatted	...	[242]
Formatted	...	[243]
Formatted	...	[244]
Formatted	...	[245]
Formatted	...	[246]
Formatted	...	[247]
Formatted	...	[248]
Formatted	...	[249]
Formatted	...	[250]
Formatted	...	[251]
Formatted	...	[252]
Formatted	...	[253]
Formatted	...	[254]
Formatted	...	[255]
Formatted	...	[256]
Formatted	...	[257]
Formatted	...	[258]
Formatted	...	[259]
Formatted	...	[260]
Formatted	...	[261]
Formatted	...	[262]
Formatted	...	[263]
Formatted	...	[264]
Formatted	...	[265]
Formatted	...	[266]
Formatted	...	[267]
Formatted	...	[268]
Formatted	...	[269]
Formatted	...	[270]
Formatted	...	[271]
Formatted	...	[272]
Formatted	...	[273]
Formatted	...	[274]
Formatted	...	[275]
Formatted	...	[276]
Formatted	...	[277]

Woodedsavannah	-13,72	-4,38	-9,77	-0,48	-0,30	-0,64
Shrubbsavannah	-15,39	-5,30	-10,66	-0,54	-0,36	-0,70
Grasslandsavannah	-20,13	-33,84	20,72	-0,73	-2,75	1,18
Bareground	-23,82	-7,19	-17,92	-0,88	-0,50	-1,23
Water surface	51,90	2,20	48,63	1,35	0,15	2,48
Zone hydromorphe	72,47	43,61	20,10	1,76	2,41	1,14

2.7. Extent of change in land use units


Figure 14 shows the extent of change in the land use units of the lake ecosystems of Bini (Fig 14 A) Dang (Fig 14 B) Mballang (Fig 14 C) and Ngaoundaba (Fig 14 D) from 1990 to 2021.

In the Bini ecosystem (Fig. 14 A), all land use units experienced significant change, both positive and negative. The agricultural area (1.16 Ha/year) followed by the residential area (0.40 Ha/year) experienced the highest acceleration. On the other hand, the lowest acceleration between 1990 and 2021 is observed in the forest gallery (0.17 Ha/year) followed by bare soil (0.03 Ha/year).


In the Dang ecosystem (Fig 14 B), the residential area (2.00 Ha/year) followed by the agricultural area (1.16 Ha/year) recorded the highest rates of change. On the other hand, the slowest rates of change were in the bare soil (-0.13 Ha/year) and the hydromorphic area (-0.10 Ha/year).

For the Mballang ecosystem (Fig 14 C), the grassy savannah (4.66 Ha/year) followed by the agricultural area (0.80 Ha/year) showed the highest rates of change. However, the lowest velocities were noted in the Forest Gallery (-0.01 Ha/year) and the water surface (-0.04 Ha/year) respectively.

For the Ngaoundaba ecosystem (Fig 14 D), the hydromorphic zone (1.57 Ha/year) followed by the water surface (0.21 Ha/year) recorded the highest velocities. On the other hand, the lowest velocities were recorded in the bare soil (-0.21 Ha/year) and the forest gallery (-0.25 Ha/year) respectively.



- Formatted ... [394]
- Formatted ... [395]
- Formatted ... [396]
- Formatted ... [397]
- Formatted ... [398]
- Formatted ... [399]
- Formatted ... [400]
- Formatted ... [401]
- Formatted ... [402]
- Formatted ... [403]
- Formatted ... [404]
- Formatted ... [405]
- Formatted ... [406]
- Formatted ... [407]
- Formatted ... [408]
- Formatted ... [409]
- Formatted ... [410]
- Formatted ... [411]
- Formatted ... [412]
- Formatted ... [413]
- Formatted ... [414]
- Formatted ... [415]
- Formatted ... [416]
- Formatted ... [417]
- Formatted ... [418]
- Formatted ... [419]
- Formatted ... [420]
- Formatted ... [421]
- Formatted ... [422]
- Formatted ... [423]
- Formatted ... [424]
- Formatted ... [425]
- Formatted ... [426]
- Formatted ... [427]
- Formatted ... [428]
- Formatted ... [429]
- Formatted ... [430]
- Formatted ... [431]
- Formatted ... [432]
- Formatted ... [433]
- Formatted ... [434]
- Formatted ... [435]
- Formatted ... [436]
- Formatted ... [437]
- Formatted ... [438]
- Formatted ... [439]
- Formatted ... [440]



AA: Agricultural area, FG: Forest gallery, TS: Tree savannah, SS: Shrub savannah, GS: Grassy savannah, BS: Bare soil, WS: Water surface, HyA: Hydromorphic area, HoA: Housing area

Figure 14: Rate of change of land use units in the lake ecosystems of Bini (A) Dang (B) Mballang (C) and Ngaoundaba (D)

2. Discussion

Spatio-temporal dynamics can be seen as a process of permanent occupation of a territory by various land use and occupation units in space and time (Reference). Diachronic analysis based on spot satellite images from 1990, 2006 and 2021 has enabled us to assess the evolution of the various landscape units over a period of thirty-one years. The overall statistical accuracy of the classified images ranges from 90.23 to 97.14% with kappa indices varying from 86.93 to 95.65%. In the present study, the Kappa index is above 50% and the lowest value is 88%. In the study of land cover, when the Kappa index evaluated in the classification operations is between 50 and 75%, the adopted classification is valid and the results can be used judiciously (Pontius, 2000). The different evaluation results of the images classified by the supervised classification method are therefore validated and could be explained by the ability of the sensors to generate good images during low cloud cover. These results could be explained by the ability of these environments to allow several elements to coexist within the ecosystem that they constitute. These results are close to 98.86% with kappa indices varying from 85.65 to 97.35% on the spatiotemporal dynamics of land cover in three protected areas in Burkina Faso. those of (recase) Moussa Ganame, (2021) who reported overall statistical accuracies of classified images ranging from 88.90% to 98.86% with kappa indices varying from 85.65 to 97.35% on the spatiotemporal dynamics of land cover in three protected areas in Burkina Faso. (recase)

The fairly large number of land use and occupancy unit classes in the four lake ecosystems (9 classes in Mballang, 9 classes in Dang, 8 classes in Bini and 7 classes in Ngaoundaba) shows a diversity in the lake ecosystem landscape. These units are not always well differentiated, probably because of the close spectral

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Red

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font: Not Bold, Font color: Red

Formatted: Font: Not Bold, Font color: Red, Strikethrough

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font: Not Bold, Font color: Red

Formatted: Font color: Red

Formatted: Font: Not Bold, Font color: Auto

responses of these natural and particularly woody plant formations. These difficulties have been encountered by several authors and even in other countries (Avakoudjo *et al.*, 2014; Mamane *et al.*, 2018). These difficulties could be related to the homogeneous plots when choosing training sites. However, despite these difficulties, the results obtained remain exploitable. Similar results, which discriminate a maximum of 9 land use and occupation unit classes on the banks of the Gbaga Lagoon in West Africa, were also reported by Ahehinnou *et al.* (2020).

The estimation of NDVI from the maps of spatiotemporal dynamics reveals a regression of the vegetation of lake ecosystems. The recorded NDVI values of the lake ecosystems varied from 0 to 0.214; from 0 to 0.214; from -0.116 to 0.326 and from -0.06 to 0.282 respectively for the lake ecosystems of Bini, Dang, Mballang and Ngaoundaba. The larger NDVI differences at the Mballang (-0.116 to 0.326) and Ngaoundaba (-0.06 to 0.282) lake ecosystems could be explained by the variation in vegetation density and edaphic parameters across these ecosystems. The authors revealed that vegetation could significantly reduce temperature if NDVI exceeded 0.35 (Mackey *et al.*, 2012). In our study, the results show a strong decrease in vegetation over time. Consequently, the reduction of vegetation cover increases the soil surface temperatures of lake ecosystems.

The evolution of land use and land cover units shows changes in the area of land use units in general and in vegetation cover in the years 1990, 2006 and 2021. In general, in 1990 the savannahs (shrub savannah, tree savannah and grass savannah) are the most extensive units. In 2006 and 2021, agricultural space, housing area, bare soil and hydromorphic area are gaining ground to the detriment of savannah and forest gallery. These results could be explained by the growth in demography and the increase in poverty, which leads respectively to the extension of agricultural areas and the search for habitable areas with low land value. The expansion of these units has encroached on areas previously occupied by natural formations such as savannahs and gallery forests. These results confirm the findings of Gauze *et al.* (2019) on Characterisation of land use dynamics and morphology of the Aby Lagoon in the Ehotile Islands National Park area; South-East of Côte d'Ivoire, Bamba *et al.* (2008) on Influence of anthropogenic actions on the spatio-temporal dynamics of land use in the Bas-Congo province (D.R.Congo).

Spatial and temporal dynamics are expressed in terms of regression, stability or progression of land-use units, hence the notion of typology. The typology of the dynamics of change from 1990 to 2021 shows that the agricultural space, the housing area, bare soil and the hydromorphic zone describe a progressive dynamic to the detriment of the forest gallery, the wooded savannah, the shrubby savannah and the water surface which have gone into regression. These results would be the result of strong pressures on the resources of the banks of the studied lake ecosystems for the satisfaction of different needs. The wetlands as a space with a high potential for life has favoured the immigration and settlement of people from

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

dry areas for grazing, watering animals and fishing. This situation leads to the uncontrolled occupation of lake ecosystems by people who largely encroach on the riparian vegetation, which provides enormous ecosystem services. These results are similar to those of Tente (2005) who worked in the forest of Eastern Cameroon and showed that other types of vegetation formation are progressing while others are regressing. This also confirms well the results of Mama *et al.* (2003), Tente (2005), Orékan (2007) and Arouna (2012) concerning the regression of dense vegetation formations in favour of agricultural spaces.

The rate of conversion and expansion of the land use units of the lake ecosystems studied shows in general that during the periods from 2021 to 1990, the rate of conversion and expansion is higher in the residential area followed by the agricultural area. This clear difference in the rate of conversion and expansion of the land use units of the four lake ecosystems studied could be explained by the fact that anthropogenic action is more likely to be directed towards agriculture and urbanisation, while neglecting the environmental aspect. Agriculture is the main source of deforestation. Similarly, similar results have been observed in several other localities in Cameroon (Solefack *et al.*, 2018; Temgoua, *et al.*, 2018b).

In general, the speed of the dynamics over the last 31 years has revealed progressive degradation in the grassy savannah, the residential area, the agricultural area and the hydromorphic area. These degradations have been at the expense of the shrub savannah, the tree savannah and the forest gallery. These plant formations have been degraded at a considerable rate. The causes of the degradation of the vegetation cover are anthropogenic and can also be climatic when the ecological zone does not benefit from the minimum rainfall that should allow the spontaneous reconstitution of plant formations, which is increasingly observed with climate change (Adjonou *et al.*, 2010). In addition, agricultural activities are the major causes of vegetation degradation (Nguimdo, 2017; Solefack *et al.*, 2018; Temgoua *et al.*, 2018b). However, this regression of shrubby savannahs, trees and forest gallery would be accompanied by the loss of biodiversity and land degradation.

Conclusion and recommendations

The study of the spatiotemporal dynamics of land use in four lake complexes in the Vina Department (Adamaoua-Cameroon) showed that the natural plant formations (shrubby savannah, trees and forest gallery) have undergone a strong regressive dynamic to the benefit of anthropogenic formations (the residential area and the agricultural space). The regression of these vegetation formations on the banks is due to several factors, the most important of which are population growth,

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

the extension of cultivated areas and logging. This destruction of plant cover leads to an imbalance in the natural resource cycle, soil degradation and, above all, the loss of ecosystem services and biodiversity.

Formatted: Font color: Auto

This should be recommended based on the findings of the study: the following recommendations are to be noted then you itemize them

These results call on the various actors or decisionmakers in charge of wetland management to consider restoration methods. This makes clear the urgent need to establish integrated and participatory management strategies at both local and regional levels to preserve and control lake ecosystems more effectively. They also contribute to a better knowledge of the predictions on the evolution of these ecosystems expected from different socio-economic scenarios and to ensure resilience.

Formatted: Font color: Red

REFERENCES

Agbanou B. T., Orekan V., Abdoulaye D., Martin P., Tente B. 2018. Spatio-temporal dynamics of land use in extensive agriculture: the case of the Natitingou-Boukoubesector in northwestern Benin. Mixtures in homage to Professors Thomas Omer, Houssou Sègè Christophe and Houndaga Cossi Jean. *La géographie au service du développement durable*, Abomey Calavi, Benin. 22-34pp. hal-02092545.

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Ahehinnou Y. M. F., Amoussou E., Allagbe B. S. Y., Vissin E. W., 2020. Spatio-Temporal Dynamics of the Occupation of the Banks of the Gbaga Lagoon (West Africa). *International Journal of Progressive Sciences and Technologies*. 22 (2), 235-243pp.

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Arouna O., 2012. Mapping and predictive modelling of spatio-temporal changes in vegetation in the Commune of Djidja, Benin: implications for land use planning. Unique PhD thesis, Geography and Environmental Management. EDP/FLASH/UAC, Benin, 246p.

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Avakoudjo J, Mama A, Toko I, Kindomhou V, Sinsin B. 2014. Land use dynamics in the National Park and its periphery in northwest Benin. *Int. J. Biol. Chem. Sci*, 8(6): 2608- 2625pp.

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Bamba I, Mama A., Neuba D. F. R., Koffi K. J., Traoré D., Visser M., Sinsin B., Lejoly J. and Bogaert J., 2008 "Influence des actions anthropiques sur la dynamique spatiotemporelle de l'occupation du sol dans la province du Bas-Congo (R.D. Congo)" *Sciences & Nature* 5 (1) 49-60pp

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Bastviken, D., Tranvik, L. J., Downing, J. A., Crill, P. M. & Enrich-Prast, A., 2011. Freshwater methane emissions offset the continental carbon sink. *Science* 331, 50p.

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Bernier B., 1992. Introduction to macroeconomics. Paris, Dunod, France, 217p.

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font color: Auto

Chaplin-Kramer R., Sharp R.P., Mandle L., Sim S., Johnson J., Butnar I., Mila C.L., Eichelberger B.A., Ramler I., Mueller C., McLachlan N., Yousefi A., King H., & Kareiva P.M., 2015. Spatial patterns of agricultural expansion determine impacts on biodiversity and carbon storage. *Proceedings of the National Academy of Sciences of the United States of America*, 112: 7402-7407pp.

Formatted: Font: Not Bold, Font color: Auto

De Graaf, G., Bartley, D., Jorgensen, J. & Marmulla, G. 2012. The scale of inland fisheries, can we do better? Alternative approaches for assessment. *Fisheries Manag. Ecol.* 1, 64-70p.

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Dodson, S. I., Arnott, S. E. & Cottingham, K. L. 2000. The relationship in lake communities between primary productivity and species richness. *Ecology* 81, 2662-2679p.

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Downing, J. A. et al. 2008. Sediment organic carbon burial in agriculturally eutrophic impoundments over the last century. *Global Biogeochem. Cycles* 22, 10-18pp.

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

FAO, 2011. State of the World's Forests 2011.

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Gauze TouaoKah Martine, KOUASSI Kouakou Lazare and MALAN Djah François, 2019. "Characterisation of the land use dynamics and morphology of the Aby lagoon in the space of the Ehotile Islands National Park; South-East of Côte d'Ivoire", *European Scientific Journal*, 15, 11-26pp.

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Halima G. and Djamel A. 2020. Estimation of the impact of green spaces and water surfaces on urban climate and soil surface temperature (mila, algeria). *Rev. Roum. Geogr./Rom. Journ. Geogr.* 64, (2), 155-174pp.

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Inoussa M.M., Mahamane A., Mbow C., Saadou M., & Yvonne B., 2011. Spatiotemporal dynamics of open forests in the W National Park of Niger (West Africa). *Sécheresse*, 22 :108-116pp.

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Leimgruber L., Kelly D.S., Steininger M.K., Brunner J., Müller T., & Songer M., 2005. Forest cover change patterns in Myanmar (1990-2000). *Environmental Conservation*, 32p.

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Mackey, Christopher, Lee, Xuhui, Smith, Ronald (2012). Remotely sensing the cooling effects of city scale efforts to reduce urban heat island, *Building and Environment*, 49, 348-358 pp.

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Mama A., Sinsin B., De C.C., & Bogaert J., 2013. Anthropisation and landscape dynamics in the Sudanian zone of northern Benin. *Tropicultura*, 31(1) :78-88pp.

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Mamane B, Amadou G, Barrage M, Comby J, Ambouta JMK. 2018. Spatiotemporal dynamics of land use in the Tamou Total Wildlife Reserve in a

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

context of climatic variability (Western Niger). *Int. J. Biol. Chem. Sci.* 12(4): 1667-1687pp.

Messenger L. M., Lehner B., Grill G., Nedeva I. & Schmitt O., 2016, Estimating the volume and age of water stored in global lakes using a geo-statistical approach. *Nature communications*, DOI: 10.1038/ncomms13603.

Moussa Ganame, 2021. Spatio-temporal dynamics and potential of the aerial carbon stock of forest ecosystems in Burkina Faso. Doctorate from the University Joseph KI-ZERBO. 167p.

Muller Schmied, H. et al, 2014. Sensitivity of simulated global-scale freshwater fluxes and storages to input data, hydrological model structure, human water use and calibration. *Hydrol. Earth. Syst. Sci.* 18, 3511-3538p.

Nguimdo V. 2017. Dynamics of deforestation, degradation and carbon stocks in the Teaching and Research Forest of the University of Dschang in BÉlabo, East Cameroon. Dissertation of Engineer of Water, Forestry and Hunting Design, University of Dschang, 79p

Norges, T. 2009. Relationships between morphometry, geographic location and water quality parameters of European lakes. *Hydrobiologia* 633, 33-43pp.

Orekan V.O.A., 2019. Dynamics of land use in the Savalou classified forest in Benin. *Regard Sud*. 19p.

Post, D. M., Pace, M. L. & Hairston, N. G. 2000. Ecosystem size determines foodchain length in lakes. *Nature* 405, 1047-1049pp.

Solefack MC, Njouonkou AL, Temgoua LF, Djouda R, Zangmene JB, Ntoupka M. 2018. Land-Use/ Land-Cover change and Anthropogenic Causes Around Koup Matapit Gallery Forest, West-Cameroon. *Journal of Geography and Geology*, 10(2): 201-219pp.

Staehr, P. A., Baastrup-Spohr, L., Sand-Jensen, K. & Stedmon, C. 2012. Lake metabolism scales with lake morphometry and catchment conditions. *Aquat. Sci.* 74, 155-169p.

Staehr, PA, Baastrup-Spohr, L., Sand-Jensen, K. & Stedmon, C. 2012. Lake metabolism changes with lake morphometry and catchment conditions. *Aquat. Sci.* 74, 155-169p.

Temgoua LF, Ajonina G, Woyu HB. 2018a. Land Use and Land Cover Change Analysis in Ajei Upland Waterched Community Forest, North West Region, Cameroon. *Journal of Geoscience and Environment Protection*, 6(09): 83-99pp.

Tente B., 2005: Recherche sur les facteurs de la diversité floristique des versants du massif de l'Atacora: Secteur Perma-Toucountouna (Bénin). Thèse de doctorat unique. EDP/FLASH/UAC, 252 p.

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Tilahun A., & Islam Z., 2015. Google earth for Land Use Land Cover Change detection in the case of Gish AbbaySekela, West Gojjam, Amhara State, Ethiopia. *International Journal of Advancement in Remote Sensing, GIS and Geography*, 3 (2):80-87pp.

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Touchart L., 2002. Limnologie physique et dynamique: une géographie des lacs et des étangs. Harmattan, Paris, p.60

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Vittek M., Brink A., Donnay F., Simonetti D., & Desclée B., 2014. Land Cover Change Monitoring Using Landsat MSS/TM Satellite Image Data over West Africa between 1975 and 1990. *Remote Sensing*, 6: 658-676pp.

Formatted: Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

UNDER PEER REVIEW

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

8/22/2022 6:54:00 AM

