

## Original Research Article

### **Effect of integrated nutrient management on the growth, yield and quality in tomato (*Solanum lycopersicum* L.).**

---

#### **ABSTRACT**

The present investigation was carried out with title Effect of integrated nutrient management on the growth, yield and quality in tomato (*Solanum lycopersicum* L.) at the Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj, Uttar Pradesh during the Kharif 2021-22 with a view to determine the effect of plant growth regulators application on tomato variety TMTM 2267 for its growth, quality and yield and to work out the economics of various treatments. Under This experiment, overall 10 treatment was taken T<sub>0</sub> (RDF of NPK(150:80:60)Kg/ha +10t FYM), T<sub>1</sub> (75% N through Urea +MOP + SSP +25% through FYM + Boron + Azotobactor +Phosphorous Solubilizing Bacteria), T<sub>2</sub> (75% N through Urea +MOP + SSP +25% through FYM +Zinc+ Azotobactor + Phosphorous Solubilizing Bacteria), T<sub>3</sub> (75% N through Urea + MOP + SSP +25% through FYM + Boron+ Zinc+ Azotobactor + Phosphorous Solubilizing Bacteria), T<sub>4</sub> (75% N through Urea + MOP + SSP + 25% through vermicompost+ B + Azotobactor + Phosphorous Solubilizing Bacteria), T<sub>5</sub> (75% N through Urea + MOP + SSP + 25% through vermicompost + Zinc + Azotobactor + Phosphorous Solubilizing Bacteria), T<sub>6</sub> (75% N through Urea+ MOP + SSP + 25% through vermicompost+ Boron + Zinc + Azotobactor + Phosphorous Solubilizing Bacteria), T<sub>7</sub> (75% N through Urea + MOP + SSP + 25% through poultry manure + Boron + Azotobactor + Phosphorous Solubilizing Bacteria), T<sub>8</sub> (75% N through Urea + MOP + SSP + 25% through poultry manure +Zinc + Azotobactor + Phosphorous Solubilizing Bacteria) and T<sub>9</sub> (75% N through Urea + MOP + SSP + 25% through poultry manure + Boron + Zinc + Azotobactor + Phosphorous Solubilizing Bacteria).. From the present investigation it was concluded that T<sub>1</sub> (75% N through Urea +MOP + SSP +25% through FYM + Boron + Azotobactor +PSB) was found to be best among all treatment in terms of yield and T<sub>6</sub> (75% N through Urea+ MOP + SSP + 25% through Vermicompost+ B + Zn + Azotobactor +PSB) in growth and quality of tomato i.e. plant spread, plant height, T.S.S. etc. The highest B:C ratio that is 3.25 was observed for T<sub>6</sub> (75% N through Urea+ MOP + SSP + 25% through Vermicompost+ B + Zn + Azotobactor +PSB).

*Keywords: Integrated Nutrient Management, PSB, Azotobactor, Benefit cost ratio.*

---

## INTRODUCTION

Tomatoes are horticulture crop belongs to the family Solanaceae bearing chromosome number  $2n=2X=24$  (Karpechenko, 1925). It originated from South America (Vavilov, 1935). The tomato plants typically grow to 1–3 meters (3–10 ft) in height and have a weak stem that often sprawls over the ground and vines over other plants. Flowers are generally borne in clusters of 4 to 8 but small fruited types may have 30 to 50 flowers per cluster. Tomato plants are dicots, and grow as a series of branching stems, with a terminal bud at the tip that does the actual growing. Tomato plays a major role in human nutrition, fruit contain 93.1% water, 1.9% protein, 0.3 g fat, 0.7% fibre, 3.6% carbohydrates, 23 calorie, 320 I.U vitamin A., 0.07 mg vitamin B1, 0.01 mg vitamin B2, 31 mg vitamin C, 20 mg calcium, 36 mg phosphorus and 0.8 mg iron.

### **Integrated Nutrient Management (INM):-**

Integrated nutrient management (INM) is an approach that involves the management of both organic and inorganic plant nutrients for optimal production of cultivated crops, forage, and tree species, while conserving the natural resource base essential for long-term sustainability. The crop is being cultivated as an important spring summer season vegetable in eastern Uttar Pradesh finding readymade market in plains of northern India fetching very remunerative prices for the farmers.

The growth, yield and fruit quality of tomato largely depend on number of various interacting factors. Among them, INM is the most crucial as well as basic factor. The continuous use of chemical fertilizer increases the concentration of heavy metals in the soil, disturbs soil health and quality which cannot support plant growth in long term basis. Integrated Nutrient Management comprises organic, inorganic component and microorganism that are highly beneficial for sustainable crop production as it ameliorates soil environment, maintains adequate level of nutrients and provides favourable conditions for high tomato yield with desired quality. The integrated use of organic and inorganic fertilizers is the need of hour and is being advocated for sustainable agriculture. When the inorganic fertilizers are not available timely due to higher prices and inadequate supply of it, organic manures can supplement the nutrients.

Nitrogen (N) plays a key role in nutrition of the plants. As a matter of fact, the plant life would not be possible without this element. Adequate amount of nitrogen are also required to

obtain good yield in vegetable crops. Phosphorous and potassium is considered as major nutrient in tomato cultivation which involves in all the metabolic process in the plant and there is considerable evidence to show that, this element plays an important role in photosynthesis and helps in building up of carbohydrate in the plant. The production of dry matter is further affected by the effect of potassium on rate of respiration.

The role of bio-fertilizers in improving soil fertility has long been studied in various crops. The bio-fertilizers such as Azotobactor, Phosphate Solubilising Bacteria and Arbuscular Mycorrhiza (AM fungi) helps to enhance overall soil fertility by modifying soil texture, soil structure integrity, aeration, increased nutrient availability there by greatly influencing plant growth and yield.

Azospirillum a diazotropic bacterium which is widely distributed in soil rhizosphere and roots of a number of plants have ability to fix nitrogen. Many reports have indicated the importance of Azospirillum in vegetable crops.

Bio-fertilizers are involved in symbiotic and associative microbial activities with higher plants. These are natural mini-fertilizer factories that are economical and safer source of plant nutrition for increasing the agricultural production and improving fertility.

## **MATERIALAND METHODS**

The present investigation entitled “Effect of integrated nutrient management on the growth, yield and quality in tomato (*Solanum lycopersicum* L.)” was done to understand the effect of Integrated nutrient management at different doses combination on fruit growth, yield and quality of tomato variety TMTH-2267. The details of the materials used and the procedures adopted in the investigation, which was carried out at Horticultural Research Farm (HRF), Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Prayagraj during the Kharif season of 2021 are described under the following heads.

### **Experimental Site**

The present investigation was carried out in the Horticultural Research Farm at Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, U.P during Kharif-2021. The University is situated on the left side of Prayagraj – Rewa National Highway, about 5 km away from Prayagraj City. All types of facilities necessary for cultivation of successful crop including field preparation, inputs, irrigation facilities and labours were provided from the Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, U.P.

## **Experimental Materials**

Variety details: TMTH-2267

VarietySource - Department of Horticulture, NAI, SHUATS

## **RESULTS AND DISCUSSION**

### **Growth Parameters**

#### **Plant height at 30 DAS**

The height of plant significantly varied among different treatment combinations. The mean Plant height at 30 DAS was observed to be 32.68 cm. The maximum plant height (35.69 cm) at 30 DAS was observed with treatment T<sub>8</sub> (75% N through Urea + MOP + SSP + 25% through Poultry Manure + Zinc + Azotobactor + Phosphorous Solubilizing Bacteria) followed by T<sub>3</sub> (75% N through Urea + MOP + SSP + 25% through FYM + Boron + Zinc + Azotobactor + Phosphorous Solubilizing Bacteria) with 33.76 cm. Minimum plant height (31.57 cm) was observed in T<sub>5</sub> (75% N through Urea + MOP + SSP + 25% through Vermicompost + Zinc + Azotobactor + Phosphorous Solubilizing Bacteria), while the remaining treatments were moderate in their growth habit.

#### **Plant height at 60 DAS**

The maximum plant height (73.60 cm) at 60 DAS was observed with treatment T<sub>9</sub> (75% N through Urea + MOP + SSP + 25% through Poultry Manure + Boron + Zinc + Azotobactor + Phosphorous Solubilizing Bacteria) followed by T<sub>6</sub> (75% N through Urea + MOP + SSP + 25% through Vermicompost + Boron + Zinc + Azotobactor + Phosphorous Solubilizing Bacteria) with 73.36 cm. Minimum plant height (52.85 cm) was observed in T<sub>0</sub> (RDF of NPK(150:80:60)Kg/ha + 10t FYM), while the remaining treatments were moderate in their growth habit.

#### **Plant height at 90 DAS**

The maximum plant height (101.05 cm) at 90 DAS was observed with treatment T<sub>5</sub> (75% N through Urea + MOP + SSP + 25% through Vermicompost + Zinc + Azotobactor + Phosphorous Solubilizing Bacteria) followed by T<sub>6</sub> (75% N through Urea + MOP + SSP + 25% through vermicompost + Boron + Zinc + Azotobactor + Phosphorous Solubilizing Bacteria) with 100.58 cm. Minimum plant height (79.20 cm) was observed

in T<sub>0</sub> (RDF of NPK(150:80:60)Kg/ha +10t FYM), while the remaining treatments were moderate in their growth habit.

### Plant height at harvest

The maximum plant height (103.99 cm) at harvest was observed with treatment T<sub>5</sub> (75% N through Urea + MOP + SSP + 25% through vermicompost + Zinc + Azotobactor + Phosphorous Solubilizing Bacteria) followed by T<sub>6</sub> (75% N through Urea+ MOP + SSP + 25% through vermicompost+ Boron + Zinc + Azotobactor + Phosphorous Solubilizing Bacteria) with 103.05 cm. Minimum plant height (82.88 cm) was observed in T<sub>0</sub> (RDF of NPK(150:80:60)Kg/ha +10t FYM), while the remaining treatments were moderate in their growth habit.

**Table 1. Effect of integrated nutrient management on plant height (cm) [30 DAS, 30, 45,60 DAT] and at harvest of the tomato.**

Treatments	Treatment Combination	30 DAS	60 DAS	90 DAS	At Harvest
T <sub>0</sub>	RDF of NPK(150:80:60)Kg/ha +10tn FYM	31.69	52.85	79.20	82.88
T <sub>1</sub>	75% N through Urea +MOP + SSP +25% through FYM + Boron + Azotobactor +PSB	33.41	59.91	87.90	91.64
T <sub>2</sub>	75% N through Urea +MOP + SSP +25% through FYM +Zn+ Azotobactor +PSB	31.55	59.14	91.41	95.13
T <sub>3</sub>	75% N through Urea + MOP + SSP +25% through FYM + B+ Zn+ Azotobactor +PSB	33.76	60.31	91.55	95.60
T <sub>4</sub>	75% N through Urea + MOP + SSP + 25% through Vermicompost+ B + Azotobactor +PSB	33.25	73.29	98.69	102.32
T <sub>5</sub>	75% N through Urea + MOP + SSP + 25% through Vermicompost+ Zn + Azotobactor + PSB	31.57	72.80	100.58	103.99
T <sub>6</sub>	75% N through Urea+ MOP + SSP + 25% through Vermicompost+ B + Zn + Azotobactor +PSB	32.27	73.16	100.58	103.58
T <sub>7</sub>	75% N through Urea + MOP + SSP + 25% through PM + B + Azotobactor + PSB	31.88	70.75	97.96	100.78

<b>T<sub>8</sub></b>	75% N through Urea + MOP + SSP + 25% through PM +Zn + Azotobactor + PSB	35.69	71.68	99.55	102.97
<b>T<sub>9</sub></b>	75% N through Urea + MOP + SSP + 25% through PM + B + Zn + Azotobactor +PSB	31.66	73.60	93.25	96.80
	<b>'F' Test</b>	<b>s</b>	<b>s</b>	<b>s</b>	<b>s</b>
	<b>C.V.</b>	<b>2.56</b>	<b>5.96</b>	<b>8.57</b>	<b>9.05</b>
	<b>S.E. (m)</b>	<b>0.86</b>	<b>2.01</b>	<b>2.88</b>	<b>3.05</b>
	<b>C.D. at 5%</b>	<b>4.57</b>	<b>5.21</b>	<b>5.31</b>	<b>5.41</b>

### Number of branches per plant

The number of branches per plant was influenced by different treatments at all successive stage of growth .It was found that T<sub>0</sub> (RDF of NPK(150:80:60)Kg/ha +10tn FYM) with maximum value i.e. 13.71 branches per plant followed by T<sub>1</sub> (75% N through Urea +MOP + SSP +25% through FYM + Boron + Azotobactor +PSB) with 12.36 branches per plant whereas the minimum score was observed in treatment T<sub>4</sub> (75% N through Urea + MOP + SSP + 25% through Vermicompost + B + Azotobactor +PSB) with 6.82.

### Plant Spread (cm)

It is also evident that the plant spread was influenced by different treatments at all successive stage of growth . It was also found that T<sub>6</sub> (75% N through Urea+ MOP + SSP + 25% through Vermicompost + B + Zn + Azotobactor +PSB) with maximum value i.e. 52.30 plant spread followed by T<sub>5</sub> (75% N through Urea + MOP + SSP + 25% through Vermicompost + Zn + Azotobactor + PSB) with 50.94 plant spread whereas the minimum score was observed in treatment T<sub>0</sub> (RDF of NPK(150:80:60)Kg/ha +10tn FYM) with 45.68.

### Days to first flowering

The results related to days to first flowering varied significantly among different treatments . Among the application of integrated nutrient management the maximum days to first flowering was seen in T<sub>8</sub> (75% N through Urea + MOP + SSP + 25% through PM +Zn + Azotobactor + PSB) with 42.88 days, followed by T<sub>3</sub> (75% N through Urea + MOP + SSP +25% through FYM + B+ Zn+ Azotobactor +PSB) with 41.31 days whereas minimum days to first flowering 30.65 days was recorded in T<sub>7</sub> (75% N through Urea + MOP + SSP + 25% through PM + B + Azotobactor + PSB).

### **Days to 1<sup>st</sup> fruit setting**

Among the application of integrated nutrient management the maximum days to first fruit setting was seen in T<sub>8</sub> (75% N through Urea + MOP + SSP + 25% through PM +Zn + Azotobactor + PSB) with 58.59 days, followed by T<sub>3</sub> (75% N through Urea + MOP + SSP +25% through FYM + B+ Zn+ Azotobactor +PSB) with 56.95 days whereas minimum days to first fruit setting 46.02 days was recorded in T<sub>7</sub> (75% N through Urea + MOP + SSP + 25% through PM + B + Azotobactor + PSB).

### **Number of fruits per plant**

The maximum number of fruits per plants 59.22 were recorded in treatment T<sub>0</sub> (RDF of NPK(150:80:60)Kg/ha +10tn FYM) followed by T<sub>1</sub> (75% N through Urea +MOP + SSP +25% through FYM + Boron + Azotobactor +PSB) i.e., 54.82 and the lowest fruits per plant 32.14 were observed in T<sub>9</sub> (75% N through Urea + MOP + SSP + 25% through PM + B + Zn + Azotobactor +PSB).

### **Days to 1<sup>st</sup> fruit picking**

Among the application of integrated nutrient management the maximum days to first fruit picking was seen in T<sub>8</sub> (75% N through Urea + MOP + SSP + 25% through PM +Zn + Azotobactor + PSB) with 71.37 days, followed by T<sub>3</sub> (75% N through Urea + MOP + SSP +25% through FYM + B+ Zn+ Azotobactor +PSB) with 69.60 days whereas minimum days to first fruit picking 58.75 days was recorded in T<sub>7</sub>

### **Fruit yield per plant (kg/plant)**

The maximum fruit yield per plants 3.44 kg/plant were recorded in treatment T<sub>1</sub> (75% N through Urea +MOP + SSP +25% through FYM + Boron + Azotobactor +PSB) followed by T<sub>0</sub> (RDF of NPK(150:80:60)Kg/ha +10tn FYM) i.e., 2.91 kg/plant and the lowest fruits per plant 1.81 were observed in T<sub>9</sub> (75% N through Urea + MOP + SSP + 25% through PM + B + Zn + Azotobactor +PSB).

**Table 2 : Effect of integrated nutrient management on fruit yield per plant (Kg/plant) of tomato.**

<b>Treatment Notation</b>	<b>Treatment Combination</b>	<b>Fruit yield per plant (kg/plant)</b>
<b>T<sub>0</sub></b>	RDF of NPK(150:80:60)Kg/ha +10tn FYM	2.91
<b>T<sub>1</sub></b>	75% N through Urea +MOP + SSP +25% through FYM + Boron + Azotobactor +PSB	3.44
<b>T<sub>2</sub></b>	75% N through Urea +MOP + SSP +25% through FYM +Zn+ Azotobactor +PSB	1.98
<b>T<sub>3</sub></b>	75% N through Urea + MOP + SSP +25% through FYM + B+ Zn+ Azotobactor +PSB	2.81
<b>T<sub>4</sub></b>	75% N through Urea + MOP + SSP + 25% through Vermicompost + B + Azotobactor +PSB	1.65
<b>T<sub>5</sub></b>	75% N through Urea + MOP + SSP + 25% through Vermicompost + Zn + Azotobactor + PSB	2.83
<b>T<sub>6</sub></b>	75% N through Urea+ MOP + SSP + 25% through Vermicompost + B + Zn + Azotobactor +PSB	2.18
<b>T<sub>7</sub></b>	75% N through Urea + MOP + SSP + 25% through PM + B + Azotobactor + PSB	2.38
<b>T<sub>8</sub></b>	75% N through Urea + MOP + SSP + 25% through PM +Zn + Azotobactor + PSB	2.23
<b>T<sub>9</sub></b>	75% N through Urea + MOP + SSP + 25% through PM + B + Zn + Azotobactor +PSB	1.81
	<b>Mean</b>	<b>2.43</b>
	<b>‘F’ Test</b>	<b>S</b>
	<b>C.V.</b>	<b>6.51</b>
	<b>S.E. (m)</b>	<b>0.09</b>
	<b>C.D. at 5%</b>	<b>0.27</b>

### **Total soluble solids (0Brix)**

The maximum T.S.S. (6.04 0Brix) was observed in treatment T<sub>6</sub> (75% N through Urea+ MOP + SSP + 25% through Vermicompost + B + Zn + Azotobactor +PSB) followed by T<sub>7</sub> (75% N through Urea + MOP + SSP + 25% through PM + B + Azotobactor + PSB) with 5.74 0Brix. The minimum T.S.S. (4.25 0Brix) was noticed in treatment T<sub>1</sub> (75% N through Urea +MOP + SSP +25% through FYM + Boron + Azotobactor +PSB).

### **Acidity (%)**

The maximum acidity (15.62%) was observed in treatment T<sub>0</sub> (RDF of NPK(150:80:60)Kg/ha +10tn FYM) followed by T<sub>3</sub> (75% N through Urea + MOP + SSP +25% through FYM + B+ Zn+ Azotobactor +PSB) with 14.66%. The minimum acidity (9.58%) was noticed in treatment T<sub>9</sub> (75% N through Urea + MOP + SSP + 25% through PM + B + Zn + Azotobactor +PSB).

## **Summary and Conclusion**

From the present investigation it was concluded that T<sub>1</sub> (75% N through Urea +MOP + SSP +25% through FYM + Boron + Azotobactor + Phosphorous Solubilizing Bacteria) was found to be best among all treatment in terms of yield and T<sub>6</sub> (75% N through Urea+ MOP + SSP + 25% through vermicompost+ Boron + Zinc + Azotobactor + Phosphorous Solubilising Bacteria) in growth and quality of tomato i.e. plant spread, plant height, T.S.S. etc. The highest B:C ratio that is 3.25 was observed for T<sub>6</sub> (75% N through Urea+ MOP + SSP + 25% through vermicompost+ Boron + Zinc + Azotobactor + Phosphorous Solubilizing Bacteria).

It is concluded from the investigation that the treatment T<sub>6</sub> (75% N through Urea+ MOP + SSP + 25% through vermicompost+ Boron + Zinc + Azotobactor + Phosphorous Solubilizing Bacteria) was found suitable for application in tomato cultivation.

## References

- Anonymous. 2013. Indian Horticulture Database-2013, National Horticulture Board, Ministry of Agriculture, Government of India. 262 pp.
- Chanda, G.K. Goutam B. and Chakraborty, S.K. 2011. The effect of vermicompost and other fertilizers on cultivation of tomato plants. *J. Hort. and Forestry*. 2011. 3(2):42-45
- CHANDRAMOHAN, S., 2002, Organic farming on cotton + black gram intercropping systems. M.Sc. (Agri.) Thesis, Tamil Nadu Agri. Univ. Coimbatore, India.
- CHEETRI, M. AND THAPA, U. 2006, Effect of bio-fertilizer and plant growth promoting bacteria on growth attributes and tuber yield of potato. *Haryana J. Hort. Sci.*, 35 (1&2): 143-145.
- CHHONKAR, D. S., PUSPENDERA, K. AND SHORTI, S. K., 2011, Efficiency of bio-fertilizer in potato. *Ann. of Hort.*, 4 (2): 197-201.
- CHINNASWAMI, C. AND MARAKULANDAI, A., 2012, Influence of organic and inorganic manures on the firmness and storage life of tomato. *South Indian Hort.*, 45 (6): 115-120.
- Çhummyani, Kanaujia, S.P., Singh, VB, and Singh, A.K. 2012. Effect of integrated nutrient management on growth, yield and quality of tomato (*Lycopersicon esculentum* Mill), *J. Soils Crops*. 22(1):65-71.
- CHUMYANI, S. P., KANAUIA, S., SINGH, V. B. AND SINGH, A. K., 2010, Effect of Integrated Nutrient Management on growth, yield and quality of tomato. *J. Soil & Crop.*, 22 (1): 67-71.
- DEKA, B. C., BORA, G. C. AND SHADEQUE, A., 1996, Effect of *Azospirillum* on growth and yield of chilli cv. Pusa Jwala. *Haryana J. Hort. Sci.*, 25 (2): 44. 47.
- DEVI, H. J., MAIY, N. C., PARIJA AND THAPA, U., 2002, Response of brinjal to different sources of nitrogen. *Veg. Sci.*, 29 (1): 45-47.
- DIBYENDU, C., AJAYA. S. AND SINGH, R. K., 2010, Fertilizer recommendations based on targeted yield concept involving Integrated Nutrient Management for potato. *Indian J. Agri. Sci.*, 80 (12): 1048-1053.
- DILIP SINGH AND S. MUKHERJEE., 2010, Effect of farm yard manure, chemical and bio-fertilizer on growth parameters and yield of brinjal (*Solanum melongena* L.). *Green Frmg.*, 1 (2): 155-157.

- GHONAME, A. AND SHAFEEK, M. R., 2005, Growth and productivity of sweet pepper (*Capsicum annuum* L.) grown in plastic house as affected by organic, mineral and bio-fertilisers. *J. of Agron.*, 4 (4): 369-372.
- GOSAVI, P. U., KAMBALE, A. B. AND PANDURE, 2010, Effect of organic and bio-fertilizer on quality of tomato fruits. *The Asian J. Hort.*, 5 (2): 376-378.
- GOVINDARAJAN, K. AND THANGARAJU, M., 2001, *Azospirillum* is a potential inoculants for horticultural crops. *South Indian Hort.*, 49: 233-235. Effect of Integrated Nutrient Management on growth, yield and quality of tomato (*Solanum lycopersicum* L.) var. Arka Rakshak 51.
- JACKSON, M. L., 1973, *Soil and Chemical Analysis*, Prentice Hall of India Private Limited, New Delhi.
- JEEVANSAB, 2000, Effect of nutrient sources on growth, yield and quality of capsicum cv. California Wonder grown under different environments. M. Sc. (Agri.) Thesis, Univ. of Agric. Sci., Dharwad, Karnataka, India.
- KADALAG, A. D., JADHAV AND RAINA, R., 2007, Yield and quality of tomato fruits as influenced by bio-fertilizers. *Asian, J. of Sci.*, 2 (2): 95-99.
- KATTIMANI, S. K. AND SHASHIDHARA, G. B., 2006, Integrated Nutrient Management in chilli genotypes. *Karnataka J. Agric. Sci.*, 19 (3): 506-512.
- KATTIMANI, S., 2004, Response of chilli (*Capsicum annuum* L.) genotypes to Integrated Nutrient Management. M.Sc. (Agri.) Thesis, Univ. Agri. Sci., Dharwad, Karnataka, India.
- KC Prativa, BP Bhattarai (2011) Effect of integrated nutrient management on the growth, yield and soil nutrient status in tomato. *Nepal Journal of Science and Technology*,
- KHAN, Z., TIYAGI, SARTAJ, A., MAHMOOD, IRSHAD, RIZVI, AND ROSE, 2011, Effects of N fertilisation, organic matter, and bio-fertilizers on the growth and yield of chilli in relation to management of plant-parasitic nematodes. Aligarh Muslim Univ., Aligarh, India.
- KONDAPA, D., RADDAR, B. M., PATIL, P. L., HEBSUR, N. S. AND ALAGUNDAGI, S. C., 2009, Effect of Integrated Nutrient Management on growth, yield and economics of chilli cv. byadagi dabbi in vertosol. *Karnataka J. Agri. Sci.*, 22 (2): 438-440.
- KUMAR, S., RAKESH, K., SUTANU, M., DEVENDRA, K. AND MANOJ, K., 2014, Effect of organic manures and bio-fertilizers on growth, flowering, yield and quality of tomato cv. Pusa Sheetal. *Int. J. Agric. Sci.*, 10 (1): 329-332.
- MADALAGERI, B. B., DHARMATTI, P. R., HOSAMANI, R. M., MEHERWADE, M. N. AND BABALAD, H. B., 2006, Effect of nutrition on the physiological maturity of fruit and seed of tomato. *Prog. Hort.*, 21 (3-4): 168-172.

- MAHATO, P., ANOOP, B. AND CHAUHAN, J, S., 2009 Effect of Azotobacter and nitrogen on growth in tomato. Res., 1 (4): 502-508.
- MALAWADI, M. N., PALLED, Y. B., BASAVARAJ, B. AND HIREMATH, S. M., 2003, Effect of secondary and micronutrients on yield and quality of chilli (*Capsicum annuum* L.). M. Sc. (Agri.) Thesis, Univ. Agri. Sci., Dharwad. Karnataka. 52 Siddaling P. Nagoni, M.Sc. (Agri.) 2015
- MANJURAL, H. M, ILLIA, M. N. AND MOLLA, A. H., 2012, Impact of Trichoderma on growth and yield of tomato. The Agri. Sci. Krishi foundation., 10 (2): 109-119.
- MANOJ, K. 2014, Impact of front line demonstration of INM on yield in tomato. Hort flora Res. Spectrum., 3 (3): 295-297.
- MEDHE, N. K., GAWADAB. V., RANDIVES. N. AND TEMKAR, B. G., 2010, Effect of Integrated Nitrogen Management on yield and economics of green chilli (*Capsicum annuum*) production. Green Frmg., 1 (3): 260-262.
- MEDHE. N. K., PATAL P. R., GAIKAWAD R. D. AND MANE V. A., 2011, Effect of Integrated Nitrogen Management on nutrient uptake of green chilli (*Capsicum annuum*) production. Green Frmg., 2 (6): 680-681.
- MOHANKUMAR A. B. AND NARASEGOWDA, N. C., 2010, Effect of different organic manurer and inorganic fertilizers on growth and yield of brinjal. (*Solanum melongena* L.). The Asian J. of Hort., 5 (2): 444 - 449.
- MUDASIR, M. M., CHATTOO, S., FAHEEMA, A. P. AND PARRY, F. A., 2009, Influence of organic and inorganic nutrients on growth and yield attributes of tomato. The Asian J. Hort., 7 (2): 337-339.
- NAIDU, A. K., KUSHWAH, S. S. AND DWIVEDI, Y. C., 2002, Influence of organic manures, chemical and bio-fertilizers on growth, yield and economics of brinjal. South Indian Hort., 50 (4-6): 370-376.

**References may be incomplete**

UNDER PEER REVIEW