

Original Research Article

Effect of Plant Growth Regulators on Yield Contributing Traits and Quality Attributes of Bottle Gourd (*Lagenaria siceraria*) cv. MGH-4

ABSTRACT

Bottle gourd cv. MGH-4 with two growth regulators having concentration NAA (30, 40, 50, 60, 70 & 80 ppm) and GA₃(30,40, 50, 60 & 70 ppm). Water was used as a control. A randomised block design with a plot size of 2 m × 2 m included three replications of each treatment. On March 30, 2019, the seeds were sown with a 2 m x 1 m planting spacing. The application of plant growth regulators has a major influence on both the reproductive and vegetative traits of the crop. The fruit yield per hectare (58.45 t/h), fruit diameter (23.89 cm), weight (1.273 kg), TSS (17.89 °Brix), vitamin C content (10.77 mg/100gm), and seed pulp ratio were all determined to be best when 40 ppm GA₃ was applied at the 2, 4 leaf stage (1:9.16).

Key words: Bottle gourd, MGH-4, PGR, NAA, GA₃, Yield and Quality.

INTRODUCTION

The bottle gourd, also referred to as lauki, kadu, ghiya, or doodhi in India (*Lagenaria siceraria* L. 2n = 2x = 22), is a popular crop. It could have originated in tropical Africa. This vegetable provides an excellent source of carbohydrates, vitamin A, vitamin C, and minerals. **Hilli et al., (2010)**. The bottle gourd is a plant that belongs to the kingdom Plantae, division Magnoliophyta, class Magnoliopsida, order Cucurbitales, and family Cucurbitaceae. The name "lagena," which translates to "the bottle," is where the genus *Lagenaria* of bottle gourds got its start **Minocha (2015)**. In tropical countries, *Lagenaria siceraria* (Molina) Stands, a vegetable crop with a fleshy fruit and seeded pepo, is widely cultivated (**Shah et al., 2010**). **De Candolle (1882)**. claims that bottle gourd has been discovered in its natural state in South Africa and India Cutler and Whitaker (1961) hypothesised that it is probably native to tropical Africa based on the diversity of seeds and fruits. It's delicious, crisp, and soft fruits are enjoyed by both rich and poor people (**Ram et al., 2006**).

Cucurbit flowering is important because it influences fruiting and production. On the same plant, staminate and pistillate flowers can be found in various arrangements and separately (**Desai et al., 2011**). Bottle gourds are normally harvested in India in two different crops: the first from mid-October seeding to mid-March harvesting, and the second from early-March sowing to mid-July harvesting (harvest). It is a monoecious annual plant with an ascending or trailing vine. Long, forked tendrils extend from the stem's hairy texture. Flowers have stalks, and the female flowers' stalks are shorter than the male flowers. They are axillary, solitary, and unisexual. It produces a wide range of morphologies of hard-shelled fruits, including long, oblong, and spherical ones.

The host of physiological processes involved in crop development, such as blooming and fruiting, are controlled by plant growth regulators (fruit set and parthenocarpy). Additionally, they are employed in the germination, inhibition of growth, and post-harvest ripening of assimilates. PGRs are employed in very tiny amounts to control plant development. Additionally, they regulate the length of the internode, blooming, fruit setting, and fruit ripening. They also govern the development of the shoot and the root. PGRs, which are frequently employed in horticulture, have been utilised to alter the timings of blooming and fruiting in a variety of fruits and vegetables (**Mishra 2014**).

NAA is an essential regulator of plant growth that enhances the development of bottle gourd plants by encouraging cell division, cell elongation, and cell enlargement in the apical region of plants (**Kore et al., 2003**). NAA is used to chemically thin fruit, reduce fruit drops, induce flowering, increase fruit size, and hence increase production. NAA interacts with genes via producing the enzymes required for the synthesis of cell wall and cytoplasmic components. NAA initiates continuous flowering. After NAA spraying, cucumbers produced the largest fruits with the thickest flesh (**Das and Rabha. 1999**).

Gibberellic acid is a vital growth regulator that may be applied in a number of ways to change plant growth, yield, and features that affect production (**Rafeekher et al., 2002**). Gibberellic acid, often known as GA₃, is the most well-known of the four main types of gibberellins. It promotes seed germination, cambial activity, growth, cell elongation, nucleic acid synthesis, protein synthesis, fruit set, leaf expansion, and dormancy break (**Devlin and Witham, 1986**).

2. MATERIAL AND METHODS

The present investigation on "**Effect of plant growth regulators on growth, yield and fruit quality of Bottle Gourd (*Lagenaria siceraria*) cv MGH-4**" was conducted at Horticulture Research Field, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj during mid- March to 2- August during the years 2019.

2.1 Experimental Design and Treatments

For the experiment, a bottle gourd cultivar called MGH-4 was employed. Three replications were used in the Randomized Block Design experiment. The bottle guard variety MGH-4 was treated with gibberellic acid (30, 40, 50, 60, 70, and 80 ppm) and naphthalene acetic acid (30, 40, 50, 60, and 70 ppm) at 2, 4, and 6 days after planting (DAS). The important aspects of the plant growth regulators used in the experiment. Throughout the course of the experiment, a total of 12 treatments including various concentrations of NAA and GA₃ were used. The treatments were T1 (Control), T2 (30 ppm NAA), T3 (40 ppm NAA), T4 (50 ppm NAA), T5 (60 ppm NAA), T6 (70 ppm NAA), T7 (80 ppm NAA), T8 (30 ppm GA₃), T9 (40 ppm GA₃), T10 (50 ppm GA₃), T11 (60 ppm GA₃) and T12 (70 ppm GA₃). Due to the size of the plant population, it was difficult to observe every plant, thus the technique of random sampling was employed to document observations of the numerous attributes of the plants during the investigation. Three plants were randomly selected from a representative sampling of the entire population from each plot.

2.2 Parameters of Study

Yield Parameters

2.2.1 Fruit Length (cm): A random sample ten fruit was taken from each plot to record fruit characteristics viz., fruit length and fruit circumference, fruit length was measured from the peduncle end of fruit up to the blossom scar point with the help of measuring tape. Total length of ten fruit was divided ten to obtain average length of the fruit variety.

2.2.2 Fruit Diameter (cm): The same fruits used for measuring the length of fruit were taken for measuring the diameter. Diameter of fruit was measured with the help of Vernier callipers and the average fruit diameter was calculated.

2.2.3 Fruit Weight (kg): The total yield of plot was recorded in kg and divided by the number of fruits of that plot to obtain average weight of edible green fruit or a variety in a particular replication.

2.2.4 Fruit Yield Per Hectare (t/ha): fruit yield at physiological maturity was recorded from the harvested fruits of net plot in each replication and the average of three replications was expressed as t/ha.

Quality parameters

2.2.5 T.S.S. (°Brix): T.S.S. (Total Soluble Solids) content of a solution is determined by the index of refraction. This is measured using a hand refractometer, and is referred to as the degrees Brix. Brix is the term used when a refractometer equipped with a scale, based on the relationship between refractive indices at 20°C and percentage by mass of total soluble solids of a pure aqueous sucrose. This tests the solids concentration of a sucrose containing solution.

2.2.6 Ascorbic Acid (Vitamin C) (mg/100gm): Ascorbic acid content of fruit was determined by diluting the known volume of juice with 3 percent meta-phosphoric acid and titrating with 2,6-dichlorophenol-indophenol solution (A.O.A.C., 1980), till the faint pink colour was obtained.

2.2.7 Seed-Pulp Ratio: Seed pulp ratio at physiological maturity was recorded by weight of fruit pulp (flesh) to weight of seed.

3. RESULTS AND DISCUSSION

The findings of the current study on the impact of plant growth regulators on bottle gourd growth characteristics. The findings have been explained in terms of the effects of several experimental treatments. The results of the experiment have been presented separately under the following headings.

Fruit Length (cm): Fruit length data shows the significant result. The maximum length of fruit recorded in T7NAA 80 ppm was 48.00 cm followed by T10 GA3 50 ppm (45.50 cm) which was found to be at par with each other and both treatments were superior to the T1 control (32.33).

Mandal and Maity (1990), observed that bottle gourd cv. Pusa Summer Prolific Long treated with the growth regulators NAA (100 or 200 ppm), GA3 (25 or 50 ppm), Ethrel (ethephon) (10 or 20 ppm), MH (100 or 200 ppm) or morphactin (25 or 50 ppm). Highest fruit length (61.6 cm) was obtained with 100 ppm NAA.

Fruit Diameter (cm): Fruit diameter data shows the significant result. The maximum diameter of fruit recorded in T9 GA3 40 ppm (23.89 cm) followed by T7 NAA 80 ppm (22.94 cm) which was found to be at par with each other. Minimum was found in treatment T1 control (13.44 cm).

Singh and Choudhary (1989) and Chovatia et al., (2010). Mandal and Maity (1990) observed that bottle gourd cv. Pusa Summer Prolific Long treated with the growth regulators NAA (100 or 200 ppm.), GA3 (25 or 50 ppm) Ethrel (ethephon) (10 or 20 ppm.), MH (100 or 200 ppm) or morphactin (25 or 50 ppm.). Highest fruit girth (22.5 cm) was obtained with 50 ppm GA3.

Fruit Weight (kg): Fruit weight data shows the significant results. The maximum weight of fruit recorded in T9 GA3 40 ppm (1.27 kg) followed by T7 NAA 80 ppm (0.96 kg) which was found to be at par with each other and both treatments were observed superior to the T1 control (0.69 kg).

Dixit and Mishra (2006) reported that application of growth regulators like GA3, (25 or 50 ppm), ethrel (ethephon) (250 or 500 ppm) and NAA (50 or 100 ppm) in bitter gourd cv. Coimbatore long (green). Among the growth regulators, GA3 at 50 ppm was found to be the most effective for higher fruit weight (1.74 kg).

Fruit Yield Per Hectare (t/ha): Total fruit yield per hectare varies from 16.85 t/ha to 58.45 t/ha. Analysis of total yields per hectare data shows the significant results. Maximum fruit yield recorded in T9 GA3 40 ppm (58.45 t/ha) followed by T7 NAA 80 ppm (42.49 t/ha) which was found to be at par with each other. Minimum fruit yield per hectare was recorded in treatment T1 control (16.85 t/ha).

Merwade et al., (2006) reported that Application of growth regulators like GA3 (25 or 50 ppm), ethrel (Ethephon) (250 or 500 ppm) and NAA (50 or 100 ppm) in bitter gourd cv. Coimbatore long (green). Among the growth regulators, NAA at 50 ppm was found to be most effective for fruit yield (18.16 t/ha). **Sharma et al., (1998)** found in his studies with the cultivar Pusa summer Prolific Long the plants were treated with several growth regulators, some were applied at the 2- and 4-true leaf stages and some 35 and 45 days after sowing. Data are tabulated on growth, flowering and yields. The highest average yields were obtained with GA3 at 50 ppm (316.8 q/ha) and with Atonik [Nitrophenolate-sodium+nitroguaincol-sodium] at 0.1% (316.2 q/ha). The control yield was 116.7 q/h.

T.S.S. (°Brix): total soluble solid data shows the significant result. The maximum T.S.S was recorded in T9 GA3 40 ppm (17.89 °Brix) followed by T7 NAA 80 ppm (16.17 °Brix) was found to be at par with each other. The minimum was recorded in treatment T1 (5.36 °Brix).

Ascorbic Acid (Vitamin C) (mg/100gm): Ascorbic acid varies from 10.77 mg/100gm to 6.83 mg/100gm. Analysis of ascorbic acid data shows the significant result. Maximum ascorbic acid recorded in T9 GA3 40 ppm (10.77 mg/100gm) followed by T7 NAA 80 ppm (9.60 mg/100gm) which was found to be at par with each other and minimum in T1 control (6.83 mg/100gm of fruit pulp).

Seed-Pulp Ratio: Seed-Pulp Ratio data shows the significant result. The maximum seed-pulp ratio was recorded in T9 GA3 40 ppm (1:9.16) followed by T7 NAA 80 ppm (1:8.68) which was found to be at par with each other as compare to control T1 (1:5.05).

TABLE 1. Effect of Plant Growth Regulators on Growth Parameters of Bottle Gourd (*Lagenaria siceraria*)cv.MGH-4

Sr. No.	Treatments	Fruit Length (cm)	Fruit Diameter (cm)	Fruit Weight (kg)	Fruit Yield Per Hectare (t/ha)	T.S.S. (°Brix)	Ascorbic Acid (Vitamin C) (mg/100gm)	Seed-Pulp Ratio
1.	T1 (Control)	32.333	13.443	0.687	16.847	5.363	6.830	5.047
2.	T2 (30 ppm NAA)	34.833	18.027	0.763	27.627	9.670	7.607	5.703
3.	T3 (40 ppm NAA)	39.667	19.250	0.867	34.687	11.633	8.143	6.680
4.	T4 (50 ppm NAA)	39.167	20.027	0.670	22.140	12.613	7.803	7.833
5.	T5 (60 ppm NAA)	42.250	18.973	0.803	30.867	13.557	8.553	8.280
6.	T6 (70 ppm NAA)	43.917	17.250	0.817	20.427	15.640	9.220	8.007
7.	T7 (80 ppm NAA)	48.000	22.943	0.957	42.487	16.170	9.603	8.677
8.	T8 (30 ppm GA ₃)	41.333	19.223	0.887	21.733	15.353	8.593	6.583
9.	T9 (40 ppm GA ₃)	39.250	23.890	1.273	58.447	17.887	10.773	9.157
10.	T10 (50 ppm GA ₃)	45.500	20.807	0.943	23.600	14.537	8.657	7.003
11.	T11 (60 ppm GA ₃)	41.167	18.973	0.863	29.220	13.343	7.900	7.037
12.	T12 (70 ppm GA ₃)	41.417	18.277	0.847	26.430	13.997	7.780	6.183
	SEm (±)	0.89	0.73	0.03	3.15	0.18	0.24	0.19
	C.D. at 5% of Level	1.87	1.53	0.07	6.56	0.37	0.51	0.41

CONCLUSION

According to the findings of the current study, "Effect of Plant Growth Regulators on Growth, Yield, and Fruit Quality of Bottle Gourd (*Lagenaria siceraria*) cv MGH-4," the treatment T₉, which involved applying 40 ppm GA₃ at the 2 and 4 leaf stages, was found to be superior in terms of bottle gourd fruit yield and quality parameters.

REFERENCES

- Chaudhary, B. R., Sharma, M. D., Shakya, S. M., and Gautam, D. M. (2006). Effect of plant growth regulators on growth, yield and quality of Chilli (*Capsicum annum* L.) at Rampur, Chitwan. *Journal of the Institute of Agriculture and Animal Science*, **27**, 65-68.
- Chovatia, R.S., Ahlawat, T.R., Kavathia, Y.A., Jivani, L.L. and Kaila, D.C., (2010). Effect of plantgrowth regulators on vegetative growth, flowering and yield of bitter gourd cv. Priya. *Indian Journal of Horticulture*, **67**:254-258.
- Devlin, R. M. and Witham, F. H. (1986). Plant physiology. CBS Publicaion and Distributors., New Sex expression and fruit yiel or bottle gourd (*Lagenaria siceraria*) (01) StandlJ Cv. Fusamusk melon (*Cucumis melo* L.). *Plant archives*, **15**(2): 899-Delhi. pp. 410-447. 91
- Das, A, and Rabha, B, K, (1999). Effect of growth regulators on Size and quality of cucumber inplastic green house during rabi season, *Crop Res.*, **18**(3): 390-396

- Hilli JS, Vyakarnahal BS, Biradar DP, Hunje R. (2010)** Effect of growth regulators and stages of spray on growth, fruit set and seed yield of ridge gourd *Luffa acutangula* L. Karnataka J Agri. Sci.; **23(2)**:239-242
- Kore, V.N., Khade, H.P., Nawale, R.N., Patil, R.S. And Mane, A.V., (2003)**, Effect of growth regulators on growth, flowering and yield of bottle gourd variety Samrat under Konkan conditions. *Journal of Soils and Crops*, **13(1)**:18- 21.
- Mandal, D., Paria, N.C. And Maity, T.K., (1990)**, Response of bottlegourd (*Lagenaria siceraria* (Molinal Standl.) to some plant growth regulators. *Crop Research*, **3(2)**: 244-246.
- Mishra, N. (2014)**. Growth and yield response of pea (*Pisum sativum* L) to Integrated Nutrient Management- A Review. *Journal of Plant and Pest Science*, **1(2)**: 87-95.
- Minocha, S., (2015)**. An overview on *Lagenaria siceraria* (bottle gourd). *Journal of Bio-medical and Pharmaceutical Research*, **4(3)**:4-10
- Rafeekher M, SA Nair, PN Sorte, GP Hatwal and PM Chandhan, (2002)**. Effect of growth regulators on growth and yield of summer cucumber. *Journal of Soils and Crops*, **12**: 108-110.
- Ram D, Rai M, Rai N, Yadav DS, Pandey S, Verma A, Lal H, Singh N, Singh S. (2006)**. Characterization and evaluation of winter fruited bottle gourd, *Lagenaria siceraria* (Mol.) USDA, 2016. National Nutrient Database for Standard Reference, Release 28 slightly revised May, 2016. Basic Report 11218, Gourd. white-flowered (calabash), raw. Report date January 29, 2018, 08:53 EST. Retrieved from <http://ndbna1.usda.gov/ndWfoods/show/2970>
- Shah, B.N., A.K. Seth, and R.V. Desai. (2010a)**. Phytopharmacological profile of *Lagenaria siceraria*: a review. *Asian J. Plant Sci.* **9(3)**:152–157