

Original Research Article

Effect of STCR Based Long-Term Fertilization on Yield Attributing Parameters and Yield of Wheat (*Triticumaestivum L.*)

Comment [h1]: Write in full STCR

ABSTRACT

A field experiment was carried out during *Rabi* season of 2020–21 and 2021–22 at the Jawaharlal Nehru KrishiVishwaVidyalaya (JNKVV), Jabalpur (Madhya Pradesh) research field to find out the effect of STCR based long-term fertilization on yield attributing parameters and yield of wheat. The treatments details were T₁; Control, T₂; 120 N: 80 P₂O₅: 60 K₂O kg ha⁻¹, T₃; Target yield 4.5 t ha⁻¹, T₄; Target yield 6.0 t ha⁻¹, T₅; Target yield 4.5 t+FYM 5 t ha⁻¹; T₆; Target yield 6.0 t+FYM 5 t ha⁻¹ and carried out in Randomized Block Design (RBD) with four replications. Research finding of long term field experiment revealed that application of soil test crop response recommendation for target yield 6 t ha⁻¹+FYM 5 t ha⁻¹ gave maximum earhead length (14.59 and 14.10 cm), number of grains ear⁻¹ (66.46 and 63.30), test weight (41.93 and 41.35 g), grain yield (5568 and 5052 kg ha⁻¹) and straw yield (7315 and 5969 kg ha⁻¹) of wheat. The balanced application of fertilizers by the soil and the crop's needs for potential growth and development may account for the increased yield under the STCR technique with and without FYM. The inclusion of FYM resulted in a potential movement of water, air, temperature, and nutrients in the soil, which may have improved the conditioning of the rhizospheric environment and further raised yield when integrated with STCR. Due to better physical, chemical, and biological characteristics, enhanced microbial activity and the conversion of inaccessible nutrients into available forms, this STCR technique also aids in higher production.

Comment [h2]: Write in full

Comment [h3]: Same as above

Keywords: *FYM, STCR, target yield, wheat, yield attributing*

1. INTRODUCTION

Around the world, a vast population highly depends on wheat (*Triticumaestivum L.*), which provides approximately 20% of people's daily protein and calories [1]. Wheat is the second most important food grain of India with an area of 32 mha, production of 107.86 mt, and an average productivity of 3370 kg ha⁻¹[2]. Due to varying meteorological conditions,

Comment [h4]: Globally,

genotypes, sowing time and techniques, and other management practices, wheat yield varies significantly across India's many agro-ecologies [3]. The wheat crop contributed significantly in food grain production for ensuring nations food security. There are reports of decreasing crop yields which raised questions on sustainability of the wheat crop. The reasons behind is farmers use excessive amounts of chemical fertilizer to increase yield attributes and yields [4] but choosing how much fertilizer to use on which crops depends on understanding how those crops will respond to nutrient application and predicted output. This approach costs farmers money in addition to harming the soil's health [5] and [6]. Additionally, the negative effects of high-analysis chemical fertilizer use on soil productivity and the environment created an energy problem and a concern with sustainability in agriculture. Environmental contamination is also brought on by the indiscriminate and ongoing use of inorganic fertilizers, Prajapat et al. [7] and Prasad [8] which have been found to have a deleterious impact on the physico-chemical and biological properties of soil [9]. Wheat production can be increased significantly using the soil test crop response (STCR) technique [10] and [11]. In this method, the recommended fertilizer doses are determined using fertilizer adjustment equations that were created after a substantial association was established between the soil test results and the additional fertilizer. Because it incorporates the combined use of soil and plant analysis, which provides details on the actual balance between applied nutrients and the nutrients that are now accessible in the soil, recommendations based on the STCR correlation idea are more quantitative, exact, and relevant [12][13]. Alternatively, organic manures (OM) are a rich source of nutrients; their single application is insufficient to cover the nutrient needs of high-producing types and frequently provides subpar crop production [11] and [14]. Due to exhaustion in soil health, employing the generally recommended dose (GRD) of fertilizer is also unable to maintain yields about the economic returns of crops, necessitating refinement for balanced crop nutrition [15]. Therefore, using only organic matter or chemical fertilizer won't help an intensive production system become more sustainable [16]. To provide balanced fertilization to crops, it is a step forward to employ an appropriate blend of organic and chemical fertilizers [17], depending on the soil fertility state [18]. This integrated nutrient management (INM) type can increase yield attributing and yield of wheat [19], [20] and [21]. According to soil test crop response based research experiment results shown very close correlation between targeted yield and yield actually obtained and an evidence of the usefulness of soil testing within limit of variation under field conditions. Keeping all this point in mind the present investigation was conducted to determine the "Effect of STCR based long-term fertilization on yield attributing parameters and yield of wheat".

2. MATERIAL AND METHODS

1.1. Experimental Site and Experimentation

Field experiments were conducted during *Rabi* season (December–April) of 2020–21 and 2021–22 at the Jawaharlal Nehru KrishiVishwaVidyalaya (JNKVV) research field under the ongoing AICRP on Soil Test Crop Response, managed by Department of Soil Science, Jabalpur at 23°13' North latitude, 79°57' East longitudes and at an elevation of 393 m above mean sea level. The treatments details were T₁; Control, T₂; 120 N:80 P₂O₅:60 K₂O kg ha⁻¹, T₃; Target yield 4.5 t ha⁻¹, T₄; Target yield 6.0 t ha⁻¹, T₅; Target yield 4.5 t+FYM 5 t ha⁻¹; T₆; Target yield 6.0 t+FYM 5 t ha⁻¹ and carried out in Randomized Block Design (RBD) with four replications. Nutrients doses (kg ha⁻¹) were calculated from the fertilizer adjustment equations for targeting yield 4.5 and 6.0 t ha⁻¹ of wheat. The fertilizer adjustment equations are given below:

$$FN=4.40 T-0.40 SN \dots\dots\dots(1)$$

$$FP_2O_5=4.00 T-5.73 SP \dots\dots\dots(2)$$

$$FK_2O=2.53 T-0.16 SK \dots\dots\dots(3)$$

Whereas,

FN=Fertilizer nitrogen (kg ha⁻¹); FP₂O₅=Fertilizer phosphorus (kg ha⁻¹); FK₂O=Fertilizer potassium (kg ha⁻¹); T=Desired yield target (t ha⁻¹); SN=Available soil nitrogen (kg ha⁻¹); SP=Available soil phosphorus (kg ha⁻¹); SK=Available soil potassium (kg ha⁻¹).

1.2. Data Collection

Wheat growth parameters such as length of earhead¹, number of grains per ear head were studied at harvest. Five randomly chosen ear heads from tagged plants were used to measure the length of earhead¹ after that threshed manually and number of grains per ear head was counted. For estimating grain and straw yield, wheat crop from a net plot area of 16 m² was harvested and sun dried. After drying, manual threshing was carried out. Grain weight and straw weight was taken from each treatment and expressed as kg ha⁻¹. 1000 grains were counted manually after threshing of each plot then weighed in electronic balance and expressed in gram.

3. RESULTS AND DISCUSSION

Comment [h5]: Rewrite as above

1.1. Effect of different treatments on yield attributes of wheat

The data pertaining to the yield attributes and yield of wheat is presented in Table 1. The significantly higher earhead length of 14.59 and 14.10 cm was observed in Target Yield 6 t ha⁻¹+FYM 5 t ha⁻¹ which were at par with Target Yield 6.0 t ha⁻¹ (12.04 and 11.56 cm) but found statistically significant over rest of the treatments during 2020–21 and 2021–22. The minimum earhead length was recorded under control 6.12 and 5.70 cm during both the year (Figure 1). Similarly, long term nutrient management of STCR on number of grains revealed that the application of nutrient as GRD, target yield of 4.5 t ha⁻¹, target yield of 6.0 t ha⁻¹ target yield of 4.5 t ha⁻¹+FYM 5 t ha⁻¹ and target yield of 6.0 t ha⁻¹+FYM 5 t ha⁻¹ significantly increased the no. of grains ear⁻¹ over control during both the years. However, application of

Table 1: Effect of long-term nutrient management of STCR approach on yield attributes in wheat

nutrient for target yield of 6.0 t ha⁻¹, target yield of 4.5 t ha⁻¹+FYM 5 t ha⁻¹ and target yield of 6.0 t ha⁻¹+FYM 5 t ha⁻¹ were found significantly superior to GRD and target Yield of 4.5 t ha⁻¹ but the treatment were found at par amongst themselves during both the years.

The maximum number of grain earhead⁻¹ during first year (66.46) and second year (63.30) were observed with target yield of 6 t ha⁻¹+FYM 5 t ha⁻¹. This might be due to supply of the nutrients in balanced amount and slowly release of nutrients through integration use of FYM which helped to produce more numbers of grains ear⁻¹ and earhead length. The spike length was directly showed the higher no of grains in panicle so that yield will be increased. These results are supported by Moharana *et al.* [22] Sheoran *et al.* [23] and Kumar *et al.* [24]. An examination of data indicated that test weight in (Table 1) target yield of 6.0 t ha⁻¹+FYM 5 t ha⁻¹ found maximum which was significantly higher than GRD (120 N: 80 P₂O₅: 60 K₂O kg ha⁻¹) and control but found to be at par with remaining treatment. According to Patel *et al.* [25], an adequate nutrient supply increases photosynthetic activities. It translocate more photosynthates in the reproductive stages of the crop, thereby promoting growth and increasing wheat's test weight—this increased wheat's test weight when chemical fertilizer, including organic manure, was used. The increased availability of nutrients to plants, first through fertilizers and later through FYM, is attributed to the rise in yield. In addition to supply nutrients, FYM also improves physical conditions of soil. Tiwari *et al.* [26], Sellamuthu *et al.* [27], Singh *et al.* [28] and Rai *et al.* [29] were also noted similar results which support the results of present experiment.

Treatment	Number of grains ear ⁻¹		Test weight (g)	
	2020-21	2021-22	2020-21	2021-22
T ₁ -Control	38.87 ^c	34.73 ^c	35.12 ^c	34.54 ^d
T ₂ - GRD (120 N: 80 P ₂ O ₅ : 60 K ₂ O kg ha ⁻¹)	47.60 ^b	45.95 ^b	38.45 ^b	37.33 ^{cd}
T ₃ - Target Yield 4.5 t ha ⁻¹	50.96 ^b	47.80 ^b	39.82 ^{ab}	38.20 ^{bc}
T ₄ - Target Yield 6.0 t ha ⁻¹	62.25 ^a	60.20 ^a	41.38 ^{ab}	41.08 ^{ab}
T ₅ -Target Yield 4.5 t ha ⁻¹ +FYM 5 t ha ⁻¹	61.61 ^a	58.82 ^a	40.83 ^{ab}	39.26 ^{abc}
T ₆ -Target Yield 6.0 t ha ⁻¹ +FYM 5 t ha ⁻¹	66.46 ^a	63.30 ^a	41.93 ^a	41.35 ^a
SEm±	2.06	1.78	1.08	1.00
CD (p=0.05)	6.22	5.36	3.25	3.03

Different letters, i.e., a, b, c within a column indicate significant difference accordingly to least significant difference (LSD) test (CD p=0.05)

Comment [h6]: Rewrite this




Figure 1: Effect of long term nutrient management of STCR approach on earhead length

3.2. Effect of different treatments on grain and straw yield of wheat

The effect of different treatments on grain and straw yield of wheat has been shown in Table 2. It is evident from the data indicated that the application of STCR based NPK level as GRD, target yield of 4.5 t ha⁻¹, target yield of 6.0 t ha⁻¹, target yield of 4.5 t ha⁻¹+FYM 5 t ha⁻¹ and target yield of 6.0 t ha⁻¹+FYM 5 t ha⁻¹ significantly increased the grain and straw yield of wheat over control during both the years. However, application of nutrient for target yield of 6 t ha⁻¹+FYM 5 t ha⁻¹, target yield of 6.0 t ha⁻¹ and target yield of 4.5 t ha⁻¹+FYM 5 t ha⁻¹ significantly increased the grain and straw yield of wheat over GRD. The application of nutrient for target yield of 6.0 t ha⁻¹+FYM 5 t ha⁻¹ also found significantly superior to target yield of 4.5 t ha⁻¹ for grain and straw yield during both the years except straw yield during second year. The maximum grain yield 5568 and 5052 kg ha⁻¹ and straw yield 7315 and 5969 kg ha⁻¹ were observed with nutrient application for target yield of 6.0 t ha⁻¹+FYM 5 t ha⁻¹

Table 2: Effect of long term nutrient management of STCR approach on yield and straw in wheat

during first year, second year. The balanced application of fertilizers by the soil and the crop's needs for potential growth and development may account for the increased yield under the STCR technique with and without FYM. The inclusion of FYM resulted in a potential movement of water, air, temperature, and nutrients in the soil, which may have improved the conditioning of the rhizospheric environment and further raised yield when integrated with STCR. Due to better physical, chemical, and biological characteristics, enhanced microbial activity and the conversion of inaccessible nutrients into available forms, this STCR technique also aids in higher production [30]. Wheat's increased yield appeared to be the result of a combination of yield characteristics supported by a balanced supply of nutrients [31],[32] and [33]. The results of present experiment confirmed the finding of Parewa et al. [14] and Moharana et al. [22].

Treatment	Grain yield (kg ha ⁻¹)		Straw yield (kg ha ⁻¹)	
	2020–21	2021–22	2020–21	2021–22
T1- Control	2145 ^d	961 ^d	3017 ^d	1445 ^c
T2- GRD (120 N: 80 P ₂ O ₅ : 60 K ₂ O kg ha ⁻¹)	4171 ^c	3898 ^c	5495 ^c	4734 ^b
T3- Target Yield 4.5 t ha ⁻¹	4683 ^{bc}	4219 ^{bc}	6079 ^{bc}	5281 ^{ab}
T4- Target Yield 6.0 t ha ⁻¹	5177 ^{ab}	4805 ^{ab}	6731 ^{ab}	5906 ^a
T5-Target Yield 4.5 t ha ⁻¹ + FYM 5 t ha ⁻¹	5095 ^{ab}	4586 ^{ab}	6688 ^{ab}	5883 ^a
T6- Target Yield 6.0 t ha ⁻¹ +FYM 5 t ha ⁻¹	5568 ^a	5052 ^a	7315 ^a	5969 ^a
SEm±	184.23	198.08	272.10	245.11
CD (p=0.05)	555.22	596.96	820.02	738.68

Different letters, i.e., a, b, c within a column indicate significant difference accordingly to least significant difference (LSD) test (CD $p=0.05$).

Comment [h7]: Pls reconcile your tables

4. Conclusion

Results of a long-term experiment concluded that STCR recommendation on wheat brought an additive effect in increasing yield attributes and yield of wheat crop. Application of STCR recommended for target yield 6.0 t ha⁻¹+FYM 5 t ha⁻¹ gave maximum yield attributes, grain and straw yields. These study results proved that STCR recommended gave better results of all yield attributing characteristics and yield of wheat in the comparison of general recommended dose.

REFERENCES

1. Shiferaw B, Smale M, Braun, HJ, Duveiller E, Reynolds M, Muricho G. Crops that feed the world past successes and future challenges to the role played by wheat in global food security. Food Security. 2013;5(3):291–317.

2. Agricultural. Statistics at a Glance. Government of India, Ministry of Agriculture and Farmers Welfare, Department of Agriculture Cooperation and Farmers Welfare, Directorate of Economics and Statistics, 2021;51.
3. Acuna T, Wade L. Genotypexenvironment interactions for root depth of wheat. *Field Crops Research* 2012;37(4):117–125.
4. John PS, George M, Jacob R, Nutrient mining in agro-climatic zones of Kerala. *Fertilizer News*.2001;46(8),45–57.
5. Saxena AK, Singh, S, Srivastava A, Gautam, P, Yield target approach for assessing the fertilizer requirement of onion in mollisols of Uttarakhand. *Indian Journal of Horticulture* 2008;65(3):302–306.
6. Chatterjee D, Srivastava A, Singh RK, Fertilizer recommendations based on targeted yield concept involving integrated nutrient management in potato in Tarai belt of Uttarakhand. *The Indian Journal of Agricultural Sciences*.2010;80(12): 1048–1063.
7. Prajapat K, Vyas AK, Dhar S, Root growth and productivity of wheat, chickpea and potato as influenced by nutrient management practices. *International Journal of Bio-resource and Stress Management*.2016;7(1):052–059.
8. Prasad, R. 2005. Rice–wheat cropping systems. *Advances in Agronomy* 86, 255–339.
9. Virmani SM. The twenty first-Dr. R.V. Tamhane memorial Lecture: UNCEED Agenda 21: The new challenge for soil research. *Journal of Indian Society of Soil Science* 1994;42:516–523.
10. Kumari N, Kharia SK, Mandeewal RL, Rojh MR, Yadav A, Kumari SC. Effect of soil test crop response approach on yield, nutrient content and uptake by wheat (*Triticumaestivum* L.). *International Journal of Plant & Soil Science*. 2022;34(14): 1–6.
11. Sharma VK, Pandey RN, Kumar S, Chobhe AK, Chandra S. Soil test crop response-based fertilizer recommendations under integrated nutrient management for higher productivity of pearl millet (*Pennisetum glaucum*) and wheat (*Triticumaestivum*) under long term experiment. *Indian Journal of Agricultural Sciences*. 2016;86(8):76–81.
12. Gulati R, Lawrence, PR, Puranam, P. Adaptation in vertical relationships: Beyond incentive conflict. *Strategic Management Journal*. 2005; 26(5):415–440.
13. Pogula S, Truptimayee, Mishra, A, Saren, S, Dey, P, Soil test based fertilizer recommendation for targeted yield of French bean (*Phaseolus vulgaris*) under

- rice–French bean cropping system. *International Journal of Bio-resource and Stress Management*.2016; 7(5):1128–1130.
14. Parewa HP, Moola R, Jain, LK, Choudhary A, Ratnoo, SD. Impact of organic nutrient management practices on yield attributes, yield and economics of wheat (*Triticumaestivum* L.). *International Journal of Bio-resource and Stress Management*.2019;10(3):257–260.
 15. Singh DK, Pandey PC, Nanda G, Gupta S. Long-term effects of inorganic fertilizer and farmyard manure application on productivity, sustainability and profitability of rice–wheat system in mollisols. *Archives of Agronomy and Soil Science*.2019;65(2), 139–151.
 16. Patel A, Neupane, MP, Nanda G, Singh, SP. 2016. Effect of NPK management and bioinoculants on growth and yield of wetland rice (*Oryza sativa* L.). *Environment & Ecology* 33(4C), 181–185.
 17. Mahajan, A., Gupta, R.D., Role of INM in sustainable rice–wheat cropping system. In: *Integrated Nutrient Management (INM) in a Sustainable Rice–Wheat Cropping System*. Springer, Dordrecht, The Netherlands, 2009;139–168.
 18. Santhi R, Bhaskaran A, Natesan R, Integrated fertilizer prescriptions for beetroot through inductive cum targeted yield model on an alfisol. *Communications in Soil Science and Plant Analysis*.2011;42(16):1905–1912.
 19. Nanda G, Sravan US, Singh A, Singh SP. Effect of NPK levels and bio-organics on growth, yield and economics of basmati rice (*Oryza sativa* L.) cv HUBR 10-9. *Environment & Ecology* 2016; 34(3C):1530–1534.
 20. Antil RS, Narwal, RP, Singh B, Singh JP. Long-term effects of FYM and N application on soil health and productivity under pearl millet–wheat cropping system. *Indian Journal of Fertilizer*.2011; 7:14–32.
 21. Chesti MH, Kohli A, Sharma, AK. Effect of integrated nutrient management on yield and nutrient uptake by wheat (*Triticumaestivum*) and soil properties under intermediate Zone of Jammu and Kashmir. *Journal of Indian Society of Soil Science*.2013; 61(1), 79–85.
 22. Moharana PC, Sharma BM, Biswas DR. Changes in the soil properties and availability of micronutrients after six-year application of organic and chemical fertilizers using STCR-based targeted yield equations under pearl millet–wheat cropping system. *Journal of Plant Nutrition*.2017;40(2),165–176.
 23. Sheoran, S., Raj, D., Antil, R.S., Mor, V.S., Dahiya, D.S. Productivity, Seed Quality and Nutrient Use Efficiency of Wheat (*Triticumaestivum*L.) under Organic, Inorganic

- and Integrated Nutrient Management Practices after Twenty Years of Fertilization. *Cereal Research Communications*.2017;45(2):315-325.
24. Kumar V, goyal V and Dey P. Impact of STCR based long term integrated management practices on soil chemical properties and yield attributing parameters of wheat and pearl millet in semi-arid North- west India. *Int. Journal of Chemical studies*. 2020;8(4):1320-1328.
 25. Patel GG, Sadhu AC, Patel HK, Shan SN, Lakum YC.Effect of organic and inorganic fertilizers in comparison with humic acid on growth and yield of wheat (*Triticumaestivum* L.). *International Journal of Agriculture Sciences*.2018;10, 6524–6527.
 26. Tiwari LB, Mishra DD, Gupta Evaluation of integrated nutrient management options in rice (*Oryza sativa*)–wheat (*Triticumaestivum*) cropping system in reclaimed sodic land. *Indian Journal of Agronomy* 2013; 58(2):137–145.
 27. Sellamuthu KM, Santhi R, Maragatham S, Dey P. Validation of soil test and yield target-based fertilizer prescription model for wheat on inceptisol. *Research on Crops* 2015;16(1):53–58.
 28. Singh YV, Singh SK, Sharma PK, Singh P. Soil Test based Integrated Fertilizer Recommendation for Wheat (*Triticumaestivum*) in an inceptisol of Eastern Plain zone of Uttar Pradesh. *Journal of the Indian Society of Soil Science* 2014; 62(3), 255–258.
 29. Rai HK, Sachidanand B, Baghel SS, Dey P. Evaluation of productivity and economics of wheat under STCR based nutrient application with and without FYM in a Vertisol. *Eco. Env. and Cons* 2016; 22: S107-S111.
 30. Sharma V.K, Pandey RN, Sharma BM. Studies on long term impact of STCR based integrated fertilizer use on pearl millet (*Pennisetum glaucum*)–wheat (*Triticumaestivum*) cropping system in semi-arid condition of India. *Journal of Environmental Biology* 2015; 36(1), 241–247.
 31. Yaduvansi NP, Sharma S, Swaroop A. Impact of integrated nutrient management on soil properties and yield of rice and wheat in a long-term experiment on a reclaimed sodic soil. *Journal of the Indian Society of Soil Science* 2013;61:188–194.
 32. Meena KB, Alam MS, Singh H, Bhat MA, Singh AK, Mishra AK, Thomas T. Influence of farmyard manure and fertilizers on soil properties and yield and nutrient uptake of wheat. *Int. J. Chem. Stud.*2018, 6(3):386-390.
 33. Bangre J, Diwedi AK, Mohanty M, SubhashDwivedi BS, DwivediSK. Effect of long-term fertilizer application on performance of wheat crop and soil properties in a Vertisol. *Ind. J. Pure App. Biosci.*2020, 8(1), 217-227.