

# **Original Research Article**

## **Estimation of Heritability and Genetic Advance in wheat (*Triticum aestivum*.L.)**

### **ABSTRACT**

An experiment was conducted to get information on genetic variability, heritability and genetic advance in the 100 genotypes including ten parents and 45 F<sub>1</sub> and 45 F<sub>2</sub> for obtained through half diallel mating design in pea during 2021-22 Oil Seed Farm, Kalyanpur, C.S. Azad University of Agriculture and Technology, Kanpur-208002 (U.P.) Quantitative analysis were carried out for all the characters which are directly or indirectly associated with the yield and yield contributing traits. Analysis of variance showed the significant variability for all the studied characters for parents and in F<sub>1</sub> significant variability observed in all the traits except spike length, number of grains per spike and protein content reflecting considerable amount of heterotic response in these attributes. High heritability was observed for all the characters in F<sub>1</sub> and F<sub>2</sub> both the generations. Highest value of GCV and PCV were observed for grain yield per plant (20.01) in F<sub>2</sub> which indicates the presence of high genetic variation. High heritability coupled with high genetic advance for protein content and yield per plant which indicate the presence of additive gene action and used for future population improvement. The genotypes with specific characters can be utilized for hybridization programme.

*Key words:* Heritability, Genetic advance, GCV and PCV

### **INTRODUCTION**

Wheat (*Triticum aestivum* L., 2n=42) belongs to the family Poaceae (Gramineae) and tribe Triticeae containing more than 15 genera and 300 species including wheat and barley. *T. aestivum* is a segmental allohexaploid (2n = 6x = 42, AABBDD) originated in the Fertile Crescent area of South-Western Asia (Lupton 1987), its geographical centre of origin and spread globally for cultivation and consumption. Allohexaploid wheat possesses three genomes and A, B, and D are three genomes. The genome "A" comes from wild einkorn wheat (*Triticum monococcum* var. *urartu*), "B" comes from an unknown species, and genome "D" comes from a weedy grass *Squarrosa Aegilops*.

Hexaploid wheat (*Triticum aestivum* L.,  $2n = 42$ ) has a haploid DNA content of around  $1.7 \times 10^{10}$  bp, which is almost 40 times that of rice. (**Bennett and Smith, 1976; Amuruganathan and Earle, 1991**). The world's most important centres of wheat and associated species variety are in Central Asia, the Near East, the Mediterranean, and Ethiopia. The Hindukush region is the epicentre of hexaploid wheat variability (**Kundu and Nagarajan, 1996**). Wheat is grown on around 221.24 million hectares worldwide, with a record yield of 771.64 million tonnes of grain and productivity is 3.49 metric tons per hectare (**USDA 2023**). India has the most wheat-growing land (14 percent), followed by Russia (12.43 percent), China (11.14 percent), and the United States (6.90 percent), accounting for around 45 percent of the global total. China, on the other hand, is the world's largest wheat producer, with 136 million tonnes produced, followed by India (98.51 million tonnes), Russia (85 million tonnes), and the United States (47.35mt). Global wheat production in 2022 is predicted to decline from the 2021 record level by 0.8 per cent, reaching 771.64 million tonnes and marking the first drop in four years. Year-on-year falls in production in Australia, India, Morocco and Ukraine will likely outweigh expected increases in Canada, Iran and Russia Further; it said that in Asia, wheat production in India is forecast at 105.5 million tonnes, down nearly 4 per cent from the record crop gathered in 2021. (**Business Standard 2022-23**).

Wheat production in 2022-23 is expected to be between 98 million and 106 million tonnes, down from 107.9 million tonnes in 2020-21 (**USDA report, 2022**). Wheat demand is anticipated to rise by 50% by 2050 compared to current levels. Meanwhile, new and more aggressive pests and viruses, dwindling water resources, limited accessible land, and unpredictable weather are all threatening the crop (heat in particular). For Africa, Asia, and Latin America, the CIMMYT's Global Wheat Program is one of the most significant public sources of high-yielding, nutritious, disease-resistant, and climate-resilient wheat varieties (**Wheat Research, CIMMYT**).

The success of our wheat varieties is up to a considerable extent due to incorporation of the Norin 10 genes *i.e.*  $Rht_1$  and  $Rht_2$  in wheat. These dwarfing genes changed the wheat plants type and it becomes more responsive to higher application of fertilizer and better crop management under practices. **In 1966, Dr. N.E. Borlaug**, a noble laureate introduced the Mexican dwarf wheat genotypes and provides the way for green revolution in India.

In most of the biometrical approaches for genetic evaluation of the crop, the diallel cross analysis became a proved and important system to provide maximum information on genetic parameters related to breeding programme of some important metric traits within considerable short time. To judge the stability performance over a wide range of environments, diallel cross analysis simultaneously evaluates the potentialities of the variance and predict the desirable types for further breeding programme. Various models of genetic analysis of diallel crosses have been given by **Jinks and Hayman 1953; Hayman 1954 a; 1954 b, Griffing, 1956 b; Gardner and Eberhart, 1966** and found suitable under limit facilities for achieving maximum genetic information.

## MATERIALS AND METHODS

The present investigation entitled “**Genetic studies for yield and its contributing characters in bread wheat (*Triticum aestivum* L.)** was conducted at Oil Seed Farm, Kalyanpur, C.S. Azad University of Agriculture and Technology, Kanpur-208002 (U.P.) during *Rabi*, 2021-22. The experimental material for present investigation comprised of forty five F<sub>1</sub>s developed by crossing 10 lines *viz.*, DBW187, K1601, HD2967, HD3249, DBW321, K1317 K0307, HI 1563, DBW107 and HD3059 following half diallel mating design. A total of 100 treatments with 10 parents (45 F<sub>1</sub>s and 45 F<sub>2</sub>s) were evaluated for the study of twelve quantitative characters in wheat.

### DIALLEL ANALYSIS:

#### Testing the validity of the hypothesis:

To test the validity of the hypothesis, i.e., the assumptions regarding diallel analysis as proposed by **Hayman (1954)**, such as (i) diploid segregation (ii) no maternal effect, (iii) no linkage (iv) no multiple allelism, (v) independent action of non-allelic genes and (vi) homozygosity of parents, the  $t^2$  test was applied as suggested by **Hayman (1954a)**:

$$t^2 = (n-2)/4 [(Var V_r - Var W_r)^2 / Var V_r \times Var W_r - Cov^2 (V_r, W_r)]$$

which is an F test with 4 and (n-2) degree of freedom.

A significant value of  $t^2$  would indicate the non-uniformity of  $W_r$ ,  $V_r$  and thus, invalidates the hypothesis postulated. The failure of hypothesis is also indicated by non-significant regression coefficient.

$$b = \frac{\text{Cov}(W_r, V_r)}{\text{Var}(V_r)}$$

Where,

$$\text{Cov.}(W_r, V_r) = \left[ \sum V_r W_r - \frac{\sum V_r \sum W_r}{n} \right] / (n-1) \text{ and}$$

$$\text{Var}(V_r) = \left[ \sum V_r^2 - \frac{(\sum V_r)^2}{n} \right] / (n-1)$$

The standard error of regression coefficient (b) was calculated as:

$$\text{SE}(b) = \left[ (\text{Var } W_r - b \text{ Cov. } W_r - V_r) / \text{Var } V_r (n-2) \right]^{0.5}$$

Where,

$$N = \text{number of parents}$$

Now the significance of differences 'b' from zero and unity was tested by using 't' value of (b-0)/SE (b) and (1-b)/SE (b) with (n-2) degree of freedom.

### Variance component analysis:

The components of variance in diallel cross were computed in F<sub>1</sub> by the use of equation given by **Hayman (1954a)**.

Expectation for F<sub>1</sub> diallel crosses is as follows:

$$V_p = \hat{D} + \hat{E}$$

$$V_r = \left( \frac{1}{4} \right) \hat{D} + \left( \frac{1}{4} \right) \hat{H}_1 - \left( \frac{1}{4} \right) \hat{F} + [(n+1)/2n] \hat{E}$$

$$W_r = \left( \frac{1}{2} \right) \hat{D} - \left( \frac{1}{4} \right) \hat{F} + (1/n) \hat{E}$$

$$V_m = \left( \frac{1}{4} \right) \hat{D} + \left( \frac{1}{4} \right) \hat{H}_1 - \left( \frac{1}{4} \right) \hat{H}_2 - \left( \frac{1}{4} \right) \hat{F} + (1/2n) \hat{E}$$

**Jinks (1956) and Hayman (1958)** gave expectations for F<sub>2</sub> diallel crosses. The expected statistics for F<sub>2</sub> generation are the same of that of F<sub>1</sub> except the contribution of h which is halved by one generation of inbreeding. Hence, the coefficient of H<sub>1</sub> and H<sub>2</sub> are (1/4) of those F<sub>1</sub> statistics while the coefficient of F is halved being second and first degree statistics h<sup>2</sup>, respectively (**Jinks, 1956; Hayman 1958; Mather and Jinks, 1971**). These expectations are as follows:

$$V_p = \hat{D} + \hat{E}$$

$$V_r = \left( \frac{1}{4} \right) \hat{D} + \left( \frac{1}{16} \right) \hat{H}_1 - \left( \frac{1}{8} \right) \hat{F} + [(n+1)/2n] \hat{E}$$

$$W_r = \left( \frac{1}{2} \right) \hat{D} - \left( \frac{1}{8} \right) \hat{F} + (1/n) \hat{E}$$

$$V_m = (1/4)\hat{D} + (1/16)\hat{H}_1 - (1/16)\hat{H}_2 - (1/8)\hat{F} + (1/2n)\hat{E}$$

Where,

$$\begin{aligned}\hat{D} &= \text{Components of variation due to additive effects of genes.} \\ &= V_0L_0 - \hat{E}\end{aligned}$$

$$\begin{aligned}\hat{H}_1 &= \text{Components of variation due to dominance effects of genes.} \\ &= V_0L_0 - 4W_0L_{01} + 4V_1L_1 - (3n-2)\hat{E}/n\end{aligned}$$

$$\hat{H}_2 = \hat{H}_1 [1-(u-v)^2] = 4V_1L_1 - 4V_0\hat{L}_1 - 2E$$

Where

$$U = \text{Proportion of positive genes in the parents.}$$

$$v = \text{Proportion of negative genes in the parents}$$

$$\hat{F} = \text{The mean of } F_r \text{ over the arrays}$$

$$F_r = 2(V_0L_0 - 4W_0L_{01} + V_1L_1 - W_r - V_r) - 2(n-2)\hat{E}/n$$

$$\hat{h}^2 = \text{Dominance effects (as the algebraic sum over all loci in heterozygous phase in all crosses)}$$

$$4(M_{L1} - M_{L0})^2 - 4(n-1)\hat{E}/n^2$$

$$\hat{E} = \text{the expected environmental component of variation}$$

$$(\text{Error SS} + \text{Replication SS/d.f.})/\text{number of replication}$$

In order to estimate of the accuracy of the components ( $\hat{D}, \hat{F}, \hat{H}_1, \hat{H}_2, \hat{h}^2$  and  $\hat{E}$ ) of variance, the term of main diagonal of matrix given **Hayman (1954)** with common multipliers  $S^2/n^5$ , was used.

Where,

$$S^2 = \frac{1}{2} \text{var. } (W_r - V_r). \text{ The formula being:}$$

$$SE(\hat{D}) = \pm [S^2 (n^5 + n^4)/n^5]^{0.5}$$

$$SE(\hat{F}) = \pm [S^2 (4n^5 + 20n^4 - 16n^3 + 16n^2)/n^5]^{0.5}$$

$$SE(\hat{H}_1) = \pm [S^2 (n^5 + 41n^4 - 12n^3 + 4n^2)/n^5]^{0.5}$$

$$SE(\hat{H}_2) = \pm [S^2 (36n^4)/n^5]^{0.5}$$

$$SE(\hat{h}^2) = \pm [S^2 (16n^2 + 16n^2 - 32n + 16n)/n^5]^{0.5}$$

$$SE(\hat{E}) = \pm [S^2 (n^4/n^5)]^{0.5}$$

After testing the significance of the components of variation, the mean degree of dominance was calculated as  $(\hat{H}_1/\hat{D})^{0.5}$  in  $F_1$  and  $[0.25 (\hat{H}_1/\hat{D})]^{0.5}$  in  $F_2$  generation. The proportion of genes with

positive and negative effects was calculated as  $H_2/4\hat{H}_1$ , the proportion of dominant and recessive genes in parents as the ratio of  $[(4\hat{D}\hat{H}_1)^{0.5} + 0.5\hat{F}] / [(4\hat{D}\hat{H}_1)^{0.5} - 0.5\hat{F}]$  in  $F_1$  and  $[0.25(4\hat{D}\hat{H}_1)^{0.5} + 0.5\hat{F}] / [0.25(4\hat{D}\hat{H}_1)^{0.5} - 0.5\hat{F}]$  in  $F_2$  generation, the number of gene groups which control the character and exhibit dominance as  $h^2 / \hat{H}_2$  and the coefficient of correlation between the parental order of dominance ( $W_r + V_r$ ) and parental measurement ( $Y_r$ ) as  $r$ .

### Heritability

**Heritability (in narrow sense)** in  $F_1$  generation was calculated by the formula proposed by **Crumpacker and Allard, (1962)**, which is as follows:

$$\text{Heritability } (\hat{h}^2) = (1/4) \hat{D} / [(1/4) \hat{D} + (1/4) \hat{H}_1 - (1/4) \hat{F} + \hat{E}]$$

Heritability in  $F_2$  generation was calculated according to the methodology proposed by **Verhalen and Murray (1969)**.

$$(\hat{h}^2) = (1/4) \hat{D} / [(1/4) \hat{D} + (1/16) \hat{H}_1 - (1/8) \hat{F} + \hat{E}]$$

or

$$1/2D/VP$$

Where,

$\hat{h}^2$  = Estimates of heritability coefficient and  $\hat{D}$ ,  $\hat{H}_1$ ,  $\hat{F}$  and  $\hat{E}$  are the same as explained earlier.

D = additive genetic variance

VP = phenotypic variance

The estimates of heritability and genetic advance were arbitrarily categorized in three classes by **Robinson in 1966** as:

- (i) High- above 30%
- (ii) Moderate- below (30-10)%
- (iii) Low below- 10%

$$\text{Heritability (\%)} = \text{Heritability coefficient} \times 100$$

### Genetic Advance:

The genetic advance was calculated by the formula given by **Robinson et al. (1949)** as:

Genetic advance (GA) = (k)x( $\hat{h}^2$ )x( $\sigma_{ph}$ ), and

Genetic advance over mean of the character

$$[GA (\%)] = \frac{GA}{\bar{x}} \times 100$$

Where

GA	=	Estimate of genetic advance
K	=	Selection differential at 5% selection intensity, i.e. 2.06
$\sigma_{ph}$	=	Phenotypic standard deviation
$\hat{h}^2$	=	Heritability coefficient in narrow sense
$\bar{X}$	=	Mean of the character concerned

#### **Genotypic and Environmental variances:**

Computed from the respective mean squares following the procedures suggested by Singh and Chaundhary (1979) and Allard (1960), thus

Genotypic variance

$$\sigma_g^2 = \frac{MSg - MSgl}{rl}$$

Genotypic by environment interaction variance

$$\sigma_{gl}^2 = \frac{MSgl - MSe}{rl}$$

Phenotypic variance

$$\sigma_p^2 = \sigma_g^2 + \left(\frac{\sigma_{2e}}{rl}\right) + \left(\frac{\sigma_{2gi}}{l}\right)$$

where,

MS<sub>g</sub> = mean square of genotype;

MS<sub>gl</sub> = meansquare due to genotype by environment interaction;

MS<sub>e</sub> = error mean square (mean square of environment);

l = number of locations;

r = number of replications.

**Genotypic (GCV), Phenotypic (PCV) and Environment (ECV) coefficients of variation (%)** estimated according to the procedure outlined by Johnson et al (1955):

$$\text{GCV}(\%) = \frac{\sqrt{\sigma^2_g}}{\bar{X}} \times 100$$

$$\text{PCV}(\%) = \frac{\sqrt{\sigma^2_p}}{\bar{X}} \times 100$$

$$\text{ECV}(\%) = \frac{\sqrt{\sigma^2_{gI}}}{\bar{X}} \times 100$$

Where,

$\sigma^2_g$  = Genotypic variance

$\sigma^2_p$  = Phenotypic variance

$\sigma^2_{gI}$  = Genotypic by environment interaction variance

$\bar{X}$  = Mean of the character concerned

The range of GCV, PCV and ECV suggested by **Subramanian & Menon (1973)**:

GCV, PCV & ECV variability categorized into three classes :-

Low = <10 %,

Moderate = 10-20%

High = > 20 %.

### **Correlation coefficients analysis:**

The estimates of phenotypic and genotypic correlation were worked out as given under:

### **Genotypic correlation**

$$r_{xy}(g) = \text{Cov}_{.xy}(g) / [V_x(g) \cdot V_y(g)]^{0.5}$$

Where,

$\text{Cov}_{.xy}(g)$  = genotypic covariance between character X and y was obtained as follows:

$$\text{Cov}_{.xy}(g) = [\text{Cov}_{.xy}(p) - \text{Cov}_{.xy}(e)]/r$$

$V_x (g)$  and  $V_y (g)$  = Genotypic variance for the characters x and y respectively

r = number of replications.

### Phenotypic correlation:

$$r_{xy} (p) = \text{Cov.}_{xy} (p) / [V_x (p) \cdot V_y (p)]^{0.5}$$

Where,

$\text{Cov.}_{xy} (p)$  = Phenotypic correlation between the character x and y and this was obtained as follows:

$$\text{Cov.}_{xy} (p) = \text{Cov.}_{xy} (g) + \text{Cov.}_{xy} (e)$$

$V_x (p)$  and  $V_y (p)$  = Phenotypic variance for the characters x and y, respectively.

$xy (e)$  = the error variance obtained from the ANNOVA of x and y characters.

## RESULTS

### HERITABILITY AND GENETIC ADVANCE:

The understanding of genetic variability present in a given crop species for the traits under improvement was imperative for the success of any plant breeding program (Sankar *et al*, 2006). The parameters such as genotypic and phenotypic coefficients of variation (GCV and PCV) are useful in detecting the amount of variability present in a given characteristic. The efficiency with which genotypic variability can be exploited by selection depends upon heritability and the genetic advance (GA) of individual trait (Bilgin *et al*, 2010). Genetic improvement of plants for quantitative traits requires reliable estimates of heritability in order to plan an efficient breeding program (Akinwale *et al*, 2011). Heritability provides information on the extent to which a particular morphogenetic character can be transmitted to successive generations (Bello *et al*, 2012).

### HERITABILITY:

Accordingly, the high magnitude of heritability (over 30%) were observed for the characters, days to 75% flowering (84.53), plant height (93.05), number of tillers per plant (94.35), number of spikelets per spike (91.48), spike length (87.27), number of grain per spike (95.14), days to

maturity (80.42), 1000 grain weight (97.20), ear density (68.54), duration of reproductive phase (75.32), protein content (98.70), and grain yield per plant (99.14) in F<sub>1</sub> generation. In F<sub>2</sub> generation high heritability (over 30%) were observed for the characters, days to 75% flowering (77.32), plant height (90.87), number of tillers per plant (92.87), number of spikelets per spike (89.37), spike length (86.23), number of grain per spike (94.45), days to maturity (47.44), 1000 grain weight (96.94), ear density (58.60), duration of reproductive phase (53.26), protein content (98.35), and grain yield per plant (99.08) (Figure 1).

### **GENETIC ADVANCE:**

In order to ascertain relative merit of different attributes, genetic advance in percent of the mean was worked out for all the twelve characters in both the generations. Estimates of genetic advance in per cent over mean ranged from 5.24 for days to maturity to 40.51 for grain yield per plant in F<sub>1</sub> generation. High genetic advance was observed for number of tillers per plant (25.64), grain yield per plant (40.51) duration of reproductive phase (22.50) in F<sub>1</sub> generation. In F<sub>2</sub> generation, range of genetic gain was recorded 3.61 for days to maturity to 41.04 for grain yield per plant. High genetic advance was observed for number of tillers per plant (26.25), grain yield per plant (41.04) and protein content (20.01) in F<sub>2</sub> generation (Figure 1).

### **GENOTYPIC, PHENOTYPIC AND ENVIRONMENT COEFFICIENT OF VARIANCE (%)**

Genotypic variation was the heritable portion of phenotypic or total variation. It's gives the variation between genotype. Environmental variation was the non-heritable portion of observable variation, suggested by **Subramanian & Menon (1973)**, GCV, PCV & ECV arbitrarily categorized into three classes - Low = <10 %, Moderate = 10-20% High= > 20 %. The estimates of GCV, PCV & ECV revealed that the values of PCV % were higher than the GCV % and ECV% for all characters in both generation F<sub>1</sub> and F<sub>2</sub>.

### **GENOTYPIC COEFFICIENT OF VARIATION (GCV %)**

Highest value of GCV was observed only in F<sub>2</sub> generation for grain yield per plant (20.01). Moderate value of GCV (%) were observed in both F<sub>1</sub> and F<sub>2</sub> generation for ), number of tillers per plant (F<sub>1</sub>-12.81, F<sub>2</sub>-13.22) and duration of reproductive phase (F<sub>1</sub>-12.58, F<sub>2</sub>-11.78) while low GCV were observed in both generation for days to 75% flowering (F<sub>1</sub>-3.97, F<sub>2</sub>-4.35), plant height (F<sub>1</sub>-5.91, F<sub>2</sub>-6.14) number of spikelets per spike (F<sub>1</sub>-5.48, F<sub>2</sub>-5.60), spike length (F<sub>1</sub>-6.88, F<sub>2</sub>-7.13), number of grain per spike (F<sub>1</sub>-8.85, F<sub>2</sub>-9.01), days to maturity (F<sub>1</sub>-2.84, F<sub>2</sub>-2.54), grain

weight (F<sub>1</sub>-7.51, F<sub>2</sub>-7.81), ear density (F<sub>1</sub>-4.15, F<sub>2</sub>-4.07), protein content (F<sub>1</sub>-9.60, F<sub>2</sub>-9.79), and grain yield per plant (F<sub>1</sub>-19.75) (Figure 2).

### **PHENOTYPIC COEFFICIENT OF VARIATION (PCV %)**

Highest value of GCV was observed only in F<sub>2</sub> generation for grain yield per plant (20.10). Moderate value of GCV (%) were observed in both F<sub>1</sub> and F<sub>2</sub> generation for ), number of tillers per plant (F<sub>1</sub>-13.19, F<sub>2</sub>-13.72) and duration of reproductive phase (F<sub>1</sub>-14.50, F<sub>2</sub>-16.15) while low GCV were observed in both generation for days to 75% flowering (F<sub>1</sub>-4.31, F<sub>2</sub>-4.94), plant height (F<sub>1</sub>-6.13, F<sub>2</sub>-6.45) number of spikelets per spike (F<sub>1</sub>-5.73, F<sub>2</sub>-5.92), spike length (F<sub>1</sub>-7.37, F<sub>2</sub>-7.68), number of grain per spike (F<sub>1</sub>-9.08, F<sub>2</sub>-9.27), days to maturity (F<sub>1</sub>-3.16, F<sub>2</sub>-3.69), grain weight (F<sub>1</sub>-7.61, F<sub>2</sub>-7.94), ear density (F<sub>1</sub>-5.02, F<sub>2</sub>-5.32), protein content (F<sub>1</sub>-9.66, F<sub>2</sub>-9.87), and grain yield per plant (F<sub>1</sub>-19.83).

### **ENVIRONMENTAL COEFFICIENT OF VARIATION (ECV %)**

The estimates of ECV (%) were observed low to very low in both F<sub>1</sub> and F<sub>2</sub> generation for all twelve characters, namely days to 75% flowering (F<sub>1</sub>-1.79, F<sub>2</sub>-2.11), plant height (F<sub>1</sub>-1.40, F<sub>2</sub>-1.44), number of tillers per plant (F<sub>1</sub>-2.43, F<sub>2</sub>-1.88), number of spikelets per spike (F<sub>1</sub>-6.38, F<sub>2</sub>-6.96), spike length (F<sub>1</sub>-5.39, F<sub>2</sub>-5.75), number of grain per spike (F<sub>1</sub>-4.73, F<sub>2</sub>-6.02), days to maturity (F<sub>1</sub>-3.15, F<sub>2</sub>-2.71), 1000 grain weight (F<sub>1</sub>-2.62, F<sub>2</sub>-2.70), ear density (F<sub>1</sub>-1.82, F<sub>2</sub>-1.56), duration of reproductive phase (F<sub>1</sub>-1.56, F<sub>2</sub>-1.23), protein content (F<sub>1</sub>-2.21, F<sub>2</sub>-1.36), and grain yield per plant (F<sub>1</sub>-1.09, F<sub>2</sub>-0.57) (Figure 2).

### **DISCUSSION**

The breakthrough in yield of wheat was achieved after Borlaug acquired a variety of dwarf wheat from Japan and cross-bred with North American strains to produce a semi-dwarf strain with a thicker, stronger stalk, capable of supporting a heavier load of grains. Crossing these with his rust-resistant strains produced ideal wheat for Mexico's needs. As his breeding techniques grew more and more sophisticated, he realized the tall thin stalks of wheat had been growing too frequently collapsed under the weight of their own grain, and success of high yielding, semi dwarf, disease resistant, lodging tolerant, photo-insensitive, fertilizer responsive genotypes carrying the 'NORIN-10' genes. Various biometrical approaches are now available for the estimation of different parameters. This is obviously a **hit-and-miss approach** depending upon the breeder's visual and phenotypic judgment and his experience rather than genetic architecture. In this experiment parents, namely., DBW187, K1601, HD2967, HD3249, DBW321, K1317 K0307, HI 1563, DBW107 and HD3059 were taken out for evaluation of different yield

attributing traits. In this present investigation the models viz., variance components analysis (**Hayman, 1954a**), combining ability analysis (**Griffing, 1956b**) and heterosis parameters (Gardner and Eberhart, 1966) have been used for different genetic analysis.

## **HERITABILITY AND GENETIC ADVANCE**

Heritability estimates have been utilized as a selection criterion in the plant improvement program. It has been widely used to estimate the degree to which a character may be transmitted from parent to off spring. In narrow sense, heritability is the ratio of additive genetic variance to the phenotypic variance. Genetic advance, though not an independent entity, has added advantage over heritability where character is to be improved through a series of segregating generations. **Johnson et al. (1965)** stated that without genetic advance, the estimates of heritability will not be of much practical importance and emphasized the importance of both the genetic advance as well as heritability estimates in selection breeding programme. According to Hanson (1963), heritability and genetic advance are two complementary concepts.

Accordingly, the high magnitude of heritability (over 30%) were observed for all the characters showed high heritability in F1 generation and days to 75% flowering (77.32), plant height (90.87), number of tillers per plant (92.87), number of spikelets per spike (89.37), spike length (86.23), number of grain per spike (94.45), days to maturity (47.44), 1000 grain weight (96.94), ear density (58.60), duration of reproductive phase (53.26), protein content (98.35), and grain yield per plant (99.08). in F2 generation(**Figure 1**). Similar estimates was also reported by **Sattar et al. (2003)**, for plant height **Aycicek and Yildirim (2006)**, **Memon et al. (2007)**, **Waqar-UI-Haq et al. (2008)**, **Mohsin et al. (2009)**, **Kamboj et al. (2010)**, **Khalid et al. (2011)**, **Verma et al. (2012)**, **Singh et al. (2017)**, **Rathwa et al. (2018)**, **umar and Kumar (2021)** for grain weight per spike; high estimates of heritability were due to greater contribution of additive genetic component. It indicates that if these characters are subjected to progeny and any other selection scheme for exploiting fixable genetic variance, a wide adopted genotype could be developed. Selection pressure should be exercised in early generations. **Singh et al. (1991)**, **Katiyar (2003)**, **Gupta et al. (2004)** and **Safi et al. (2017)**

The estimates of heritability was low (below 10%) for the character spike length in F2 generation. It indicated that selection would be more effective in early segregating generations. Similar result observed by **Ibrahim and Quick (2001)**, **Sattar et al. (2003)**, **Ahmed et al. (2004)**, **Aycicek and Yildirim (2006)**, **Memon et al. (2007)**, **Waqar-UI-Haq et al. (2008)**, **Saleem et al. (2016)**, **Malbhage et al. (2020)** and **Kumar and Kumar (2021)**.

High genetic advance was observed for number of tillers per plant (25.64), grain yield per plant (40.51) duration of reproductive phase (22.50) in F1 generation. High genetic advance was observed for number of tillers per plant (26.25), grain yield per plant (41.04) and protein content (20.01) in F2 generation. It indicated that manifestation of these traits was primarily governed by additive genetic effects which were fixable, and the desired selection gain could be achieved in

early generations. Moderate genetic advance was reported by **Kumar et al. (1991) and Yadav and Singh (2002)**, for number of grains per spike and biological yield; Singh et al. (1991), Alpay Balkan (2018), **Rathwa et al. (2018)** for biological yield and number of productive tillers per plant.

Genotypic variation is the heritable portion of phenotypic or total variation. Environmental variation is the non-heritable portion of observable variation, suggested by **Subramanian & Menon (1973)**. GCV, PCV & ECV were categorized into three classes, viz., low (less than 10%), moderate (10-20%) and high (more than 20%). Highest value of GCV was observed only in F2 generation for grain yield per plant (20.01) (**Figure 2**). **Similar results were also reported by Bhushan et al. (2013), Dutamo et al. (2015), Sarfraz et al. (2016), Arya et al. (2017), Jaiswal et al. (2019), (2021), Kumar and Kumar (2021).**

## REFERENCES


1. **Akinwale, M. G., Gregorio, G., Nwilene, F., Akinyele, B. O., Ogunbayo, S. A., & Odiyi, A. C. (2011)**. Heritability and correlation coefficient analysis for yield and its components in rice (*Oryza sativa* L.). *African Journal of plant science*, **5**(3), 207-212.
2. **Allard, R.W. (1960)**. Components of genetic variances, Principles of Plant Breeding. *John Wiley and Sons Inc.*, New York, USA, p. 485.
3. **Amuruganathan, E. and Earle, E.D. (1991)**. Nuclear DNA content of some important plants species. *Plant Mol. Biol. Rep.*, **9**: 208-218.
4. **Arya, V.K.; Singh, J.; Kumar, L.; Kumar, R.; Kumar, P. and Chand, P. (2017)**. Genetic variability and diversity analysis for yield and its components in wheat (*Triticum aestivum* L.). *Ind. J. Agric. Res.*, **51**(2): 128-134.
5. **Arya, V.K.; Singh, J.; Kumar, L.; Kumar, R.; Kumar, P. and Chand, P. (2017)**. Genetic variability and diversity analysis for yield and its components in wheat (*Triticum aestivum* L.). *Ind. J. Agric. Res.*, **51**(2): 128-134.
6. **Asif, R.; Mujahid, M.Y.; Kisana, N.S.; Mustafa, S.Z. and Ahmad, I. (2004)**. Heritability, genetic variability and path-coefficient of some traits in spring wheat. *Sar. J. of Agril. Pakistan*, **20** (1): 87-91.
7. **Aycicek, M. and Yildirim, T. (2006)**. Heritability of yield and yield component in bread wheat (*Triticum aestivum* L.) genotypes. *Bangl. J. Bot.*, **35** (1): 17-22.

8. **Bello, O. B., Ige, S.A., Azeez, M.A., Afolabi, M.S., Abdul-maliq, S.Y., Mahamood, J., (2012).** Heritability and genetic advance for grain yield and its component character in Maize (*Zea mays* L.). *International Journal of Plant Research*, 2: 138–145. DOI: 10.5923/j.plant.20120205.01
9. **Bennett, M.D. and Smith, J.B. (1976).** Nuclear DNA amounts in angiosperms. *Phil. Trans. R. Soc. Lond. B.*, **274** (933): 227-274.
10. **Bhushan, B.; Gaurav, S.S.; Kumar, R.; Pal, R.; Pandey, M.; Kumar, A.; Bharti, S.; Nagar, S.S. and Rahul, V.P. (2013).** Genetic variability, heritability and genetic advance in bread wheat (*Triticum aestivum* L.). *Envi. & Eco.*, **31** (2): 405-407.
11. **Bilgin O, Korkut KZ, Baser I, Dalioglu O, Ozturk I, Kahraman T, Balkan A, (2010).** Variation and Heritability for Some Semolina Characteristics and Grain Yield and their Relations in Durum Wheat (*Triticum durum* Desf.). *World J Agric Sci* 6: 301- 308
12. **Borlaug, N.E. (1966).** Preliminary reports of the first three Inter-American and the first two Near East-American spring wheat yield nurseries. *Biblio. Infor.*, **130**(2): 1-5.
13. [Business-Standard.com/article/international/with-ban-exceptions-india-s-wheat-export-forecast-to-be-7-mt-in-2022-23-122061000357\\_1.html](https://www.business-standard.com/article/international/with-ban-exceptions-india-s-wheat-export-forecast-to-be-7-mt-in-2022-23-122061000357_1.html)
14. **Crumpacker, D. and Allard, R. (1962).** A diallel cross analysis of heading date in wheat. *Hilga*. **32**(6): 275-318.
15. **Dutamo, D.; Alamerew, S.; Eticha, F. and Assefa, E. (2015).** Genetic variability in bread wheat (*Triticum aestivum* L.) germplasm for yield and yield component Traits. *Journal of Biology, Agriculture and Healthcare*, **5**(17): 140–147.
16. **Gardner, C.D. and Eberhart, S.A. (1966).** Analysis and interpretation of the variety cross diallel and related population. *Biometrics.*, **22**: 439-452.
17. **Griffing, B. (1956b).** Concept of general and specific combining ability in relation to diallel crossing system. *Australian J. Bioll Sci*, **9**:463-493.
18. **Gupta, S. K., Rathore, P., & Singh, K. (2004).** Genetic variability in mungbean [*Vigna radiata* (L.) Wilczek]. *Agric Sci Digest*, **24**, 136-138.
19. **Hayman, B. I. (1954).** The analysis of variance of diallel tables. *Biometrics*, **10**(2), 235-244.
20. **Hayman, B.I. (1954a).** The theory and analysis of diallel crosses. *Genetics*, **39**: 789-809

21. **Hayman, B.I. (1954b).** The analysis of variance of diallel tables. *Biometrics*, **10**:235-244.
22. **Hayman, B.I. (1958).** The theory and analysis of diallel crosses –II *Genetics*, **43**:63-85.
23. **Ibrahim, A.M.H. and Quick, J.S. (2001).** Heritability of heat tolerance in winter and spring wheat. *Crop Sci.*, **41**(5): 1401-1405.
24. **Jaiswal, A., Singh, V., Lal, K., Prasad, D., Yadav, K., & Yadav, V. P. (2019).** Study on genetic variability and divergence under sodic soil in indigenous lines of wheat (*Triticum aestivum* L. em. Thell). *Journal of Pharmacognosy and Phytochemistry*, **8**(3), 1752-1756.
25. **Jinks, J.L. and Hayman, B.I. (1953).** The analysis of diallel cross. *Maize Genet. Newsletter*, **27**: 48-54.
26. **Jinks, J.L., (1956).** The F<sub>2</sub> and back cross generations from a set of diallel crosses. *Heredity*, **10**: 1-30.
27. **Johnson, H.W.; Robinson, H.F. and Comstock, R.E. (1955).** Genotypic and phenotypic correlations in soybean and their implications in selection. *Agron. J.*, **47**: 477-483.
28. **Kamboj, R.K. (2010).** Genetic variability, heritability and genetic advance in bread wheat (*Triticum aestivum* L.em. Thell.) under salinity stress conditions. *Madras Agric. J.*, **97** (1-3): 29-30.
29. **Katiyar, M. (2003).** Study of heritability and genetic advance over environments in bread wheat (*Triticum aestivum* L.). *Farm Sci. J*, **12**(2), 176-177.
30. **Khalid, M; Khalil, I.H.; Farhatullah; Bari, A.; Tahir, M.; Ali, S.; Anwar, S.; Ali, S. and Ismail, M. (2011).** Assessment of heritability estimates for some yield traits in winter wheat (*Triticum aestivum* L.). *Pak. J. Bot.*, **43** (6): 2733-2736.
31. **Kumar, A., Ahmad, Z. and Singh K. N. (1991).** selection parameters for yield and quality attributes in breed wheat Proc. Golden Jubilee symp. *Indian Soc. Genet. Plant Breed.*, New Delhi, Abstr. **II**, p: 347
32. **Kumar, M. and Kumar, S. (2021).** Estimation of heritability and genetic advance in 24 genotypes of bread wheat (*Triticum aestivum* L.). *J. of Pharma. and Phytochem.*, **10** (1): 1110-1113.

- 33. Kundu S, Nagarajan S. (1996).** Distinguishing characters of Indian wheat varieties. Research Bulletin No. 4. Directorate of Wheat Research. Karnal, India.
- 34. Malbhage, A.B.; Malbhage, M.M.; Shekhawat, V.S. and Mehta, V.R. (2020).** Genetic variability, heritability and genetic advance in durum wheat (*Triticum durum* L.). *Journal of Pharma. and Phytochem.*, **9** (4): 3233-3236.
- 35. Mather, K. and Jinks, J.L. (1971).** Biometrical Genetics. 2<sup>nd</sup> (Ed.) Chapman and Hall Ltd., London.
- 36. Memon, S.; Qureshi, M.; Ansari, B.A. and Sial, M.A. (2007).** Genetic heritability for grain yield and its related characters in spring wheat (*Triticum aestivum* L.) *Pak. J. Bot.*, **39** (5): 1503-1509.
- 37. Mohsin, T; Khan, N. and Naqvi, F.N. (2009).** Heritability, phenotypic correlation and path coefficient studies for some agronomic characters in synthetic elite lines of wheat. *Journal of Food, Agric. & Env.*, **7** (3&4): 278-282.
- 38. Rathwa, H.K.; Pansuriya, A.G.; Patel, J.B. and Jalu, R.K. (2018).** Genetic variability, heritability and genetic advance in durum wheat (*Triticum durum* Desf.). *Int. J. Curr. Microbiol. App. Sci.*, **7** (1): 1208-1215.
- 39. Robinson, H.F.; Comstock, R.E. and Harvey, P.H. (1949).** Estimates of heritability and the degree of dominance in corn. *Agron. J.*, **41**(8): 353-359.
- 40. Safi, L.; Singh, R. and Abraham, T (2017).** Assessment of heritability and genetic parameters in wheat (*Triticum aestivum* L.) based on agronomic and morphological traits. *Journal of Pharmacognosy and Phytochemistry*, **6** (5): 18-21.
- 41. Saleem, B.; Khan, A.S.; Shahzad, M.T. and Ijaz, Fahid (2016).** Estimation of heritability and genetic advance for various metric traits in seven F<sub>2</sub> populations of bread wheat (*Triticum Aestivum* L.). *Journal of Agricultural Sciences*, **61** (1): 1-9.
- 42. Sankar PD, Sheeba A, Anbumalarmathi J, (2006).** Vari-ability and Character Association Studies in Rice. *Agric Sci Digest* 26: 182-184
- 43. Sarfraz, Z., Shah, M. M., & Iqbal, M. S. (2016).** Genetic variability, heritability and genetic advance for agronomic traits among A-genome donor wheat genotypes. *J. Agric. Res*, **54**(1), 15-20.

- 44. Sattar, A.; Chowdhry, M.A. and Kashif, M. (2003).** Estimation of heritability and genetic gain of some metric traits in six hybrid population of spring wheat. *Asian J. of Plant Sci.*, **2** (6): 495-497.
- 45. Satyawart; Yadav, R.K. and Singh, G.R. (2002).** Variability and heritability estimates in bread wheat. *Env. and Eco.*, **20** (3): 548-550.
- 46. Singh R.K. and Chaudhary, B.D. (1979).** Biometrical methods in quantitative genetic analysis. pp. 9-10. KalyaniPublishers, New Delhi, India.
- 47. Singh, T. and Rai, R.K. (1991).** Genetics of yield and some morpho-physiological traits in bread wheat. Paper presented in golden Jubilee Symposium Indian Soc. *Genet. P.I. Breed New Delhi* Abstr. **II**, p. 357.
- 48. Sivasubramanian, S. and Menon, (M. 1973).**Heterosis and inbreeding depression in rice. *Madras Agric. J* 60:1139.
- 49. United States Department of Agriculture (USDA).** Released (July 12, 2022) by the National Agricultural Statistics Service (NASS), Agricultural Statistics Board.
- 50. Verhalen, L. M., & Murray, J. C. (1969).** A Diallel Analysis of Several Fiber Property Traits in Upland Cotton (*Gossypium birsutum* L.) II 1. *Crop science*, **9**(3), 311-315.
- 51. Verma S., Singh S.V. and Singh M.P. (2012).** Heritability, genetic advance and correlation studies for quality and yield contributing traits in bread wheat (*Triticum aestivum* L.). *Progressive Res.* **7**(special) : 314-317.
- 52. Waqar-Ul-Haq; Malik, M. F.; Munir, M. and Akram, Z. (2008).** Evaluation and estimation of heritability and genetic advancement for yield related attributes in wheat lines. *Pak. J. Bot.*, **40** (4): 1699-1702.



**Figure 1: Estimates of grand mean (a), heritability (b), genetic advance(c) and genetic advance in per cent of mean (d) for 12 characters in a 10 parent diallel cross of F<sub>1</sub> and F<sub>2</sub> generation in wheat.**




Figure 2: Estimates of Grand mean(a), GCV(b), PCV (c)and ECV(d) for 12 different characters in wheat.