

Original Research Article

Effect of Different Drying Methods on Physico-Chemical Attributes of Guava's Fruit Pulp Powder

Comment [U1]: Properties, change this in whole text

Comment [U2]: Whole Manuscript should be corrected for a grammar mistakes from a native or professional English speaker. Text should be technically edited according to the Journal Guidelines, too.

ABSTRACT

Guava (*Psidium guajava L.*) is one of the most popular fruits in tropical and subtropical regions of India. It is popularly known as "Apple of tropics", belongs to the family Myrtaceae and often considered as "super fruit". It is rich in vitamin A and C, omega-3 and 6 polyunsaturated fatty acids and contains high levels of dietary fiber and other nutritional properties. Guava is a highly perishable fruit crop and has a very limited shelf life, so there is a need to preserve the fruits in dried form, whose powder can be used for processing purposes, value addition in off season demands. Therefore, we conducted an experiment to study the effect of different drying methods on physico-chemical attributes of guava's fruit pulp powder. Fruits of guava pulp were used to prepare powder by different drying methods like freeze drying, sun drying, air drying and oven drying. Results showed that bulk density and moisture percent was highest recorded in sun drying method followed by air drying while minimum was recorded with oven drying followed by freeze drying. Maximum nutritional value like Fiber (5.97%), Ash (1.38%), Protein percent (1.82%), Total sugar (32.88%), reducing (23.14%) & non-reducing sugar (9.73%), ascorbic acid (227.90 mg/100g), Titratable acidity (2.20%), Total phenols (250.66) and DPPH radical scavenging activity (83.40) was observed maximum with freeze drying followed by air drying. Taste, flavour, aroma and overall acceptability were better in freeze drying followed by Air drying.

Keywords: Guava, DPPH, Vitamin -C, Total Phenols, Pulp Powder, Drying Methods.

1. INTRODUCTION

The guava (*Psidium guajava L.*) is one of the most important commercial fruit crops in India and belong to the family Myrtaceace. It is believed to be native in central America

and the southern part of Mexico (Somogyi et.al. 1996). Guava fruit is considered as a highly nutritious fruit because it contains a high level of ascorbic acid (200-299mg/100g of fresh weight), which is five to six time higher than oranges. Guava is a very popular fruit for their easy adaptability in diverse agro-climatic condition. It is available throughout the year except during the summer season. Therefore, it is commonly referred to as 'Poor man's Apple' and Apple of the Tropics'. In india, it ranks fourth in total cultivated area after mango, banana, and citrus. Uttar Pradesh is the largest producer of guava in india and Allahabad has the reputation of producing the world's best guavas (Mitra and Bose, 1990).

It is also an excellent source of beta carotene, lycopene, calcium, potassium, zinc, and Soluble fiber. Guavas are very rich in antioxidants which can acts against the free radicals that damage cells and cause cancer, diabetes, and coronary disease. Guava can be consumed fresh or can be processed form like jelly, jam, juice, nectar, pulpand slices in syrup, fruit bar or dehydrated products, as well as being used as an additive to other fruit juices and pulp(Anonymous, 2014).

The major objective of drying is to preserve the fruits, it also lowers the cost of transportation and storage by reducing both weight and volume of final product. Freeze drying, is the best drying technique to produce highest quality food products in terms of appearance and retention of nutrient and sensory characteristics (Murthy and Joshi 2007).

Being perishable in nature, guava fruits had a very limited shelf life so there is need to preserve the fruits in dehydrated form. Therefore, present work was undertaken to study the Effect of different drying methods on Physico-Chemical attributes of Guava's fruit Pulp Powder prepared by different drying methods such as Freeze drying, Sun drying, Air drying and Oven drying and their effect on storage.

2. MATERIALS AND METHODS

2.1 Location of Experiment

Experiment was conducted in the University Laboratory, RVSKVV, Gwalior, MP, India during the period of February 2022 to March, 2022.

2.2 Collection of Fruit Samples

Fully mature fruits were collected from the guava Orchard, they were thoroughly washed under tap water to remove dust & impurities and segments were cut and separated from the seed with the help of knife.

Comment [U3]: And, avoid to use & in the formal text

2.3 Drying methods and preparation of fruit pulp powder

In Gwalior during February, it was mild cold sunny days when the fruits were dried under sun, for Sun drying fruit pulp was dried under the sun for five to six days. In Oven Drying the pulp was kept in the hot air oven at 55 °C and for Air drying, pulp pieces were kept under air circulated incubator at constant 25 °C until the moisture is reduced to minimum 10 %. For freeze drying, fruit pulp segments were kept at -40 °C for 5 hours and the freeze dried at -50°C and 52 pa pressure for 22 hours. The dried pieces were grinded to fine powder and filtered using muslin cloth. The powder was kept under plastic sealed bags and stored at 4 °C until the analysis (Verma et.al.,2015).

2.4 Physico-Chemical Analysis

2.4.1 Moisture, Bulk Density, Total soluble solids, and pH

The moisture content was determined by drying at 70°C up to constant weight and expressed in terms of the percent wet basis (100X kg water/kg wet material).Bulk density of powder was obtained by measuring the volume of a determined weight of the powder in a 100 ml graduated glass cylinder. The ratio of mass of the powder and the volume occupied in the cylinder determines the bulk density value(g/cm³)(Goula et al., 2004).Total soluble solids (TSS)(°Brix) of the juice of fresh pulp was determined by using hand refractometer and pH of pulp juice was determined by digital pH meter.

2.4.2 Ascorbic Acid, Titratable Acidity

Ascorbic acid (Vitamin-C) content of the sample was analyzed by using 2, 6-dichlorophenol indophenol dye method as described by Ranganna (2000). And expressed in mg per 100g.

Titrateable acidity (%) of guava powder samples was measured by AOAC, (2005) method by boiling the sample for 1 h in water and making up the volume up to 100 ml and then titrating it against 0.1 N sodium hydroxide solution using phenolphthalein indicator.

2.4.3 Ash, Fiber and Protein

Protein was estimated as per the method described by Lowry et al. (1951).

2.4.4 Total Sugar, Reducing Sugar and Non-Reducing Sugar

Total sugar and reducing sugar content were measured using methods of Lane and Eynon as described by Ranganna. Non-reducing sugar was estimated by subtracting reducing sugar from total sugar.

2.4.5 Total Phenols and DPPH Radical Scavenging Activity

The 10 g powder sample was homogenized along with 40 ml of methanol at a magnetic stirrer at 300 rpm, supernatant was increased by adding 50 ml methanol, and this extract was reduced in the volume by rotary vacuum evaporator. The concentrated solution was filtered through filter paper and was stored in plastic bottles at 4°C until analyzed. This extract was used to determine phenols and antioxidant activity.

Total phenolic content (TPC) in fresh or freeze-dried guava samples was determined by the Folin-Ciocalteu method (Singleton et al., 1999). The TPC was expressed as mg gallic acid equivalent (mg GAE/g) from the standard curve prepared by using pure gallic acid.

Free radical scavenging activity of extracts was measured by the slightly modified method of Allothman et al., 2009. The antioxidant capacity of the fruit extracts was studied through the evaluation of the free radical-scavenging effect on the 1, 1-diphenyl-2-picrylhydrazyl (DPPH) radical. An aliquot (100 µl) of fruit extract was mixed

with 3.9 ml of 0.1 mM DPPH methanolic solution. The mixture was thoroughly vortex-mixed and kept in the dark for 30 min. The absorbance was measured later, at 517 nm, against a blank of methanol and ascorbic acid as standard. Results were expressed as percentage of inhibition of the DPPH radical.

Sensory Analysis

Sensory analysis of samples was done at 9.0-point hedonic scale. Ten semi trained panelists aged between 21 and 45 years, were selected to take part in the sensory panel. The panel measured the taste, flavor, aroma and overall acceptability.

Statistical Analysis

The data were obtained in triplicates (n=3) for each parameter and their mean was calculated. Data significantly was analyzed statistically using one-way ANOVA. F-test was adopted for the level of significance at 5%. Standard error of difference (SEd) and Critical difference (CD) were also obtained.

3. Results and Discussion

3.1 Physico-Chemical Attributes of Fresh Guava's Fruit Pulp

The data represented in (table 1) showing the Physico-Chemical Attributes of Fresh Fruit of Guava Pulp that were used for preparation of dried guava powder. Analysis revealed that the moisture content of fresh guava fruit is (85.83%) is reported and Bulk density (0.97 g/cm³). The chemical analyses of the fresh fruit revealed that it contains pH, TSS (Total Soluble Solids) and titratable acidity was found 5.01, 11.04°Brix and 0.585% respectively. Guava having also a good amount of dietary fiber (2.88%), protein (0.50%), ash (0.66%), total sugar (10.85%), reducing sugar (7.64%), non-reducing sugar (3.21%) and ascorbic acid (280.66mg/100g).

Table 1. Physico-Chemical Attributes of Fresh Guava's Fruit Pulp

S.No.	Parameters	Mean ± SD*
1.	Weight of pulp per fruit (g)	111.6± 3.51
2.	Volume of pulp per fruit(cm ³)	114.46±1.24
3.	Bulk density(g/cm ³)	0.97±0.021
4.	Moisture%	85.83±0.40
5.	pH	5.01±0.68
6.	TSS(Total Soluble Solids)°Brix	11.04±0.26
7.	Titratable acidity%	0.58±0.04
8.	Dietary fiber%	2.88±0.35
9.	Protein%	0.50±0.05
10.	Ash%	0.66±0.02

Comment [U4]: This part should be corrected.

1. For all the Tables: under each table, mark clearly the statistical analysis, which variables are compared? Make it clear
2. The statistical analysis should be more discussed. Which parameters are statistically significant higher/lower compared to another one? Why is this important?
3. Different drying methods should be discussed in terms of their efficiency. Which one showed the best results and why? Which one is economically the best way for using and why? What are your recommendation according to this study?

11.	Total sugar%	10.85±0.53
12.	Reducing sugar%	7.64±0.43
13.	Non-reducing sugar%	3.21±0.19
14.	Ascorbic acid(mg/100g)	280.66±16.01

**Data presented in Mean± Standard Deviation*

3.2 Physico-chemical Attributes of guava's fruit pulp powder

3.2.1 Moisture and bulk density

The drying methods showed significant results, highest moisture content (6.73%) was found in sun drying followed by air drying (6.26%) methods (Table 2). Because drying under uncontrolled environments which cause more chances of environmental affect. Bulk density was reported highest in sun drying (0.76) followed by air drying (0.67) and both methods differed significantly. More moisture tends to put more bulk density .While oven drying possessed minimum moisture percent and bulk density because of temperature is higher than other methods and environmental condition is controlled.

Table 2. Effect of drying methods on physico-chemical attributes and proximate of guava'sfruit pulp powder

S. No	Drying methods	Parameters						
		Moisture %	Weight of powder per fruit (g)	Volume of powder per fruit (cm3)	Bulk density(g/cm3)	Ash%	Fiber %	Protein %
1.	Sun drying	6.73	41.56	54.2	0.76	1.04	4.52	0.75
2.	Oven drying	3.85	17.52	36.17	0.48	1.14	4.94	0.94
3.	Air drying	6.26	31.56	46.67	0.67	1.27	5.53	1.06
4.	Freeze drying	4.17	24.86	42.94	0.57	1.38	5.97	1.82
	SEd±	0.15	0.84	1.40	0.02	0.01	0.51	0.08
	C.D.(P=.05)	0.35	1.95	3.23	0.06	0.04	1.19	0.20

3.2.2 Ash, Fiber and protein

Ash, fiber and protein content were significantly differed among different drying methods. However, oven drying &air-drying methods were found at par with each other.

Significantly highest content of ash (1.38), fiber (5.97) and protein (1.82) were found in pulp powder when freeze dried followed by air dried. The minimum content was found with sun drying method (Table 2). As opposed to sun drying, which is influenced by environmental fluctuation, and oven drying, where temperature is high and more component degradation occurs, freeze drying and air-drying methods are characterised by low temperature treatment under controlled conditions that eliminates environmental fluctuation and results in minimal component degradation, such as ash, fibre and protein. Results are in accordance with the reports of (Alkandari et al.), (Karam et al.); (Radojčin et al.), (Shuen et al.); (Verma et al.), they reported freeze drying method as best as compared to other methods like oven drying, sun drying etc.

Comment [U5]: Units? Check this in whole Discussion part

3.2.3 Titratable acidity, total sugar, reducing and non-reducing sugar

Significant results were obtained. The maximum titratable acidity (2.20), total sugar (32.88), reducing sugar (23.14) and non-reducing (9.73) were found with freeze drying method followed by air drying (Table 3). Minimum titratable acidity (1.38) was reported with oven drying method while minimum total sugar (22.79), reducing sugar (16.04) and non-reducing sugar (6.74) content was reported under sun drying method. Oven drying and air drying methods were at par with each other. Reduced sugar content and acidity results from browning of sugars during sun drying and oven drying under the influence of heat. As opposed to freeze drying, which keeps a high sugar content while considerably reducing the likelihood of browning,

Comment [U6]: Units?

Table 3. Effect of drying methods on Titratable acidity, total sugar, reducing and non reducing sugars of guava fruit pulp powder

S.No.	Drying methods	Parameters			
		Titratable acidity%	Total Sugar%	Reducing Sugar%	Non-Reducing Sugar%
1.	Sun drying	1.64	22.79	16.04	6.74
2.	Oven drying	1.38	26.80	18.87	7.93
3.	Air drying	1.92	30.17	21.23	8.93
4.	Freeze drying	2.20	32.88	23.14	9.73
	SEd±	0.08	1.13	0.93	0.41
	C.D.(P=.05)	0.18	2.62	2.15	0.96

3.2.4 Ascorbic acid, total phenols and DPPH radical scavenging activity

Data analysis revealed highly significant results, the highest recovery of ascorbic acid was reported with freeze drying (227.90) followed by air drying (204.11) and oven drying (143.15) respectively, while minimum was reported with sun drying (122.97). The maximum total phenol content was reported with freeze drying (250.66) followed by air drying (207.9), while minimum was reported under oven drying(162.42) method. Highest inhibition of DPPH radical was observed under freeze drying method(83.40), succeeding air drying method (69.57)(Table4).Ascorbic acid and phenols are less likely to be retained during sun drying because of increased oxidation and breakdown, whereas oven drying uses high heat and also results in decreased retention of these chemicals. Ascorbic acid is a delicate substance that greatly decreases after heat treatment, as well as when exposed to sunlight and other environmental changes. The issues of oxidation, high temperatures, and environmental effects are practically all eliminated by freeze drying and air drying, while low temperatures enable improved ascorbic acid and phenol retention.(12-14,16-18).According to studies [19–21], the presence of phenols, ascorbic acid, and other bioactive substances has a direct correlation with antioxidant activity. More antioxidant activity may be due to the greater ascorbic acid and phenolic content discovered during freeze drying and air drying (DPPH Radical Scavenging Activity

Table 4.Effect of drying methods on Ascorbic acid, total phenols and DPPH radical scavenging activity of guava fruit pulp powder

S.No.	Drying methods	Parameters		
		Ascorbic acid(mg/100g)	Total phenols(mg/100g GAE)	DPPH radical scavenging activity%
1.	Sun drying	122.97	188.92	54.86
2.	Oven drying	143.15	162.42	48.04
3.	Air drying	204.11	207.9	69.57
4.	Freeze drying	227.90	250.66	83.40
	SEd±	6.72	1.62	0.61
	C.D.(P=.05)	15.51	3.74	1.42

3.2.5 Sensory analysis of guava's fruit pulp powder

Organoleptic characters significantly affected by different drying methods, and they are significantly differed from each other except oven drying and air drying which were at par with other. Score for taste (7.38), flavour (7.96), aroma (8.25) and overall acceptability 7.93() were maximum under freeze drying followed by air drying (Table 5). Minimum organoleptic score was found with sun drying method. Due to the extremely low risk of the fruit pulp's originality degrading, freeze drying and air drying kept greater taste, aroma, flavour, and overall acceptability [13,16].

Table 5. Effect of drying methods on sensory aspects of guava's fruit pulp powder

S.No.	Drying methods	Parameters			
		Taste	Flavour	Aroma	Overall acceptability
1.	Sun drying	6.43	6.75	7.09	6.56
2.	Oven drying	6.66	7.06	7.35	7.02
3.	Air drying	6.96	7.46	7.86	7.55
4.	Freeze drying	7.38	7.96	8.25	7.93
	SEd±	0.10	0.07	0.07	0.15
	C.D.(P=.05)	0.25	0.16	0.18	0.36

4. CONCLUSION

Although it was quite hygroscopic in nature, freeze drying generated the finest quality guava powder in terms of ascorbic acid, polyphenolic content, antioxidant activity, and flavour retention. It may be concluded that guava powder is an excellent source of natural antioxidants and can be used in place of synthetic antioxidants. According to the study, freeze-dried guava powder can be used as a component in the creation of food products with added value because it has a high total phenolic content together with other nutrients. In comparison to conventional drying technologies, this technology has the capability to dry delicate material while maintaining the highest level of nutritional stability. The outcomes for guava powder provided grounds for more research into its potential as a functional food.

REFERENCES

1. Alkandari D, Sarfraz H and Sidhu JS(2019). Development of a functional food (pan bread) using amla fruit powder. *J. Food Sci. Technol.*, 56(4):2287–2295.
2. Alothman M, Bhat R, Karim AA (2009) Antioxidant capacity and phenolic content of selected tropical fruits from Malaysia, extracted with different solvents. *Food Chem* 115:785–788.
3. Anonymous. <http://nhb.gov.in/area-pro/Indian>, Horticulture, 2013-14.
4. AOAC (2005) official methods of analysis, 18th edn. Association of official Analytical Chemists, Washington DC.
5. Goula AM, Adamopoulos KG, Kazakis NA (2004) Influence of spray drying conditions on tomato powder properties. *Dry Technol* 22(5):1129–1151.
6. Goyal, R.K., Patil, R.T., Kingsly, A.R.P., Walia, H. and Kumar, P. (2008). Status of post-harvest technology of aonla in India-A Review. *American Journal of Food Technology*, 3(1): 13-23.
7. Guiné RPF. (2018). The Drying of Foods and Its Effect on the Physical-Chemical, Sensorial and Nutritional Properties. *ETP Int. J. Food Eng.*, 4(2):93–100.
8. Karam MC, Petit J, Zimmer D, Baudelaire Djantou E, ScherJJ.(2016). Effects of drying and grinding in production of fruit and vegetable powders: A review. *J. Food Eng.*, 188:32–49.
9. Kaur C, Kapoor HC(2002). Anti-oxidant activity and total phenolic content of some Asian vegetables. *Int. J. Food Sci. Technol.*,37(2):153–161.
10. Kritkar, K.R. and Basu, B.D. (1993). *Phylanthusemblica*, In: Indian medicinal plants, 2nd ed. Lalit Mohan Basu (ed.), Calcutta. pp: 2220.
11. Lowry OH, Rosebrough NJ, Farr AL and Randall RJ.(1951). Protein measurement with the folin phenol reagent. *J. Biol. Chem.* 193(1):265-275.
12. Mishra P, Srivastava V, Verma D, Chauhan OP and Rai GK.(2009). Physico-chemical properties of Chakiya variety of Amla (*Emblicaofficinalis*) and effect of different dehydration methods on quality of powder. *African J. Food Sci.*, 3(10): 303–306.

13. Mitra SK, Bose T.(1990). Guava. In: Fruits Tropical and Subtropical., Eds., Naya Prakash, Calcutta,,: 278-303.
14. Murthy ZVP, Joshi D (2007) Fluidized bed drying of Aonla (*Emblicaofficinalis*). *Dry Technol* 25(5):883–889.
15. Nirmala Prabhu Reshmi, Goutam Mandal and Rocky Thokchom. (2018). Effect of Pre-Treatment of Salt Solution and Drying Methods on the Quality of Processed Aonla (*Emblicaofficinalis*Gaertn.). *Int.J.Curr.Microbiol.App.Sci.* 7(08): 1438-1446.
16. Pragati, Dahiya S and Dhawan SS. (2003).Effect of drying methods on nutritional composition of dehydrated aonla fruit (*Emblicaofficinalis* Garten) during storage. *Plant Foods Hum. Nutr.*, 58(3):1– 9.
17. Radojčin M, Pavkov I, BursaćKovačević D, Putnik P, Wiktor A, Stamenković, Z (2021). Effect of selected drying methods and emerging drying intensification technologies on the quality of dried fruit: A review. *Processes.*,9(1):132.
18. Ranganna S (2000) Handbook of analysis and quality control for fruit and vegetable products, 2nd edn. Tata McGraw-Hill Publ Co. Ltd,New Delhi.
19. Scartezzini P, Antognoni F, Raggi MA, Poli F and SabbioniC.(2006). Vitamin C content and antioxidant activity of the fruit and of the Ayurvedic preparation of *Emblicaofficinalis*Gaertn. *J. Ethnopharmacol.* ,104(1-2):113–118.
20. Shuen GW, Yi LY, Ying TS, Von GCY, Yusof YAB and PhingPL.(2021). Effects of drying methods on the physicochemical properties and antioxidant capacity of Kuini powder. *Brazilian J. Food Technol.*,24:1–14.
21. Singleton VL, OrthoferRandLamuela-Raventos RM.(1999). Analysis of total phenols an other oxidation substrates and anti-oxidants by means of folin-cocalteu reagent. *Scientia Horticulturae.*213(1974):152-178.
22. Somogyi LP, Barrett DM and Hui YH.(1996). *Major Processed Product.* 2 US: Technomic Publishing Co. Inc.
23. Sultana Bushra(2012). Effect of drying techniques on the total phenolic contents and antioxidant activity of selected fruits. *J. Med. Plants Res.* ,6(1):161–167.
24. Verma M, Singh J, Kaur D, Mishra V, Rai GK.(2015). Effect of various dehydration methods and storage on physicochemical properties of guava powder. *J. Food Sci. Technol.*,52(1):528–534.

UNDER PEER REVIEW

