

Original Research Article

Effect of Zn and B on Lentil (*Lens culinaris*) growth characteristics, yield, and available nutrients in the soil.

Abstract

To assess the effects of Zn and B treatment, a field experiment was carried out during the rabi season 2020–21 at the Crop Research Center of the Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut (U.P.). Ten treatments, each with a different combination of control, RDF, Zn, and B, were examined using a randomised block design with three replications. The experimental results revealed that growth attributing traits *viz.* Plant population (ha^{-1}), Plant height (cm), Number of branches plant^{-1} , Dry matter accumulation (g m^{-2}), Effective nodules (No. plant^{-1}), Nodules dry weight (mg plant^{-1}), yield *viz.* grain yield, straw yield, biological yield and harvest index and Available nutrient in soil N, P, K, Zn, S and B in lentil differed significantly among different treatments. Growth parameters and yield were significantly better in the treatment T₁₀ (RDF + Boron 2 kg ha^{-1} + Zinc 5 kg ha^{-1}). The highest grain yield was recorded in T₁₀ RDF + Boron 2 kg ha^{-1} + Zinc 5 kg ha^{-1} was applied with Zn and B and was statistically at par with T₈. From the study it may be concluded that the application of RDF + Boron 2 kg ha^{-1} + Zinc 5 kg ha^{-1} with Zn and B (T₁₀) gave best results (Grain yield increased by 26.7%, 25.7%, 21%, 22.9%, 17.2% and 59.1% over T₁, T₃, T₄, T₅, T₆ and T₇, respectively) and proved to be beneficial for *rabi* lentil followed by RDF + Boron 2 kg ha^{-1} + Zinc 5 kg ha^{-1} (T₉) also gave better results. Zn and B, together with N, P, K, and S, were used in lentils in a balanced manner to preserve soil fertility while also improving growth characteristics and yield.

Keywords: Micronutrient, Boron, Zinc, Sulphur, Lentil

1. INTRODUCTION

A nutrient-rich food legume, lentil (*Lens culinaris* Medic.), is a member of the *Fabaceae* family and is frequently referred to as an old crop in modern times. It is a grain legume with a high nutrient content that is grown in temperate climates. Its seed is a good source of fibre (4 g), ash (2.1 g), calcium (68 mg), phosphorus (325 mg), iron (7.0 mg), sodium (29 mg), potassium (780 mg), thiamine (0.46 mg), riboflavin (0.33 mg), and niacin (1.3 mg). It also has a relatively higher content of dietary protein (340–346 g). Among all the winter season legumes, lentil is the most abundant supply of vital amino acids (lysine, arginine, leucine, and other S-containing amino acids) (Sharma *et al.*, 2014). Pulses are particularly significant for food security in low-income nations since they provide 5% of the daily energy intake and 10% of the daily protein intake, respectively (Neacsu *et al.*, 2017). Pulses produced 87.40 million tonnes of grain with an average productivity of 1023 kg ha^{-1} on 85.40 million hectares of worldwide cropland in 2017. Still, though, the output of pulses is not keeping up with the daily minimum protein requirement of 60 g (Anonymous 2018). Around 6.58 million ha of land were planted with lentils globally in 2017, producing 7.59 million tonnes with an average yield of 1153 kg ha^{-1} (FAO, 2019). The area under pulses was >29 million ha with a total production of 25.23 million tonnes at a productivity of 841 kg ha^{-1} during 2017–18. India is the world's largest producer, consumer, and grower of pulses, accounting for 34%

of total acreage, 26% of total production, and approximately 30% (23–24 million tonnes) of the total consumption worldwide (Anonymous, 2018). After Canada, India is the second-largest producer of lentils. India, Canada, Turkey, Bangladesh, Iran, China, Nepal, and Syria are among the biggest lentil-producing nations in the world (Ahlawat, 2012). Lentil output in India reached an all-time high of 1.61 million tonnes from an area of 1.55 million ha at the all-time high yield level of 1034 kg ha⁻¹ (Anonymous, 2018). According to Singh *et al.* (2014), lentils are the third-most significant pulse crop in North India and are mostly farmed as a rainfed crop in the states of Uttar Pradesh, Uttarakhand, Madhya Pradesh, Jharkhand, Bihar, and West Bengal. Lentil, which ranks second among all legumes in terms of protein content per calorie behind soybean, is a key component of the diets of developing nations (Mudryj *et al.*, 2014). By influencing the plant itself and the symbiotic nitrogen fixation process, micronutrients have a significant impact on the yield of pulses. Due to their deficiency in soil, which is primarily caused by the removal of micronutrients from long-term crop production, increased use of only high analysis fertilizers, higher micronutrient requirements associated with higher crop yields, decreased use of animal manures in crop production, and use of high P concentrations from long-term applications, there is a great need for micronutrients today. About 200 enzymes and transcription factors require zinc, a crucial trace metal (Kabata, 2004). Zn is essential for the metabolism of auxin, nitrogen, and other nutrients as well as for the production of cytochrome C, stabilising ribosomal fractions, protecting cells from oxidative stress, and influencing the activities of many enzymes (Obata *et al.*, 1999). Field crops with a zinc shortage develop interveinal chlorosis and lower leaf necrosis with poor growth. Low-Zn seeds may have produced plants that are extremely vulnerable to biotic and abiotic stress (Mishra *et al.*, 2018). All plant meristems need a constant supply of boron from the soil since it is non-mobile in plants and necessary for the synthesis of cell walls, lignification, and structural integrity of bio membranes, stabilizing the ratio of sugars to starches, pollination, and seed generation (Kumar *et al.*, 2018). The production of pods and seeds, as well as cell division, depends greatly on boron (Singh *et al.*, 2015). Through seed treatment, foliar sprays, and soil fertilization, these micronutrients can be supplied to crops. Each technique has the potential to have an impact on the micronutrient nutrition of plants, both directly in the treated plant and in the offspring plants through nutrient enrichment of the parent plant's seeds.

2. MATERIALS AND METHODS

In the Indo-Gangetic plains of Western Uttar Pradesh, during the *rabi* season of 2020–21, a field experiment was carried out at Sardar Vallabh bhai Patel University of Agriculture and Technology. At an altitude of 230 metres above mean sea level, 29° 5' 34" N latitude, 77° 41' 58" E longitudes. While the mean weekly low temperature varied from 4.9 °C to 16.63 °C, the weekly maximum temperature ranged from 18.70 °C to 32.99 °C. Between 94.86 and 32.86%, the mean relative humidity ranged. Whereas, 39.8 mm of rain fell overall during the crop period. Before the experiment began, a composite soil sample (0–15 cm depth) was taken from the test site and examined for its physico-chemical composition. The soil in the test field was a sandy clay loam with a mildly alkaline response. The soil had a low amount of nitrogen readily available but a medium amount of available phosphorus and potassium.

2.1 Variety and nutrient

The experiment used the (Pusa Vaibhav) variety, which was released from IARI New Delhi in 1996 and is suitable for NWPZ (Punjab, Haryana, Delhi, and West UP), with a seed rate of @50 kg ha⁻¹ and a furrow depth of approximately 5 cm that was manually opened at a row

distance of 30 cm using a furrow opener. It has a production potential of 20–24 q ha⁻¹, is small-seeded, rust- and wilt-tolerant, and typically matures in 130–135 days. In this experiment, the recommended doses of N, P, K, and S were administered using DAP, MOP, urea, and bentonite, respectively, at 20, 50, 20, and 40 kg ha⁻¹. According to the treatments, zinc and boron were applied by zinc sulphate monohydrate and borax as a basal dose.

2.2 Growth attribute

2.2.1 Plant population

Plant populations per meter row length were recorded at harvest stages in all plots and then averaged.

2.2.2 Plant height (cm)

The height of five randomly selected plants were measured from the ground level to the base of apical bud with the help of a meter scale at harvest and mean height was computed.

2.2.3 Number of branches per plant

The number of basal branches arising from main shoot were counted in all five randomly marked plants at harvest and then mean were determined for each stage.

2.2.4 Number of nodules and dry weight of nodules

Five plants from each plot were uprooted thoroughly and kept in a tub filled with water. The roots of the plants was be washed in the tub and then nodules per plant was counted. All the nodules from the roots were detached and kept for drying in hot air oven at 60⁰C. At that the dry weight per plant were recorded.

2.2.5 Dry matter accumulation (g m⁻²)

The plant samples were cut close to the ground at harvest. The plant samples was sun dried and then dried in oven at 70⁰C till the constant weight obtained. Therefore, final weight was recorded.

2.3 Yield Study

Biological yield, seed yield and straw yield obtained from each plot was added to obtain biological yield in kilogram from each plot and converted to quintal per hectare. For seed yield the weight of clean seeds obtained from each plot was recorded on double pan balance. Finally, the seed yield plot⁻¹ was converted into yield ha⁻¹ by multiplying with appropriate value. Stover yield was determined by subtracting the seed yield from the biological yield of each net plot under a particular treatment. Then, the value was converted into Stover yields ha⁻¹ by using the appropriate value for each plot, which was used in case of conversion of seed yields ha⁻¹. Harvest Index (%) refers to the ratio of economic yield (seed yield) to the biological (Seed + Stover) yield under a particular treatment and expressed in percentage. It was be computed by using the following formula (Donald (1962)).

$$\text{Harvest index} = \frac{\text{Economic yield (kg ha}^{-1}\text{)}}{\text{Biological yield (kg ha}^{-1}\text{)}} \times 100$$

2.4 Soil analysis

2.4.1 Available nitrogen:

Available nitrogen in soil was determined by the procedure outlined by (Subbiahand Asija, 1956).

2.4.2 Available phosphorus:

Available P was extracted from the soil with 0.5 M NaHCO₃ solutions, pH 8.5 (Olsen *et al.*, 1954). Phosphorus in the extract was determined by developing blue colour with reduction of phospho-molybdate complex and the colour intensity was measured calorimetrically at 660 nm wave lengths.

2.4.3 Available potassium:

Exchangeable K content of soil was determined on 1 N NH₄OAC (pH 7.0) extract of the soil by using flame photometer. (Hanway and Heidel 1952).

2.4.4 Available Zinc:

Available Zn in soil was estimated by using DTPA as an extractant (Lindsay and Norvell, 1978) and its concentration were read on Atomic Absorption Spectrophotometer (GBC-Avanta PM Model).

2.4.5 Available Sulphur:

Available sulphur is present in mineral soil in the form of adsorbed SO₄-ion. To replace this adsorbed SO₄⁻ ions calcium chloride was used. Then SO₄ content of the extract was determined by turbidimetric method as proposed by (Williams and Steinbergs, 1959).

2.4.5 Available Boron:

Available B in soil was estimated by extraction with water directly on a hot plate. Use of azomethine-H in place of carmine or curcumin has further simplified the determination of hot-water soluble B (Gupta, 1967).

3 RESULTS AND DISCUSSION

3.1 Growth Attribute

3.1.1 Plant population

Plant population of lentil as influenced by different nutrient management at maturity Table 1 with the application of (T₁₀) RDF + Boron 2 kg ha⁻¹ + Zinc 5 kg ha⁻¹ was significantly higher than the rest of treatment, while minimum plant population (280014 ha⁻¹) was recorded under (T₁ control). Higher plant stand may be due to better management of micronutrients. Similar type of result was obtained by (Dhuppar *et al.*, 2012).

3.1.2 Plant Height

At harvest of lentil, the plant height (Table 1 and Fig. 1) was ranged from 38.5 to 47.9 cm. The maximum plant height (47.9 cm) found in T₁₀ was significantly superior to control and

statistically at par with rest of treatments, and increased by 24.4 % over control (T_1), while minimum (38.5 cm) was observed in control (T_1). The basal application of chemical fertilizers meets the nutritional requirement of crop for proper establishment and growth during the initial period. The increased plant height might be due to the involvement of nutrients in cell wall development and cell differentiation which resulted in elongation of shoot and root in plants. Similar results were obtained by (Muhammad and Khattak, 2009) who had reported that an appropriate supply of nutrients through inorganic sources increased the plant height of lentil through active photosynthesis. The results are in agreement with the findings of (Biswash *et al.*, 2014) and (Singh and Singh, 2014).

3.1.1.3 Number of Branches

At harvest branches decreased slightly and ranged from 2.8 to 4.1 plant⁻¹ (Table 1). The highest number of branches plant⁻¹ (4.1) recorded in T_{10} RDF + Boron 2 kg ha⁻¹ + Zinc 5 kg ha⁻¹ T_8 has same value which was statistically at par to T_8 while significantly higher than rest of treatments and lowest were 2.8 plant⁻¹ T_1 in (control). The increase in branches in (T_{10}) per plant over (T_1) control was 46.4%. The increased number of branches could be attributed to nutritional participation in cell wall formation and cell differentiation, which resulted in plant shoot and root elongation. The findings are consistent with those of (Singh *et al.*, 2005) and (Biswash *et al.*, 2014).

3.1.4 Dry Matter Accumulation

At harvest dry matter accumulation ranged from 177.1 to 269.7 (g m²) (Table 1 and Fig. 1). The highest dry matter accumulation recorded in T_{10} RDF + Boron 2 kg ha⁻¹ + Zinc 5 kg ha⁻¹ which was significantly higher to (T_1) control while statistically at par to rest of treatments and lowest were in T_1 (control). The dry matter accumulation in (T_{10}) RDF + Boron 2 kg ha⁻¹ + Zinc 5 kg ha⁻¹ (g m²) over (T_1) control was 52.3%. Similar kind of trend was observed by (Kumari and Ushakumari, 2002) who reported that the application of @ 30 kg P₂O₅ in cow pea significantly improved the dry matter. (Jat and Ahlawat, 2006) also reported the application of RDF + nitrogen + @ 26 P₂O₅ kg ha⁻¹ and S @ 40 kg ha⁻¹ in lentil, significantly increased dry matter accumulation in lentil. The results reported by (Meena, 2013) are also in close conformities with these findings.

3.1.5 Effective nodules (No. plant⁻¹) and Nodules dry weight (mg plant⁻¹)

The number of effective nodules plant⁻¹ and nodules dry weight in lentil was significantly influence by Zn and B application on effective nodules at 45 DAS (Table 1 and Fig. 1). The maximum number of effective nodules plant⁻¹ was recorded under T_{10} (37.7) which was statistically at par to treatment T_8 RDF + Boron 2 kg ha⁻¹ + Zinc 2.5 kg ha⁻¹ and significantly higher than the rest of treatment. While minimum (25.3 plant⁻¹) in control (T_1). Number of effective nodules plant⁻¹ increased by 49% in T_{10} RDF + Boron 2 kg ha⁻¹ + Zinc 5 kg ha⁻¹ over control (T_1) 45 DAS. The nodules dry weight of lentil was did not differ significantly under the influence of different treatments. The maximum nodules dry weight (67.1 mg plant⁻¹) observed with the application of (T_{10}) RDF + Boron 2 kg ha⁻¹ + Zinc 5 kg ha⁻¹ and minimum (55.3 mg plant⁻¹) in control. Effective nodule and nodule dry weight may be increased due to customized management of micronutrients. Similar type of results was also found by (Mohammad *et al.*, 2016) and (Singh, 2017).

3.2 Yield

The data pertaining to yield presented in Table 2 and Fig. 2 *ie.* Grain, straw, biological yield and harvest index. Grain yield of lentil under different treatments ranged from 10.5 to 16.7 q ha⁻¹. Maximum grain yield (16.7 q ha⁻¹) was noticed in T_{10} statistically at par to T_9 (RDF +

Boron 1 kg ha⁻¹ + Zinc 5 kg ha⁻¹), T₈ (RDF + Boron 2 kg ha⁻¹ + Zinc 2.5 kg ha⁻¹) and significantly higher than remaining treatments was found in T₁₀ (RDF + Boron 2 kg ha⁻¹ + Zinc 5 kg ha⁻¹). Minimum grain yield (10.5 q ha⁻¹) was observed under T₁. Significantly higher yield was obtained with Zn and B application of RDF + Boron 1 kg ha⁻¹ + Zinc 5 kg ha⁻¹ and RDF + Boron 2 kg ha⁻¹ + Zinc 5 kg ha⁻¹. Grain yield in T₉ and T₁₀ was higher by 46.66% and 59.0%, respectively over control. Result revealed that the grain yield increase by Zn and B application. Straw yield varied from 11.6 to 16.8 q ha⁻¹ under different treatments. Maximum straw yield (16.8 q ha⁻¹) statistically at par to T₉, T₈, T₇ and T₆ and significantly higher than remaining treatments was found in T₁₀ (RDF + Boron 2 kg ha⁻¹ + Zinc 5 kg ha⁻¹), while minimum (11.6 q ha⁻¹) in control (T₁). In comparison to T₁ (Control) straw yield increased by 44.8% in T₁₀. Biological yields ranged from 22.0 to 33.4 q ha⁻¹ under different treatments. Maximum biological yield (33.4 q ha⁻¹) register with T₁₀ which was statistically at par to the treatments T₉ (RDF+ Boron 1 kg ha⁻¹ + Zinc 5 kg ha⁻¹), T₈ (RDF + Boron 2 kg ha⁻¹ + Zinc 2.5 kg ha⁻¹), T₇ (RDF + Boron 1 kg ha⁻¹ + Zinc 2.5 kg ha⁻¹) and T₆ (RDF + Zinc 5 kg ha⁻¹) and significantly higher than remaining treatments was found in T₁₀ (RDF + Boron 2 kg ha⁻¹ + Zinc 5 kg ha⁻¹), while minimum significantly lower than the rest of treatments in control (T₁). Harvest index express proportion of economic yield in total biological yield did not differ significantly by the Zn and B application during the experimentation. Numerically maximum harvest index value (49.9%) was observed in T₁₀ (RDF + Boron 2 kg ha⁻¹ + Zinc 5 kg ha⁻¹) than rest of the treatments during year of study. Lowest harvest index (47.5%) was recorded in control (T₁). The proper mobilization of dry matter production towards the sink (seed yield) is an important factor for economic yield. The variation in seed yield of different lentil cultivars has been reported by various researchers like (Zike *et al.*, 2017), (Iliger and Alagundagi, 2017) and (Biswas *et al.*, 2014).

3.3 Available nutrients in soil

Available nutrients in soil as present in Table 3 and Fig. 3 Shows that available nitrogen revealed that numerically value of available nitrogen was observed with in application of RDF + Boron 2 kg ha⁻¹ + Zinc 5 kg ha⁻¹ (T₁₀) over remaining treatments. After crop harvest available nitrogen in soil ranged from 175.2 to 210.5 kg ha⁻¹. Maximum available nitrogen (210.5 kg ha⁻¹) was observed in T₁₀ (RDF + Boron 2 kg ha⁻¹ + Zinc 5 kg ha⁻¹), while minimum (210.5 kg ha⁻¹) in control (T₁). For available phosphorus the data presented revealed that numerical value of available phosphorus was observed with in application of RDF + Boron 2 kg ha⁻¹ + Zinc 5 kg ha⁻¹ (T₁₀) over remaining treatments. It is clear from the data available soil phosphorus at harvest differed significantly under the influence of different treatments. After crop harvested available P in soil ranged from 11.5 to 14.9 kg ha⁻¹ under different treatments. Maximum available phosphorus (14.9 kg ha⁻¹) statistically at par in treatments T₉ (RDF + Boron 1 kg ha⁻¹ + Zinc 5 kg ha⁻¹), T₈ (RDF + Boron 2 kg ha⁻¹ + Zinc 2.5 kg ha⁻¹), T₇ (RDF + Boron 1 kg ha⁻¹ + Zinc 2.5 kg ha⁻¹) and T₆ (RDF + Zinc 5 kg ha⁻¹) and significantly higher than the rest treatments was found in RDF + Boron 2 kg ha⁻¹ + Zinc 5 kg ha⁻¹ (T₁₀), while minimum (11.5 kg ha⁻¹) in control (T₁). For available potassium the data presented revealed that numerical value of available potassium was observed with in application of RDF + Boron 2 kg ha⁻¹ + Zinc 5 kg ha⁻¹ (T₁₀) over remaining treatments. It is clear from the data available soil potassium at harvest differed significantly under the influence of different treatments. After crop harvested available K in soil ranged from 180.7 to 210.8 kg ha⁻¹ under different treatments. Maximum available potassium (210.8 kg ha⁻¹) was did not differed significantly by the application of Zn and B during experimentation. The

available potassium minimum (180.7 kg ha^{-1}) in control (T_1). For available zinc the data presented revealed that numerically value of available zinc was observed with in application of RDF + Boron 2 kg ha^{-1} + Zinc 5 kg ha^{-1} (T_{10}) over remaining treatments. It is clear from the data available soil zinc at harvest differed significantly under the influence of different treatments. After crop harvested available Zn in soil ranged from 0.77 to 0.80 g ha^{-1} under different treatments. Maximum available zinc (0.80 g ha^{-1}) was did not differed significantly by the application of Zn and B during experimentation. The available zinc minimum (0.77 g ha^{-1}) in control (T_1). For available Sulphur the data presented revealed that numerically value of available sulphur was observed with in application of RDF + Boron 2 kg ha^{-1} + Zinc 5 kg ha^{-1} (T_{10}) over remaining treatments. It is clear from the data available soil sulphur at harvest differed significantly under the influence of different treatments. After crop harvested available S in soil ranged from 7.6 to 10.6 kg ha^{-1} under different treatments. Maximum available sulphur (10.6 kg ha^{-1}) statistically at par in treatments T_9 (RDF + Boron 1 kg ha^{-1} + Zinc 5 kg ha^{-1}), T_8 (RDF Boron 2 kg ha^{-1} + Zinc 2.5 kg ha^{-1}) and T_7 (RDF + Boron 1 kg ha^{-1} + Zinc 2.5 kg ha^{-1}) and significantly higher than the rest treatments was found in RDF + Boron 2 kg ha^{-1} + Zinc 5 kg ha^{-1} (T_{10}), while minimum (7.6 kg ha^{-1}) in control (T_1). For available boron the data presented revealed that numerically value of available boron was observed with in application of RDF + Boron 2 kg ha^{-1} + Zinc 5 kg ha^{-1} (T_{10}) over remaining treatments. It is clear from the data available soil boron at harvest differs significantly under the influence of different treatments. After crop harvested available B in soil ranged from 28.2 to 32.4 g ha^{-1} under different treatments. Maximum available boron (32.4 g ha^{-1}) statistically at par in treatments T_9 (RDF + Boron 1 kg ha^{-1} + Zinc 5 kg ha^{-1}), T_8 (RDF + Boron 2 kg ha^{-1} + Zinc 2.5 kg ha^{-1}) and T_7 (RDF + Boron 1 kg ha^{-1} + Zinc 2.5 kg ha^{-1}) and significantly higher than the rest treatments was found in RDF + Boron 2 g ha^{-1} + Zinc 5 kg ha^{-1} (T_{10}), while minimum (28.2 g ha^{-1}) in control (T_1). Increased availability of N, P, K, S, Zn and B in the soil might be due to balanced combined application of Zn and B. The combination of Zn and B customized fertilizer also facilitated more availability of N, P, K, S, Zn and B in soil and also more uptakes by crop. Results of similar kind have also been reported by (Nasser *et.al.*, 2008). This might be due to its increased availability with the supplementation of these nutrients in soil. Increase in Zn and B content in soil by its application in soil has also been reported by (Jat *et.al.*, 2006) and (Arya *et al.*, 2007).

Table: 1 Effect of Zn and B on Lentil Plant population (ha⁻¹), Plant height (cm), Number of branches plant⁻¹, Dry matter accumulation (g m⁻²), Effective nodules (No. plant⁻¹), Nodules dry weight (mg plant⁻¹) at harvest

Symbol	Treatments	Plant population (ha ⁻¹)	Plant height (cm)	Number of branches plant ⁻¹	Dry matter accumulation (g m ⁻²)	Effective nodules (No. plant ⁻¹)	Nodules dry weight (mg plant ⁻¹)
T ₁	Control	280014	38.5	2.8	177.1	25.3	55.3
T ₂	RDF (N:P:K:S)	286518	44.1	3.3	244.0	29.8	60.2
T ₃	RDF + Boron 1 kg ha ⁻¹	287290	45.8	3.4	245.6	31.3	61.4
T ₄	RDF + Boron 2 kg ha ⁻¹	288340	46.3	3.4	254.6	31.5	62.8
T ₅	RDF + Zinc 2.5 kg ha ⁻¹	287801	46.2	3.4	250.7	31.0	61.8
T ₆	RDF + Zinc 5 kg ha ⁻¹	289380	46.2	3.5	250.7	32.0	63.5
T ₇	RDF + Boron 1 kg ha ⁻¹ + Zinc 2.5 kg ha ⁻¹	291104	46.4	3.5	267.6	32.5	63.9
T ₈	RDF + Boron 2 kg ha ⁻¹ + Zinc 2.5 kg ha ⁻¹	295589	47.4	4.1	269.7	34.2	65.9
T ₉	RDF + Boron 1 kg ha ⁻¹ + Zinc 5 kg ha ⁻¹	292475	46.6	3.5	269.7	33.6	64.1
T ₁₀	RDF + Boron 2 kg ha ⁻¹ + Zinc 5 kg ha ⁻¹	296590	47.9	4.1	269.7	37.7	67.1
	SEm (±)	1948	1.6	0.11	9.08	1.17	2.24
	C.D. (P=0.05)	5476	4.8	0.35	27.18	3.51	NS

Table. 2 Effect of Zn and B on yield on Lentil

Symbol	Treatments	Yields (q/ha)			Harvest Index (%)
		Grain	Straw	Biological	
T ₁	Control	10.5	11.6	22.0	47.5
T ₂	RDF (N:P:K:S)	13.9	14.9	28.7	48.2
T ₃	RDF + Boron 1 kg ha ⁻¹	14.0	14.9	28.9	48.4
T ₄	RDF + Boron 2 kg ha ⁻¹	14.5	15.2	29.7	48.8
T ₅	RDF + Zinc 2.5 kg ha ⁻¹	14.3	15.1	29.4	48.5
T ₆	RDF + Zinc 5 kg ha ⁻¹	14.9	15.5	30.4	49.0
T ₇	RDF + Boron 1 kg ha ⁻¹ + Zinc 2.5 kg ha ⁻¹	15.1	15.8	31.0	49.1
T ₈	RDF + Boron 2 kg ha ⁻¹ + Zinc 2.5 kg ha ⁻¹	15.4	15.8	31.2	49.5
T ₉	RDF + Boron 1 kg ha ⁻¹ + Zinc 5 kg ha ⁻¹	15.4	15.9	31.4	49.2
T ₁₀	RDF + Boron 2 kg ha ⁻¹ + Zinc 5 kg ha ⁻¹	16.7	16.8	33.4	49.8
	SEm (±)	0.5	0.5	1.0	1.7
	C.D. (P=0.05)	1.5	1.6	3.1	NS

Table: 3 Effect of Zn and B on available Nitrogen, Phosphorus, Zinc, Sulphur, Boron in Lentil at harvest

Symbol	Treatments	Nitrogen (kg ha ⁻¹)	Phosphorus (kg ha ⁻¹)	Potassium (kg ha ⁻¹)	Zinc (g ha ⁻¹)	Sulphur (kg ha ⁻¹)	Boron (g ha ⁻¹)
T ₁	Control	175.2	11.5	180.7	0.77	7.6	28.2
T ₂	RDF (N:P:K:S)	192.5	12.6	195.6	0.77	9.3	29.4
T ₃	RDF + Boron 1 kg ha ⁻¹	193.5	12.8	190.8	0.73	9.3	32.7
T ₄	RDF + Boron 2 kg ha ⁻¹	195.4	12.9	192.4	0.77	9.2	34.8
T ₅	RDF + Zinc 2.5 kg ha ⁻¹	195.5	13.2	198.2	0.80	9.4	29.8
T ₆	RDF + Zinc 5 kg ha ⁻¹	198.2	13.6	200.1	0.83	9.5	28.5
T ₇	RDF + Boron 1 kg ha ⁻¹ + Zinc 2.5 kg ha ⁻¹	206.8	13.7	204.7	0.77	10.2	31.4
T ₈	RDF + Boron 2 kg ha ⁻¹ + Zinc 2.5 kg ha ⁻¹	209.5	13.6	207.4	0.77	10.5	33.2
T ₉	RDF + Boron 1 kg ha ⁻¹ + Zinc 5 kg ha ⁻¹	208.5	14.1	208.7	0.83	10.4	31.4
T ₁₀	RDF + Boron 2 kg ha ⁻¹ + Zinc 5 kg ha ⁻¹	210.5	14.9	210.8	0.80	10.6	32.4
	SEm (±)	7.12	0.48	7.10	0.037	0.34	1.14
	C.D. (P=0.05)	NS	1.43	NS	NS	1.02	3.42




Fig.1 Effect of Zn and B on, Plant height (cm), Dry matter accumulation (g m⁻²), Effective nodules (No. plant⁻¹), Nodules dry weight (mg plant⁻¹) at harvest

UNDER REVIEW




Fig. 2 Effect of Zn and B on yield on Lentil

UNDER




Fig. 3 Effect of Zn and B on available Nitrogen, Phosphorus, Zinc, Sulphur, Boron in Lentil at harvest

4. Conclusion

Based on the results of this study, it can be stated that farmers can use the (T₁₀)-RDF (N:P: K:S @ 20, 50, 20, and 40 kg ha⁻¹) + Boron 2 kg ha⁻¹ + Zinc 5 kg ha⁻¹ combination, followed by either the (T₈) -RDF + Boron 2 kg ha⁻¹ + Zinc 2.5 kg ha⁻¹ or the (T₉)- RDF + Boron 1 kg ha⁻¹ + Zinc 5 kg ha⁻¹. The observations are based on data from one season, so it is advised that the experiment be repeated in the future on the same soil with the same layout to obtain more precise information.

5. REFERENCES

Ahlawat IP S. Agronomy – rabi crops, Lentil. Division of Agronomy Indian Agricultural Research Institute, New Delhi – 110 012 *Agronomy* ; 2012.

Anonymous, 2018. Economic Survey of India. annual report.

Arya, R.L., Varshney J.G. and Kumar Lalit (2007). Effect of integrated nutrient application in chickpea+mustard intercropping system in the semi-arid tropics of North India. *Comm. Soil Science and Plant Analysis* 38 (1/2): 229-240.

Biswash, M.R., Rahman, M.W., Haque, M.M., Sharmin, M. and Barua, R. (2014). Effect of Potassium and Vermicompost on the Growth, Yield and Nutrient Contents of Mungbean (BARI Mung 5). *Open Science Journal of Bioscience and Bioengineering*. 1(3): 33-39.

Biswash, M.R., Rahman, M.W., Haque, M.M., Sharmin, M. and Barua, R. (2014). Effect of Potassium and Vermicompost on the Growth, Yield and Nutrient Contents of Mungbean (BARI Mung 5). *Open Science Journal of Bioscience and Bioengineering*. 1(3): 33-39.

Dhuppar, P., Biyan, S., Chintapalli, B., & Rao, S. (2012). Lentil crop production in the context of climate change: an appraisal. *Indian Research Journal of Extension Education*, 2 (Special Issue), 33-35.

Donald CM. In search of yield. *Journal of the Australian Institute of Agricultural Sciences* 1962;28:171-178.

FAO. FAOSTAT statistical database of the *United Nations Food and Agriculture Organization* (FAO); 2019.

Gupta, U. C.. A simplified method for determining hot-watersoluble boron in podzol soils. *Soil science*, 1967;103(6), 424-428.

Hanway JJ. Heidel H. Soil analysis methods as used in Iowa state college soil testing laboratory. *Iowa Agriculture*. 1952;57:1-31

Iliger, M. D., Alagundagi, S. C., Patil, M. B., & Vijayakumar, A. G. (2017). Influence of seed rate and fertilizer levels on growth and yield of lentil (*Lens culinaris* Medik.) genotypes under dry land situation. *Journal of Pharmacognosy and Phytochemistry*, 6(6), 2019-2022.

Jat, R.S. and Ahlawat, I.P.S. 2006. Direct and residual effect of vermicompost sulphur fertilization on soil nutrient dynamics and productivity of mungbean, maize sequence. *Journal of sustainable Agriculture*. 28(1): 41-54.

Kabata-Pendias, A. (2004). Soil–plant transfer of trace elements—an environmental issue. *Geoderma*, 122(2-4): 143-149.

- Kumar, J., Gupta, S., Gupta, D. S., & Singh, N. P. (2018). Identification of QTLs for agronomic traits using association mapping in lentil. *Euphytica*, 214(4), 1-15.
- Lindsay WL, Norvell WA. Development of a DTPA soil test for zinc, iron, manganese, and copper. *Soil Science society of America Journal*.1978;42(3):421-428.
- Mishra, U. S., Sharma, D., & Raghubanshi, B. P. S. (2018). Effect of zinc and boron on yield, nutrient content and quality of blackgram (*Vigna mungo* L.). *Research on Crops*, 19(1), 34-37.
- Mohammed, Y. A., Chen, C., McPhee, K., Miller, P., McVay, K., Eckhoff, J., ... & Holmes, J. (2016). Yield performance and stability of dry pea and lentil genotypes in semi-arid cereal dominated cropping systems. *Field Crops Research*, 188, 31-40.
- Mudryj, A. N., Yu, N., and Aukema, H. M. (2014). Nutritional and health benefits of pulses. *Applied Physiology, Nutrition, and Metabolism*, 39(11): 1197-1204.
- Muhammad, D. and Khattak, R.A. (2009). Growth and nutrients concentrations of maize in pressmud treated saline sodic soils. *Soil Environment*, 28(2):145-155.
- Nasser, R. R., Fuller, M. P., & Jellings, A. J. (2008). Effect of elevated CO₂ and nitrogen levels on lentil growth and nodulation. *Agronomy for Sustainable Development*, 28(2), 175-180.
- Neacsu, M., McBey, D., & Johnstone, A. M. (2017). Meat reduction and plant-based food: replacement of meat: nutritional, health, and social aspects. *Sustainable protein sources*, 359-375.
- Obata, H., S. Kawamura, K. Senoo and A. Tanaka (1999). Changes in the level of protein and activity of Cu/Zn superoxide dismutase in zinc deficient rice plant, *Oryza sativa* L. *Soil Science Plant Nutrition* 45: 891-896.
- Olsen SR. Estimation of available phosphorus in soils by extraction with sodium bicarbonate (No.939). *US Department of Agriculture*; 1954.
- Sharma, A.K., Raghubanshi, B.P.S. and Sirothia, P. (2014). Response of chickpea to levels of zinc and phosphorus. *Annals of Plant and Soil Research*, 16(2): 172- 173.
- Singh, D, Khare, A., Sharma, R., & Chauhan, S. S. (2015). Effect of boron on yield quality and uptake of nutrients by lentil. *Annals of Plant and Soil Research*, 17(4), 385-387.
- Singh, D. and Singh, H. (2012). Effect of phosphorus and zinc nutrition on yield, nutrient uptake and quality of chickpea. *Annals of Plant and Soil Research*, 14(1): 71- 74.
- Singh, D. and Singh, R.P. (2014). Effect of integrated nutrient management on growth, physiological parameters and productivity of lentil (*Lens culinaris* Medik.). *International Journal of Agricultural Sciences*, 10(1): 175-178.

Singh, K.K., Srinivasarao, C. and Ali, M. (2005). Root growth, nodulation, grain yield, and phosphorus use efficiency of lentil as influenced by phosphorus, irrigation, and inoculation. *Communications in Soil Science and Plant Analysis*, 36(13/14): 1919-1929.

Subbiah B, Asija GL. Alkaline permanganate method of available nitrogen determination. *Current Science*. 1956;25:259-260.

Williams, C. H., & Steinbergs, A.. Soil sulphur fractions as chemical indices of available sulphur in some Australian soils. *Australian Journal of Agricultural Research*, 1959; 10(3), 340-352.

Zike, T., Abera, T. and Hamza, I. (2017). Response of improved lentil (*Lens culinaris* Medik) varieties to phosphorus nutrition on vertisols of West Showa, Central Highlands of Ethiopia. *Advances in Crop Science and Technology*, 5(6): 315.

UNDER PEER REVIEW