

**Effect of Spacing and Panchagavya on growth and yield attributes of
Chickpea(*Cicer arietinum* L.).**

ABSTRACT

A field experiment was conducted during *Rabi* season 2021-22 at the experimental field of the Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology And Sciences, Prayagraj, Uttar Pradesh, India. The Experiment was laid out in Randomized Block Design with Nine treatments replicated thrice. The treatments consisted of three levels of spacing 20cm x 10cm, 30cm x 10cm, 40cm x 10cm and three levels of panchagavya – 2%, 3%, 5%. The results revealed that significantly higher plant height (51.1 cm), maximum number of nodules/plant (11.7) and higher plant dry weight (25.7 g). Whereas the maximum number of pods / plant (62.0), higher seeds / pod (1.86), maximum seed index (22.9 g), higher seed yield (2.58 t/ha), higher stover yield (5.34 t/ha) and higher harvest index (36.6%) were recorded in the treatment-5 (30cm x 10 cm + Panchagavya 3%).

Keywords: *Chickpea*, Spacing, Panchagavya, Growth and Yield.

INTRODUCTION

Pulses are valuable source of dietary protein and have a specific ability to sustain and preserve soil fertility by adding biological nitrogen fixation. Pulses are restorative crops and leaves quite enough as 30 kg/ha nitrogen in soil (**Reddy and Reddy, 2010**). Among the pulses, chickpea is an important *rabi* season crop with high acceptability and wider use in nutritional food basket. Chickpea contains 18-22 per cent protein, 52-70 per cent carbohydrate, 4-10 percent fat and sufficient quantity of minerals, calcium, phosphorus, iron and vitamins. In India pulses are grown nearly in 28.83 m ha with an annual production of 25.72 m t and productivity of 0.8 tha. Some of the states like Uttarpradesh is about 8.24 m ha with an annual production of 9.97 m t and productivity of 1.08 t ha major producer of chickpea in India as advocated by Ministry of agriculture and Farmer Welfare (**GOI 2020-21**). Besides, it is also important for sustainable agriculture as it improves the physio-chemical and biological properties of the soil. Its deep roots also open the soil, which ensure better aeration and heavy leaf drop increases the organic matter in the soil. It can fix about 25-30 kg N/ ha through symbiosis and these minimize dependency on chemical fertilizers. Thus, chickpea plays a vital role in improving the soil health (**DES 2018**).

Row spacing is one of the important characters which can be manipulated to attain the maximum production from per unit land area. The optimum row spacing with proper geometry of planting is dependent on variety, its growth habit and agro climatic condition. The seed yield of chickpea is highly dependent on plant population. Seed yield increases with decrease in row spacing up to an optimum limit which changes according to genotypes. Row spacing is also one of the important factor which ultimately effect nutrient uptake growth and yield of plant. Increase in spacing decreases the total population, but with more nutrition to the individual plants grows better and yield more and vice versa. The increases or decreases of plant population has definite pattern in relation to the yield (**Nawange et al., 2015**).

Panchagavya is an organic product that can be used as manure. This is prepared by using five main ingredients cow dung, cow urine, cow milk, Desi cow ghee, and curd. Panchagavya is used for organic farming as it plays a major role in the growth of plants and also in increasing the immunity of plants. Panchagavya, an organic product has the potential to play the role of promoting growth and providing immunity in plant system. Panchagavya consists of five products viz. cow dung, cow urine, milk, curd, ghee. When suitably mixed and used, these have miraculous effects.

Plants sprayed with Panchagavya invariably produce bigger leaves and develop denser canopy. The photosynthetic system is activated for enhanced biological efficiency, enabling synthesis of maximum metabolites and photosynthates. The key feature of Panchagavya is its efficacy to restore the yield level of all crops when the land is converted from inorganic cultural system to organic culture from the very first year. The harvest is advanced by 15 days in all the crops. It not only enhances the shelf life of vegetables, fruits and grains, but also improves the taste (**Selvaraj *et al.*, 2007**).

MATERIALS AND METHODS

The experiment was carried out during *rabi* season of 2021-22 at Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, SHUATS, Prayagraj (U.P.). which is located at 25° 30' 42''N latitude, 81° 60' 56'' E longitude, and a height of 98 metres above sea level. The soil texture in the experimental plot was sandy loam, with a practically neutral soil (PH 7.1), low in organic carbon (0.44%) available N (171.48 kg/ha), available P (27.0 kg/ha) and K (291.2 kg/ha). The crop was sown on 13 december 2021 using variety NBeG-49. The experiment was set up in a Randomized Block Design with three replications and nine treatments in total *Viz.*, T₁: 20cm x 10 cm + Panchagavya 2%, T₂: 30cm x 10 cm + Panchagavya 2%, T₃: 40cm x 10 cm + Panchagavya 2%, T₄: P20cm x 10 cm + Panchagavya 3%, T₅: 30cm x 10 cm + Panchagavya 3%, T₆: 40cm x 10 cm + Panchagavya 3%, T₇: 20cm x 10 cm + Panchagavya 5%, T₈: 30cm x 10 cm + Panchagavya 5% and T₉: 40cm x 10 cm + Panchagavya 5%. The growth Parameters and yield attributes were recorded at harvest form randomly selected plants in each plot. The data was computed and analysed by following statistical method of **Gomez and Gomez (1984)**.

RESULTS AND DISSCUSSION

Effect of Spacing and Panchagavya on growth attributes of chickpea.

Plant height (cm)

At harvest the significant and higher plant height (51.1 cm) was recorded in treatment 5 (Spacing 30cm x 10 cm + Panchagavya 3%). However, treatment 6 (40cm x 10 cm + Panchagavya 3%) and treatment 8 (30cm x 10 cm + Panchagavya 5%) was found statistically at par with treatment 5 (Spacing 30cm x 10 cm + Panchagavya 3%). The significantly higher plant height was under treatment of spacing 30 x 10 cm this might be due to plant get sufficient space under wider spacing for light, air and nutrition for better growth and development. Similar

findings also reported by **Chaudhary *et al.*, (2015)**.

Number of nodules/plant

Significantly maximum number of nodules / plant (11.7) was recorded in treatment 5 (Spacing 30cm x 10 cm + Panchagavya 3%). However, treatment 6 (40cm x 10 cm + Panchagavya 3%) and treatment 8 (30cm x 10 cm + Panchagavya 5%) were found statistically at par with treatment 5 (Spacing 30cm x 10 cm + Panchagavya 3%). Significant and maximum number of root nodules / plant might be due to the better availability of nutrients from panchagavya which has many nutrients and effective conversion of nutrients from organics such as Fe, Mg and Zn available at the site of photosynthesis. application of panchagavya 3 per cent noticed higher number of nodules and effective root nodules which resulted in manifestation of higher growth and yield attributing characters and higher yield of chickpea . Similar findings also reported by **Yadav *et al.*, (2017)**.

Plant Dry weight (g)

Significantly maximum plant dry weight (25.7 g) was recorded in treatment 5 (Spacing 30cm x 10 cm + Panchagavya 3%). However, treatment 6 (40cm x 10 cm + Panchagavya 3%) and treatment 8 (30cm x 10 cm + Panchagavya 5%) were found statistically at par with treatment 5 (Spacing 30cm x 10 cm + Panchagavya 3%).. Significant increase in plant dry weight might be due to foliar spray of panchagavya made improvement in dry matter accumulation, chlorophyll content and nitrogen content described above may be ascribed to higher yield and yield attributes with Panchagavya. Similar findings also reported by **Kumawat *et al.*, (2009) and Saranraj *et al.*,(2010)**.

Effect of Spacing and Panchagavya on yield and yield attributes of chickpea.

Number of pods/plant

Treatment-5 (30cm x 10 cm + Panchagavya 3%) was recorded significantly maximum number of pods / plant (62.0) which was superior over all other treatments. However, treatment 6 (40cm x 10 cm + Panchagavya 3%) and treatment 8 (30cm x 10 cm + Panchagavya 5%) was found statistically at par with treatment 5 (30cm x 10 cm + Panchagavya 3%). Significant increase number of pods/plant this might be due panchagavya which is an efficient plant growth stimulant that enhanced the biological efficiency of crops. Similar findings also reported by **Kulkarni *et al.*, (2015)**.

Number of seeds/ pod

Treatment-5 (30cm x 10 cm + Panchagavya 3%) was recorded significantly maximum number of seeds per pod (1.86) which was superior over all other treatments However, treatment 6 (40cm x 10 cm +

Panchagavya 3%) and treatment 8 (30cm x 10 cm + Panchagavya 5%) were found statistically at par with treatment 5 (30cm x 10 cm + Panchagavya 3%). Significant increase in number of seeds/pod Probably may be due to favorable effect of panchagavya on reproductive growth viz., pods / plant which is one of the important yield attributes having significant positive correlation with seed & straw yield. Similar findings also reported by **Choudhary *et al.*, (2017)**.

Seed index (g)

Treatment-5 (30cm x 10 cm + Panchagavya 3%) was recorded significantly maximum seed index (22.9g) which was superior over all other treatments. However, treatment 6 (40cm x 10 cm + Panchagavya 3%) and treatment 8 (30cm x 10 cm + Panchagavya 5%) were found statistically at par with treatment 5 (30cm x 10 cm + Panchagavya 3%). The higher seed index might be due to favorable effect of the cow dung present in the panchagavya acts as a medium for the growth of beneficial microbes and cow urine provides nitrogen which is essential for crop growth upon fermentation with other ingredients in panchagavya has beneficial effect on growth and yield **Patil *et al.*, (2012)**.

Seed yield (t/ha)

Treatment-5 (30cm x 10 cm + Panchagavya 3%) was recorded significantly maximum Seed yield (2.58 t/ha) which was superior over all other treatments. However, treatment 6 (40cm x 10 cm + Panchagavya 3%) was found statistically at par with treatment 5 (30cm x 10 cm + Panchagavya 3%). The significant and maximum seed yield was recorded with 30 cm of row spacing which was significantly more over others row spacing followed in experiment. It might be due to sufficient plant population which caused for higher growth and development along with proper utilization of production inputs and ultimate results was maximum yield. Similar findings also reported by **Verma *et al.*, (2008)**.

Stover yield (t/ha)

Treatment-5 (30cm x 10 cm + Panchagavya 3%) was recorded significantly maximum Stover yield (5.34 t/ha) which was superior over all other treatments. However, treatment 6 (40cm x 10 cm + Panchagavya 3%) was found statistically at par with treatment 5 (30cm x 10 cm + Panchagavya 3%). The maximum stover yield might be due to the IAA and GA present in panchagavya when applied as foliar spray could have created stimuli in the plant system and increased the production of growth regulators in cell system and the action of growth regulators in plant system ultimately stimulated the necessary growth and development This might be attributed to the favourable effect on vegetative growth, i.e. plant height and reproductive growth parameters. The similar findings were also reported by **Panchal *et al.* (2017), Chakraborty and**

Sarkar (2019).**Harvest Index (%)**

Treatment-5 (30cm x 10 cm + Panchagavya 3%) was recorded significantly maximum harvest index (36.6%) which was superior over all other treatments. The maximum harvest index suggests that individual plant under higher plant population performed better than plant under lower plant population the improvement in yield attributing parameters and yield per plant of individual plant under lower plant population was not sufficient enough to compensate the loss in density for higher seed yield. The similar findings also reported by **Nawange *et al.*, (2015)**.

Table 1. Effect of Spacing and Panchagavya on growth and yield of Chickpea.

S. No.	Treatments	Plant Height (cm)	Number of Nodules/plant	Plant Dry weight (g)
1	20cm x 10 cm + Panchagavya 2%	45.4	6.5	19.2
2	30cm x 10 cm + Panchagavya 2%	46.0	8.2	24.0
3	40cm x 10 cm + Panchagavya 2%	44.6	7.3	23.0
4	20cm x 10 cm + Panchagavya 3%	44.0	7.8	20.7
5	30cm x 10 cm + Panchagavya 3%	51.1	11.7	25.7
6	40cm x 10 cm + Panchagavya 3%	49.6	11.1	25.1
7	20cm x 10 cm + Panchagavya 5%	43.5	9.0	20.1
8	30cm x 10 cm + Panchagavya 5%	49.0	10.1	24.8
9	40cm x 10 cm + Panchagavya 5%	45.6	9.2	22.3
	F test	S	S	S
	SEm (\pm)	0.76	0.56	0.55
	CD (P=0.05)	2.28	1.68	1.66

Table 2. Effect of Spacing and Panchagavya on yield attributes of chickpea.

S. No.	Treatments	No. of Pods/plant	No. of Seeds/pod	Seed Index (g)	Seed yield (t/ha)	Stover yield (t/ha)	Harvest index (%)
1	20cm x 10 cm + Panchagavya 2%	54.3	1.13	19.1	2.00	3.88	35.1
2	30cm x 10 cm + Panchagavya 2%	59.0	1.46	21.3	2.46	4.48	34.0
3	40cm x 10 cm + Panchagavya 2%	58.7	1.46	20.9	2.31	4.20	34.5
4	20cm x 10 cm + Panchagavya 3%	57.8	1.20	20.0	2.20	4.14	36.0
5	30cm x 10 cm + Panchagavya 3%	62.0	1.86	22.9	2.58	5.34	36.6
6	40cm x 10 cm + Panchagavya 3%	61.0	1.79	22.3	2.52	5.23	36.3
7	20cm x 10 cm + Panchagavya 5%	54.5	1.26	19.6	2.05	4.05	34.7
8	30cm x 10 cm + Panchagavya 5%	60.9	1.73	22.2	2.43	4.84	35.5
9	40cm x 10 cm + Panchagavya 5%	58.8	1.55	21.4	2.30	4.34	35.8
	F test	S	S	S	S	S	S
	SEm (\pm)	0.30	0.09	0.35	0.03	0.16	0.32
	CD (P=0.05)	2.51	0.29	1.07	0.11	0.50	1.07

CONCLUSION

Based on the research done in one season, it is concluded that for obtaining higher yield in chickpea, the treatment (T5) with Spacing 30cm x 10cm and Panchagavya 3% is advised as it is recorded higher in growth and yield.

REFERENCES

- Chakraborty, Bishal and Sarkar, Indrajit. (2019). Quality Analysis and Characterization of Panchagavya, Jeevumrutha and Sasyamrutha. *Int.J.Curr.Microbiol.App.Sci.* **8(5)**: 2018-2026. doi: <https://doi.org/10.20546/ijemas.2019.805.234>.
- Chaudhary,N.A, Vihol,J.K, Chaudhary,H.J, Mor,B.V and Desai,J.L. (2015). Influence of spacing and scheduling of irrigation on growth, yield, yield attributes and economics of summer greengram (*Vigna radiata* L.). *Eco. Env. & Cons.* **21** (December Suppl.) : 2015; pp. S357-S361.
- DES, Ministry of Agriculture & FW, (2018). Success report of 2018-2019, pulses Revolution – From Food to Nutritional Security. Available online at <https://farmers.gov.in/SucessReport2018-19> and retrieved on July 17, 2020.
- Fisher, R.A. and Yates, F. (1963). Statistical tables, Oliver and Boyd, Edinburgh, Tweed date Court, London (U.K).
- GOI, 2020-21. Economics and Statistics of Indian Agriculture. Annual report, Government of India, New Delhi.
- Gomez, K.A. and Gomez, A.A., (1984). Statistical procedures for agricultural research (2 ed.). John Wiley and Sons, New York, 680p.
- Choudhary, Lal. Gopal, Sharma,S.K, Singh Pal Kendra, Choudhary Sanju and Bazaya, B.R. (2017). Effect of Panchagavya on Growth and Yield of Organic Blackgram [*Vigna mungo*(L.)Hepper]. *Int.J.Curr.Microbiol.App.Sci.* **6(10)**:1627-1632. doi: <https://doi.org/10.20546/ijemas.2017.610.1950> .
- Jackson, M.L. (1973). Soil Chemical Analysis (II Edition). Prentice Hall of India Private Limited. New Delhi, India.
- Kulkarni, Shyamrao, Upperi, S.N. and Jadhav, R.L.. (2016). Greengram productivity enhancement through foliar spray of nutrients. *Legume Research - An International Journal.* **39**. 814-816. 10.18805/lr.v0iOF.9386.
- Kumawat R.N, Mahajan S.S and Mertia R.S. (2009). Growth and development of Groundnut (*Arachis hypogaea*) under foliar application of Panchagavya and leaf extracts of endemic plants. *Indian Journal of Agronomy* **54(3)**: 324-331.
- Nawange,D.D, Verma,H.D. and Verma Hemlata.(2015).Relative performance of Kabuli chickpea (*Cicer arietinum* L.) genotypes under different plant densities and fertility levels.. *Agric. Sci. Digest.*, **36** (2) 2016:160-162.

row spacing on growth and yield of chickpea (*Cicer arietinum* L.). *Agricultural Science Digest*. (2008). **28** (2):139 – 140.

Panchal Pratik Patel, Pravinchandra , Patel .G,A, and Desai, Ajit. (2017). Effect of Panchagavya on growth, yield and economics of chickpea (*Cicer arietinum*). *International Journal of Chemical Studies* 2017; **5**(2): 265-267.

Saranraj.P (2011). Growth and development of blackgram (*Vigna mungo*) under foliar application of Panchagavya as organic source of nutrient. *Current Botany*.

Selvaraj,N.,Anitha.B.,Anusha.B and Guru Saraswathi.M. (2007). Organic Horticulture. Horticultural Research Station, Tamil Nadu Agricultural University, Udhagamandalam-643 001.

Yadav, J. K., Sharma, M., Yadav, R.N., Yadav S.K. and Yadav.S (2017) Effect of different organic manures on growth and yield of chickpea (*Cicer arietinum* L.). *Journal of Pharmacognosy and Phytochemistry*; **6**(5): 1857-1860.