

Minireview Article

Reducing the Carbon footprint by cultivating and consuming Spirulina- A mini review

Abstract:

Increased carbon emissions have increased ~~the~~ global warming resulting in tremendous changes in the climate factor. Climate change has brought drought, cyclones, floods, tsunami, irregular rainfall patterns threatening food security, clean water, etc. To reduce the impacts of global warming that are detrimental to humanity and ~~the~~ biosphere, global nations have agreed to reduce carbon emissions and go for ~~a~~ low carbon footprint in all industrial and commercial activities. This study analyzes ~~the~~ cultivation and consuming *spirulina* (*Arthrospira platensis*) as a win-win situation for sustainable development in terms of GDP as well as carbon capture & storage (CCS).

Keywords:

Sustainability, carbon capture, ~~blue~~ blue-green algae, ~~nutrient~~ nutrient-rich bioresource, phycocyanin.



Fig. 1. *Spirulina* culture



Fig. 2. Collection technique Fig. 3. Role of *Spirulina* in different sectors

1.Introduction:

Herzog et al.,(2001) Natural ways of carbon sequestration are by vegetation in soil, ~~in~~ forests, and in oceans as natural sinks. Artificial carbon capture can be done in Geological sinks that can hold thousands of gigatons of carbon in depleted oil and gas reservoirs, deep saline formations, and unminable coal seams. CO₂ injection into geological formations for enhanced oil recovery (EOR) is a prominent technology that reduces the viscosity of heavy oil resulting in the release of trapped oil. **Song et al.,(2016)** Phytoliths are silicified forms of plants that get accumulated in soils and sediments for hundreds to thousand years. Organic carbon present in phytoliths is mainly captured from atmospheric carbon dioxide during photosynthesis. Phytolith carbon sequestration in croplands, grasslands, forests, and bamboo lands can be an efficient measure for carbon capture. **Sayre et al.,(2010)** Microalgae are recognized as the most productive biological systems for biomass production than terrestrial systems due to their short maturation cycle. Carbon dioxide or bicarbonate capturing efficiencies in their cell structure makes them apt for carbon capture from stationary point sources like cement kilns or power stations and from nonpoint sources like atmospheric carbon. **Schipperet al.,(2013)** Constructing algal ponds near industries is a necessary step followed by industries to reduce the carbon footprint. **Vonshaket al.,(2004)** *Spirulina* is a blue-green algae with spiral filaments under the genus *Arthrospira*, and phylum Oscillatoriaceae. It is a cyanobacteria and contains two pigments green (chlorophyll) that does photosynthesis and fixes carbon dioxide and blue (phycocyanin) in its cell structure. It is a unicellular micro algae growing in high alkaline conditions 10-12 pH introduced to modern society by Kanembu tribes of Africa, near Lake Chad. In 1965, the botanist Jean Leonard confirmed that ~~it~~ he consumed by the Kanembu tribe is made up of *spirulina*, and he cultured it in a sodium

Formatted: Highlight

Formatted: Font: Not Italic

hydroxide production facility in laboratory conditions. From that time till date *spirulina* is cultured and consumed by trillions of people as it is a GRAS level edible algae. This study involves the particular species *Arthrospira platensis* biomass useful to humanity in infinite ways.

2. Nutrient profile of *Spirulina*:

Capelli et al. (2010), Grosshagauer et al. (2020), *Spirulina* is rich in exopolysaccharides (15-21%), fatty acids, and easily digestible protein (60-70%). It contains a wide array of micronutrients and vitamins. GLA- gamma linolenic acid present in mother's milk is an important constituent that makes *spirulina* nature's mother's milk to humans. As it is rich in chlorophyll, a pigment resembling the structure of hemoglobin in the blood it can fight anemia. It is a rich vegetative source of vitamin B12, cyanocobalamin and is rich in other vitamins also. The micronutrients present are calcium, potassium, chromium, copper, iron, magnesium, manganese, phosphorus, selenium, sodium, zinc, molybdenum, chloride, germanium, and boron. Phycocyanin the blue polypeptide present in *spirulina* regulates the production of white blood cells and stem cells thus imparting immunoregulatory nature to the microalgae. Phycocyanin also helps in the formation of blood-hematopoiesis. This versatile nutrient profile of *spirulina* has its approval approved by WHO as an efficient tool against malnutrition and anemia in children and women. Beta carotene an anti-anti-oxidant is present that improves eye health and other antioxidants rich property helps to fight free radicals reducing cancer risk and heart attacks. **Yüce tepe et al. (2016)** In this study compares conventional protein sources like egg and soybean with the nutritional benefits of bioactive peptides present in *spirulina* based upon the reports of FAO. All these qualities have given *spirulina* the nutraceutical value making it the **Great king of microalgal kingdom.**

Formatted: Highlight

Amino acid	Per 10 gm	% of total	Amino acid	Per 10 gm	% of total
Isoleucine	350 mg	5.6	Cystine	60 mg	1.0
Leucine	540 mg	8.7	Arginine	430 mg	6.9
Lycine	290 mg	4.7	Histidine	100 mg	1.6
Phenylalanine	280 mg	4.5	Threonine	320 mg	5.2
Tyrosine	300 mg	4.8	Proline	270 mg	4.3
Methionine	140 mg	2.3	Valine	400 mg	6.5
Glutamic acid	910 mg	14.6	Alanine	470 mg	7.6
Aspartic acid	610 mg	9.8	Glycine	320 mg	5.2
Tryptophan	90 mg	1.5	Serine	320 mg	5.2

Table 1 : “Aminoacids present in *Spirulina*”

3. *Spirulina* in Aquaculture:

Calcium-Spirulan is a unique polymerized sugar that does not allow a virus to penetrate the cell membrane to infect the cell and other polysaccharides of *spirulina* are involved in DNA repair mechanisms and enzymes synthesis in cells (Fig. 1). Thus supplementing *spirulina* as a feed, prevents the animal from viral attack increasing its immune system. **Vonshak et al. (2004)** By using *spirulina* as fish feed reduces the major production cost which is the feed cost (40-60%) in aquaculture. It also improves the health of

the cultured species, makes the fish species more fit for human consumption by reducing antibiotics and hormones use in the culture, increases disease resistance, and the survival rate of the larvae. **Cardoso et al.,(2021)**The *Spirulina* sp. LEB 18 was cultivated in aquaculture wastewater supplemented with 25% zarrouks medium to obtain biomass of T-25 assay with the highest concentrations of protein (65.73%), phycocyanin (16.60 mg/mL), PUFA (38.20%), and γ -linolenic acid GLA (23.29%). It also showed 90% bioremediation potential with the removal of COD, sulphate, phosphate, and bromine and proved to be ideal for biodiesel applications.

4.Spirulina as animal feed:

Holman et al.,(2013) *Spirulina* is a promising new feed resource supporting rations of agriculturally significant animals like cattle,cows,pigs,rabbits,poultry showing improvements in productivity, health and product quality.**Panjaitan et al.,(2010)**Approximately 20% of dietary *Spirulina* bypasses rumen degradation and is available for direct absorption within the abomasum and increases microbial crude protein production, and reduces its retention time within the rumen.**Jin et al.,(2020)**observed *spirulina* powder supplement in feeds for 125 days culture of abalones and found enhancement in the shell size and protein content of abalones.

5.Spirulina as a Biofertilizer:

Dineshkumar,et al.,(2018)This study used *chlorella vulgaris* and *spirulina platensis* for sustainable agriculture reducing the use of chemical fertilizers polluting the environment to get 7-20.9% increased yield in rice crops by nitrogen fixation by the microalgae.**Mahapatra et al.,(2018)**Cultivation of *Spirulina* sp. isolated from urban wastewater-fed lakes in outdoor rooftop batch cultures with concentrated wastewaters are a typical zero waste economy converting macronutrients C:N:P in the wastewater into algal biomass with 100% efficiency making them potential biofertilizers. **Jadhav,et al.,(2020)**studied the enhancement of soya bean plant growth in terms of plant height, number of branches, and number of leaves by applying BGA fertilizer and obtained positive results.**Dinesh kumar et al.,(2019)**applied *spirulina* and *chlorella* along with cowdung to the seeds of crops Maize, Onion, Green gram, Black gram, Tomato and Paddy and concludes that shifting to organic agriculture is possible.Thus sustainable agriculture is close to our hands by using BGA biofertilisers.**Wuang, et al.,(2016)**By applying *spirulina* fertilizer observed enhanced growth of Chinese Cabbage, Chinese broccoli, and Protea White Crown plants from seed germination.**Hamouda et al.,(2022)**Agricultural lands are degraded mostly due to extensive use of chemical fertilizers for the long term and also groundwater table going very low in certain areas causing salt water penetration.The study was conducted with *Triticum aestivum* L. against salinity concentrations of seawater 10% and 25%.With Aliquots of 2% liquid extracts from *A.platensis*, there was stimulated an enhancement in protein, antioxidants, carbohydrates, and total phenols of the plant in both concentrations of the sea water.Thus *spirulina* biofertiliser can be effective against salinity stress in crop growth. **Gonçalves et al.,(2021)**In this review discusses in detail about different mechanisms involved in applying *spirulina* biofertiliser for plant growth. Polysaccharides, long-chain fatty acids, and enzymes secreted by *A.platensis* are antagonistic to plant pathogens as a biocide.It can act as a biostimulant by secretion of growth-promoting phyto hormones like auxins. Various bioactive molecules and antioxidants secreted by the microalgae can act as stress tolerance boosters. **Bioremediation—The bioremediation** potential of *spirulina* improves soil quality by chelating toxic metal ions and the presence of surplus micronutrients and P and nitrogen-fixing capacity enhances plant growth. Improving *spirulina* biofertiliser using nanotechnology as algal biochar nanofertiliser is an upgrowing field that improves the phycoprosects of the fertilizer leading to sustainable agricultural practices (Fig. 2).

6.Spirulina in the cosmetics industry:

Formatted: Font: Italic

Ragusa, et al., (2021) in their detailed study of *spirulina* extracts in wide range of wound healing, antiaging, anti-anti-acne, skin care products concludes it as-is a booster in all products with no side effects. In beauty parlours, *spirulina* face packs and lip balms are used as anti-anti-wrinkle agents. Costa, et al., (2017) The biologically active metabolites in *spirulina* are interesting ingredients for nutraceutical formulations and are important for skin care and anti-age-aging.

7. Spirulina in the Pigments industry:

The pigments present in spirulina are Phycocyanin (Blue): 14%, Chlorophyll (Green): 1% , Carotenoids (Orange/ Red): 47%. (Ciferri, 1983) *Spirulina* contains two biliproteins with high economical value: c-phycocyanin and allophycocyanin with absorption maxima at 615-620nm and 650nm in the visible region respectively. The chromophore is phycobilin, an open tetrapyrrole. Bousiba and Richmond (1980) Lina blue is commercialized by Dainippon Ink & Chemicals of Japan which constitutes of spirulina Phycocyanin. It is used as a food colorant in the coloring of candy, ice cream, dairy products, and soft drinks as well as in cosmetics like eye shadow, eyeliner, and lipstick and in-textiles such as natural blue. Ranjitha et al., (2020) used phycocyanin pigments from *spirulina* as a natural photosensitizer for bio-sensitized solar cells (BSSC). Silver-Silver-doped TiO₂ nanoparticles were prepared by sol-gel technique and along with pigments of phycocyanin extracted from *A. platensis* was fabricated in a solar cells to obtain high efficiency making this combination ideal for future BSSC applications. Moldovan et al., (2022) Sustainability in the textile industry leads us to shift in the use of phycocyanin blue from *A. platensis* than to chemical and traditional dyes. The pre-mordanted cotton and bleached wool with phycocyanin-rich extract, representing the sustainable blue dye were tested for color characterization and fastness. The results validated the sustainable character of spirulina-based phycocyanin in the dyeing process yielding low oxygen demanding effluent waste according to the international standards thus less polluting the environment. Jeyaraja et al., (2022) The authors observed photocatalytic degradation of organic dye Malachite green of concentration 25ppm by phycocyanin extracts of *A. platensis* in sunlight. After 3 hours, 100% of the dye was degraded that which is confirmed by UV absorbance studies which showed no peak in 620nm proving dye degradation. Ho et al., (2019) The light-light-harvesting pigment phycocyanin of *spirulina* residue was converted into biochar by pyrolysis at 900 degrees and it is activated by peroxydisulfate to obtain an efficient long durable carbon green catalyst for wastewater treatment that showed marvelous bactericidal properties on *Escherichia coli*.

8. Spirulina vs Global warming – The Conclusion:

Global algal biodiesel market is approximately USD 6.95 billions in the year 2020 with an annual growth rate of 8%. Due to the richness of nutrients, *Spirulina* can be used as a prominent substrate for industrially important biomolecules production. The ease of production and its capacity to harvest sun's energy makes this microalgae a renewable source of energy. As a biofertiliser, it enhances the agricultural productivity, and its metal binding nature and presence of long-long-chain fatty acids, holds it a special place in water treatment methods (Fig. 3). Thus one species of this microalgae can achieve seven sustainable development goals - 7SDGs like sustainable agriculture yielding food security, clean water (used in WWT), clean energy production, upgrading degraded land (heavy metal removal and salt tolerance), Clean air (carbon sequestration & mitigation of climate change), good health (nutraceutical value), eradicating poverty by an improved economy. This wonder product can be produced in larger quantities by exploring novel methods like symbiosis and co-culturing techniques with other microorganisms like *Rhodotorula* yeast, *Pseudomonas stutzeri*, *Azospirillum brasilense*, and *Lactobacillus plantarum*, which enhances the yield manifolds so that it reaches every human hand at a low cost. Literature - The literature says 1 gram of spirulina biomass uses 0.4-2 grams of carbon dioxide. In other words, one acre of algae can remove 2.7 tonnes of carbon dioxide which is 10-50% more

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

efficient than terrestrial plants. If we follow the footprints of SPIRULINA in anyone of the above aspects, it will reduce the carbon footprint which is our prior duty to mother EARTH.

References:

- [1].Boussiba, Samy, and Amos E. Richmond. "C-phycoyanin as a storage protein in the blue-green alga *Spirulina platensis*." *Archives of Microbiology* 125.1 (1980): 143-147.
- [2].Capelli, Bob, and Gerald R. Cysewski. "Potential health benefits of *Spirulina* Microalgae." *Nutrafoods* 9.2 (2010): 19-26.
- [3].Cardoso, Lucas Guimarães, et al. "*Spirulina* sp. as a bioremediation agent for aquaculture wastewater: production of high added value compounds and estimation of theoretical biodiesel." *BioEnergy Research* 14.1 (2021): 254-264.
- [4].Ciferri, Orio. "*Spirulina*, the edible microorganism." *Microbiological reviews* 47.4 (1983): 551-578.
- [5].Costa, Jorge Alberto Vieira, et al. "The potential of spirulina and its bioactive metabolites as ingested agents for skin care." *Industrial Biotechnology* 13.5 (2017): 244-252.
- [6].Dineshkumar, R., et al. "Marine microalgal extracts on cultivable crops as a considerable bio-fertilizer: A Review." (2019).
- [7].Dineshkumar, R., et al. "Microalgae as bio-fertilizers for rice growth and seed yield productivity." *Waste and biomass valorization* 9.5 (2018): 793-800.
- [8].Gonçalves, Ana L. "The use of microalgae and cyanobacteria in the improvement of agricultural practices: a review on their biofertilising, biostimulating and biopesticide roles." *Applied Sciences* 11.2 (2021): 871.
- [9].Grosshagauer, Silke, Klaus Kraemer, and Veronika Somoza. "The true value of *Spirulina*." *Journal of agricultural and food chemistry* 68.14 (2020): 4109-4115.
- [10].Hamouda, Ragaa A., et al. "Protective role of *Spirulina platensis* liquid extract against salinity stress effects on *Triticum aestivum* L." *Green Processing and Synthesis* 11.1 (2022): 648-658.
- [11].Herzog, Howard. "What future for carbon capture and sequestration?." *Environmental Science and Technology-Columbus* 35.7 (2001): 148A.
- [12].Ho, Shih-Hsin, et al. "N-doped graphitic biochars from C-phycoyanin extracted *Spirulina* residue for catalytic persulfate activation toward nonradical disinfection and organic oxidation." *Water research* 159 (2019): 77-86.
- [13].[https://en.wikipedia.org/wiki/Spirulina_\(dietary_supplement\)](https://en.wikipedia.org/wiki/Spirulina_(dietary_supplement))
- [14].<https://thefishsite.com/articles/the-multifunctional-dietary-properties-of-spirulina-and-its-use-in-aquaculture>
- [15].Jadhav, S. R., and S. M. Talekar. "Growth of soyabean [*Glycine Max* L.(Merr)] under the influence of blue green algal (BGA) Biofertilizer." *BIOINFOLET-A Quarterly Journal of Life Sciences* 17.1a (2020): 87-89.
- [16].Jeyaraja, Sharmila, et al. "Phycocyanin from *Spirulina platensis* bio-mimics quantum dots photocatalytic activity: A novel approach for dye degradation." *Environmental Science and Pollution Research* (2022): 1-13.

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Field Code Changed

Formatted: Portuguese (Brazil)

- [17].Jin, Su-Eon, et al. "Spirulina powder as a feed supplement to enhance abalone growth." *Aquaculture Reports* 17 (2020): 100318.
- [18].Mahapatra, Durga Madhab, et al. "Algae-based biofertilizers: a biorefinery approach." *Microorganisms for green revolution*. Springer, Singapore, 2018. 177-196.
- [19].Moldovan, Simona, et al. "Wastewater effluents analysis from sustainable algae-based blue dyeing with phycocyanin." *Textile Research Journal* (2022): 00405175221119419.
- [20].Panjaitan, T., et al. "Effect of the concentration of Spirulina (*Spirulina platensis*) algae in the drinking water on water intake by cattle and the proportion of algae bypassing the rumen." *Animal Production Science* 50.6 (2010): 405-409.
- [21].Ragusa, Irene, et al. "Spirulina for skin care: A bright blue future." *Cosmetics* 8.1 (2021): 7.
- [22].Ranjitha, S., et al. "Synthesis and development of novel sensitizer from spirulina pigment with silver doped TiO₂ nano particles for bio-sensitized solar cells." *Biomass and Bioenergy* 141 (2020): 105733.
- [23].Sayre, Richard. "Microalgae: the potential for carbon capture." *Bioscience* 60.9 (2010): 722-727.
- [24].Schipper, Kira, et al. "New methodologies for the integration of power plants with algae ponds." *Energy procedia* 37 (2013): 6687-6695.
- [25].Song, Zhaoliang, Kim McGrouther, and Hailong Wang. "Occurrence, turnover and carbon sequestration potential of phytoliths in terrestrial ecosystems." *Earth-Science Reviews* 158 (2016): 19-30.
- [26].Vonshak, Avigad, and Giuseppe Torzillo. "Environmental stress physiology." *Handbook of microalgal culture: biotechnology and applied phycology* 57 (2004).
- [27].Wuang, S.C.; Khin, M.C.; Chua, P.Q.D.; Luo, Y.D. Use of Spirulina biomass produced from treatment of aquaculture wastewater as agricultural fertilizers. *Algal Res.* 2016, 15, 59–64.
- [28].Yücepete, Aysun, and BeraatÖzçelik. "Bioactive peptides isolated from microalgae *Spirulina platensis* and their biofunctional activities." *AkademikGıda* 14.4 (2016): 412-417.