

A Study on Bio-efficacy of Dinotefuran 20% SG against Brown plant hopper *Nilaparvatha lugens* (Stall) on paddy *Oryza sativa* L in Raichur district of Karnataka State

Comment [h1]: delete

Abstract

In rice cultivation, the Brown plant hopper causes damage to growth of plants by sucking cell sap and it causes in drying of plants and crop fields results into yellow color later it turns brown. In order to overcome this kind of damage particularly in rice cultivation, a field trial was conducted during kharif 2017 at Agricultural Research Station Gangavathi to assess the bio-efficacy of Dinotefuran 20% SG against brown plant hopper *Nilaparvatha lugens* (Stall) on paddy. The results revealed that, the new insecticide Dinotefuran 20% SG 40 g a.i./ha showed lowest population of 0.86 and 0.73, 0.67 and 0.57 BPH/hill at 3, 7,10 and 15 days after spray (DAS) respectively and this treatment was significantly superior over rest of the treatments. Further, it was on par with lower dose of Dinotefuran 20% SG @ 30 g a.i./ha which is recorded 1.63, 1.60, 1.49 and 1.42 BPH/hill at 3, 7, 10 days and 15 DAS respectively, followed by standard check buprofezin @ 187.5g a.i./ha recorded 1.63, 1.90, 2.05 and 2.15 and BPH/hill at 3, 7,10 and 15 DAS respectively. Another insecticide imidacloprid @ 22.5g a.i./ ha was found inferior to all other treatments by harbouring 4.43,4.53, 4.76 and 4.87 BPH/hill, where as the untreated control recorded highest BPH population of 21.70 to 25.64 BPH/hill through out the observational period from one day before to 15 DAS respectively. The maximum additional yield was 22.37 % was recorded over control (64.28 q/ha) was found in case of Treatment 4 (T4) with an incremental benefit cost ratio of 3.72.

Comment [h2]: Thereby resulting in drying and yellowing of plants and crop fields which later on turn brown.

Comment [h3]: manage

Comment [h4]: pest

Comment [h5]: italicize

Comment [h6]: observed at

Comment [h7]: remove is

Comment [h8]: remove resp

Comment [h9]: which

Comment [h10]: the bioefficacy of Imidaclopride...

Comment [h11]: of

Comment [h12]: in case of treatment (T4) with an.....

Key words: *Brown Plant hopper, Nilaparvatha lugens, bio-efficacy and Paddy*

Introduction:

The study of efficacy of Insecticide emerged in the year 1960's especially for species of plant and leaf hoppers, the chemicals such as organophosphate, cyclodiene and organochlorine carbamate were the main insecticides used for controlling these hoppers among chemical methods. The Brown Plant hopper is sucking pest and is known as significant pests in Asia especially in rice growing area (Rao et al., 2003). Rice is staple food crop due this reason there was long term evolution on efficacy of insecticides studies were discovered and still going on. The insect has capacity to develop fastest growth particularly in tropical areas, it can complete 12 generations within in single year and it is categorized as migratory pest (Cheng J, 2009). Conversely, nearly 100 species of insects including more than 20 economic pests are competent to damage to rice plants (Rahaman et al 2016, Pathak. M.D. and Dhaliwal. G.S.1981). Insect pests continue to pose threat to rice farming and they are major constraints to rice production and coexist with rice growth (Savary S et al 2000 and Nirmala B and Muthuraman P, 2016, Dhawan et al., 2011, Pasalu and Katti, 2006, and Alam and Das, 2017). Hence, under these dynamic situations of both

Comment [h13]: rewrite the introduction again start with importance of rice crop first then insect pest spectrum and management and the need of the study.

Comment [h14]: bio-

Comment [h15]: insecticides

Comment [h16]: organochlorines, cyclodiene, organophosphate and carbamates

Comment [h17]: one of the

Comment [h18]: Rice being a staple food crop the long term evolution on bioefficacy studies of insecticides is still going on due to the severe insect pest load particularly brown plant hoppers

Comment [h19]: Achieve fastest growth and development particularly....

Comment [h20]: a

Comment [h21]: remove

insect damage and climatic conditions, monitoring plant health will be always challenging task. This has to be addressed with due care in order to stabilize the both yield and plant health in future.

Rice *Oryza sativa* a cereal crop, belong to the family Graminae is one of the most important staple food crop in the world. for more than half of the world population. Rice constitutes 55 per cent of total cereal production and 52 per cent of the total food grain production in India (Saxena. R.C and Singh. R.K.2003). Rice is the major staple food in many developing countries. It is an important crop because it contains high nutritive value each 100 gm of rice is consists of energy 1,527 KJ (365 Kcal) and carbohydrate around 76.7gm as major nutrition. Other edible form of rice used as puffed rice, rice flakes, rice wafers and canned rice. The by products of rice starch is largely used in beverage industries (Salim et al. 2003 and FAO 2011). Brown plant hopper is major pests across the country especially in irrigated rice where intensive paddy cropping is being done. Three species of plant hoppers reported on rice are white backed plant hopper (WBPH), *Sogatella furcifera* (Horvath), Smaller brown plant hopper (SBPH), *Laodelphax striatellus* (Fallen) and brown plant hopper (BPH) *Nilaparvath lugens* (Stall). First and third of these are of economic importance. Brown plant hopper is the most destructive pest of rice in Uttar Pradesh, Madhya Pradesh, West Bengal, Andra Pradesh, Karnataka and Tamil Nadu (Chung et al.,1982, Liu et al., 2003).

The plant hopper suck the plant sap from the phloem vessels through their proboscis. Due to this plant start wilting with outer most leaves start drying first and then the entire plant dries up- a symptom often called “hopper burn” (Patcharin,2011). BPH and WBPH casuses huge crop loss in grain yield ranging from 10 - 70 per cent (Kulshreshta, 1974) and 35 – 95 per cent (Sindhu, 1970), respectively. Hence these two pests combination (BPH &WBPH) have been emerged as the number one pest which limit the rice production in India. Among the major pests brown plant hopper constitutes one of the most important causing substantial yield losses. Uses of chemical insecticides forms one of the effective management practices and an important Integrated Pest management (IPM) component besides cultural and Biological methods of pest control. Synthetic insecticides are proved to be the only option where we can rely for critical management of insect pests reaching on or beyond ETL level (Singh.S.P., 2000). Indiscriminate use of broad spectrum of chemical insecticides also reduce the biodiversity of natural ecosystem their by reduce the natural enemies population and induce out break of secondary pests and polluted the natural eco-system results in resurgence of brown plant hopper. But still chemical control forms the first line of defence.(I.C.Pasalu et al.,2002). So there is a need to evaluate the new groups, new formulations of insecticides and their combinations for their effect on target and non target insects (Whalon et al.,2008). There fore present investigation was carried out to evaluate new insecticide molecules against BPH infesting rice.

Comment [h22]: s

Comment [h23]: space

Materials and methods: Field trial conducted during *kharif* 2017 at Agricultural Research Station Gangavathi to assess the bio-efficacy of Dinotefuran 20% SG against brown plant hopper *Nilaparvatha lugens* (Stall) on paddy for the variety Sona Masuri (BPT-5204). There were totally 7 treatments among them Dinotefuran 20% SG at 4 different concentrations viz., @ 20g, 25g, 30g, and 40g were tested and these were compared with Imidacloprid 17.8 SL @ 22.5g, standard check Buprofezin 25% SC @187.5g and control. Further, to assess the grain yield and benefit-cost ratio of different treatments were also calculated. The per cent increase in yield was calculated by following formula and it was adopted by Matharu and Tanwar (2020). Economics of different treatments was analyzed, using the cost of insecticide, its application cost and other charges etc. during the field trial. The data on grain yield per hectare and its prevalent market price were used to work out the benefit derived from each treatment / ha. Based on Incremental benefit in yield over control and the cost involved, Incremental Cost Benefit Ratio (ICBR) was worked out to establish economic ranking of various treatments.

Comment [h24]: an experimental field trial was...

Comment [h25]: the grain yieldwas also calculated

Comment [h26]: given

Comment [h27]: on per hectare basis.

Increase of yield (%) = $\frac{\text{Yield in treatment} - \text{Yield in control}}{\text{Yield in Control}} \times 100$.

Fourty days' old seedlings of rice BPT 5204 were transplanted in the plots measuring 15.75sqm, arranged in Randomized Block Design, at a spacing of 20 X 10 cm. The recommended N:P:K @ 120:60:60 Kg/ha were applied to the crop. The entire amount of P₂O₅ and K₂O and one fourth of the N₂ were given as basal and rest of N₂ was given in two splits. Irrigation was provided as per requirement of the crop. Other crop production measures were taken as usual. The crop was inspected daily for observing the incidence of rice hoppers. The crop was infested by five species of leaf and plant hoppers (Table 2) of which only Brown plant hopper (BPH), *Nilaparvata lugens* was found at above ETL level. The population of other species was scanty. The insecticidal spraying was given with the help of an high volume knapsack sprayer using 500 litres of water. The spraying was done at about eighty one days after transplanting when the population of BPH reached the ETL (20 insects/hill) in most of the plots; the second spraying was given after 10 days of the first spray. Five randomly selected hills were selected in each treatment plots. Each treatment was replicated thrice in randomized complete block design (RCBD). Observations were taken at one day, three day and seven days, tendays and fifteen days after spray.

Comment [h28]: particularly BPH

Comment [h29]: delete

Comment [h30]: delete

Comment [h31]: per ha

Comment [h32]: after

Comment [h33]: after 1, 3, 7, 10 and 15th days

Treatment wise yield was also recorded and subjected to statistical analysis.

Table 1: Name of the insecticides tested along with their dosages and manufacturers.

Treatments. No.	Insecticide	Dosage (g a.i. / ha	Manufacturer
T1	Dinotefuran 20% SG	20 g	P. I. Industries Ltd
T2	Dinotefuran 20% SG	25 g	P. I. Industries Ltd
T3	Dinotefuran 20% SG	30 g	P. I. Industries Ltd
T4	Dinotefuran 20% SG	40g	P. I. Industries Ltd
T5	Imidacloprid 17.8% SL	22.5g	Bayer Crop Care India Ltd.
T6	Buprofezin 25% SC	187.5 g	Syngenta India Ltd.
T7	Control	-	-

Table 2: Incremental Cost-Benefit Ratio of Dinotefuran treatments and its combinations against *Nilaparvatha lugens* (Stall) in paddy (2017)

Treatments	Yield (q/ha)	Increase in yield over Control (%)	Additional cost due to application of insecticide (Rs/ha)	Additional returns from Produce over Control (Rs/ha)	Additional net profit from Produce over Control (Rs/ha)	Incremental benefit cost ratio	Rank
T1	70.95	10.38	1400	6136.4	4736.4	3.38	IV
T2	72.33	12.52	1600	7406	5806	3.63	III
T3	74.9	16.52	2100	9770.4	7670.4	3.65	II
T4	78.66	22.37	2800	13229.6	10429.6	3.72	I
T5	70.33	9.41	1600	5566	3966	2.48	V
T6	67.51	5.02	1300	2971.6	1671.6	1.29	VI
T7*	64.28	-	-	-	-	-	-

*Control Treatment (T7)

Table 3: Number of BPH(adults and nymph/hill in different treatments and dosages along with yield

Treatments	Dosage g/ha	BPH/hill Kharif 2017 (1 st spray)						BPH/hill Kharif 2017 (2 nd spray)						Yield qt/ha
		ADBS	3DAS	7DAS	10DAS	15DAS	%ROC	ADBS	3DAS	7DAS	10DAS	15DAS	%ROC	
T1 Dinotefuran 20% SG	20	20.42 (4.52)	4.37 (2.08)	3.87 (1.95)	3.25 (2.06)	3.11 (2.02)	87.88	19.25 (4.50)	2.83 (1.67)	3.12 (1.78)	3.04 (2.01)	2.93 (1.99)	89.34	70.95
T2 Dinotefuran 20% SG	25	21.27 (4.60)	2.53 (1.58)	2.27 (1.51)	2.18 (1.78)	2.06 (1.75)	91.97	10.69 (3.42)	1.80 (1.34)	1.76 (1.34)	1.66 (1.63)	1.51 (1.58)	94.51	72.33
T3 Dinotefuran 20% SG	30	19.37 (4.38)	1.63 (1.27)	1.60 (1.26)	1.49 (1.57)	1.42 (1.55)	94.47	2.22 (1.79)	1.17 (1.07)	1.25 (1.11)	1.18 (1.47)	1.06 (1.43)	96.15	74.90
T4 Dinotefuran 20% SG	40	22.00 (4.57)	0.86 (0.93)	0.73 (0.85)	0.67 (1.29)	0.57 (1.25)	97.98	0.93 (1.39)	0.63 (0.77)	0.82 (0.92)	0.70 (1.30)	0.62 (1.27)	97.75	78.66
T5 Imidacloprid 17.8 SL	22.5	21.50 (4.63)	4.53 (2.35)	4.43 (2.23)	4.76 (2.19)	4.87 (2.20)	82.73	7.41 (2.90)	3.34 (1.82)	3.47 (1.86)	3.32 (2.07)	3.34 (1.82)	87.85	70.33
T6 Buprofezin 25% SC	187.5	23.37 (4.83)	1.90 (1.37)	2.06 (1.75)	2.53 (1.58)	3.87 (1.95)	84.91	17.93 (4.35)	2.93 (1.99)	3.11 (2.02)	3.24 (2.06)	3.25 (2.06)	88.74	67.51
T7 Control	----	21.70 (4.66)	22.6 (4.85)	23.5 (4.76)	24.18 (5.01)	25.64 (5.16)	-----	64.58 (8.09)	18.23 (4.28)	25.97 (5.09)	26.73 (5.26)	27.48 (5.33)	-----	64.28
S _e m _± CD @ 5% CV		NS	0.10 0.31 4.08	0.12 0.36 5.45	0.24 0.76 7.82	0.21 0.65 6.58		0.46 1.45 4.61	0.14 0.43 6.05	0.46 1.45 4.61	0.06 0.20 3.11	0.11 0.36 3.68		0.28 599.11 4.72

Values in parantheses are $\sqrt{x+1}$ transformed values

Table 4: Effect of insecticides on natural enemies associated with *Nilaparvata lugens* during kharif 2017-18

Treatments	Dosage g or ml a.i/ha	Pre-treatment					15 days after treatment 1 st Spray				15 days after treatment 2 nd spray			
		BPH/hill	MB/hill	Spider/hill	BPH/MB	BPH/Spider	BPH/hill	Mean MB/hill	BPH/MB	Mean Spider/hill	BPH/hill	Mean MB/hill	BPH/MB	Mean Spider/ hill
T1 Dinotefuran 20% SG	20	20.42 (4.62)	3.07 (2.01)	3.06 (2.00)	6.65	6.67	3.11 (2.02)	2.23 (1.79)	1.39	3.43 (2.06)	19.25 (4.50)	2.65 (1.91)	7.26	3.43 (2.09)
T2 Dinotefuran 20% SG	25	21.24 (4.71)	2.93 (1.98)	3.30 (2.06)	7.25	6.43	2.06 (1.75)	1.71 (1.64)	1.20	3.14 (1.92)	10.69 (3.42)	1.50 (1.58)	7.12	3.14 (1.98)
T3 Dinotefuran 20% SG	30	19.37 (4.50)	3.12 (2.03)	2.70 (1.87)	6.20	7.17	1.42 (1.55)	1.26 (1.50)	1.12	3.11 (1.97)	2.22 (1.79)	1.47 (1.57)	1.51	3.14 (1.98)
T4 Dinotefuran 20% SG	40	21.33 (4.70)	3.01 (2.00)	2.59 (1.84)	7.30	8.23	0.57 (1.25)	0.98 (1.40)	0.58	3.02 (1.99)	0.93 (1.39)	1.44 (1.56)	0.64	3.02 (1.93)
T5 Imidacloprid 17.8 SL	22.5	21.50 (4.71)	3.27 (2.06)	3.55 (2.13)	6.57	6.05	4.87 (2.20)	0.51 (1.22)	9.54	2.35 (1.81)	7.41 (2.90)	0.62 (1.27)	11.95	2.35 (1.76)
T6 Buprofezin 25% SC	187.5	23.37 (4.93)	2.83 (1.95)	3.02 (1.93)	8.25	7.73	3.87 (1.95)	0.85 (1.36)	4.55	3.59 (2.05)	17.93 (4.35)	0.97 (1.40)	18.48	3.59 (2.14)
T7 Control	----	21.70 (4.74)	2.81 (1.95)	3.16 (1.94)	7.72	6.86	25.64 (5.16)	3.02 (2.00)	8.49	3.59 (2.08)	64.58 (8.09)	6.64 (2.76)	9.72	3.59 (2.14)
S.em±		2.18	0.05	1.00			0.21	0.05		0.89	0.46	0.12		0.69
CD		NS	0.17	NS			0.65	0.17		NS	1.45	0.38		NS
CV		17.78	6.52	56.73			6.58	6.52		48.60	4.61	9.75		37.82

Values in paranteses are $\sqrt{x+1}$ transformed values

Results and Discussion: The insecticides were tested under field condition on the basis of number of BPH per hill. It is clear from the result that the brown plant hopper population did not vary significantly among the treatments before the application of insecticides. At 3 day after spraying of the Dinotefuran 20% SG at 40 and 30 g ai/ha recorded lowest number of BPH per hill followed by Buprofezin. Upto 15 days after 1st spray Dinotefuran 20% SG at 40 and 30 g a.i/ha maintained the brownplant hopper under normal limit. Highest per cent reduction of BPH @97.98 was recorded by Dinotefuran 20SG @40 g both at 1st spray followed by next higher dose Dinotefuran 20SG @30g recorded 94.47 per cent reduction over control at 1st spray. Same trend was noticed after 2nd spray also. In both the sprayings population of brown plant hopper considerably reduced after 3 day of spraying and continued even after 15 days. Lowest population was recorded in Dinotefuran 20% SG at 40 and 30g ai/ha which were statistically at par through out the observation. Dinotefuran 20% SG and 30 g ai/ha were recorded as the best treatments over Imidacloprid and Buprofezin. The grain yield in the treatment Dinotefuran 20% SG @ 40g a.i./ha recorded significantly highest yield of 78.66q/ha followed by Dinotefuran 20% SG @ 30g a.i./ha (74.90q/ha) and buprofezin @ 187.5g a.i./ha (67.51q/ha) in the lower dose of Dinotefuran 20% SG@ 20g a.i./ha and imidacloprid @ 22.5g a.i./ha recorded yield of 70.95 and 70.33q/ha respectively. The lowest yield of 64.28q /ha was recorded in the untreated control.

The results of the present study are on par with the findings in the management of BPH namely Kharbade et al., (2015), Wang et al., (2008), Shashank *et al.*, (2012) and Kumar et al., (2017) Further, several reports in the literature indicated the effectiveness of insecticides for management of BPH such as Kendappa et al., 2005; Hegde and Nidagundi, 2009; Suri et al., 2012) and Konchada et al., (2017) also studied different dose of insecticides and its impact on control over BPH and also on grain yield of rice.

Effect of insecticides on natural enemies associated with Brownplant hopper: population of natural enemies viz., mired bug and spider found to be moderate to good throughout the experimental period. Fluctuations in the population level of mired bug and spider population noticed among all the insecticidal treatments. It was mainly due to toxicity of insecticides that implies the survivability of natural enemies. Population of mired bug was found to be highly dependent on the availability of brown planthopper for preying. The population of spiders and

Comment [h34]: bioefficacy of insecticides was evaluated

Comment [h35]: the results revealed that

Comment [h36]: The lowest number of BPH population per hill was recorded after 3rd day of spraying.....followed by..... the first spray ofwas observed to maintain the population below ETL upto 15th day after spray.

Comment [h37]: Mean per cent

Comment [h38]: Which

Comment [h39]: @

Comment [h40]: was

Comment [h41]: Rewrite as, the highest grain yield() was recorded in Dinotefuran 20% SG @ 40g a.i./ha followed byand..... in comparison to untreated control...

Comment [h42]: Were observed In conformity with the findings in the management of BPH (.....)

Comment [h43]: Mirid

mired bug is directly proportional to brown planthopper population noticed on crop. The population of natural enemies was more with availability of brown planthopper and vice versa in untreated plot. It is evident from the table 3 that mean number of mired bug per hill after 15 days after first spray was comparatively low in all insecticide treated plot than the untreated control plot. Considerable increase in mired bug population ratio was maintained in dinotofuran treated plot that implies its safety to mired bug. Same trend was noticed after second spray also. Table 3 showed that up to 15 days after both sprays there was no significant effect of insecticides on the mean number of spider populations.

Comment [h44]: Was observed

Comment [h45]: proportional

Comment [h46]: the data presented in Table 3 revealed that.....

Comment [h47]: the results further revealed that (Table 3) there was no significant effect of insecticides on mean number of spider population upto 15 days after both the sprayings.

Incremental Cost-Benefit Ratio

The study has made an attempt to analyse the Incremental Cost-Benefit Ratio of Dinotofuran treatments and its combinations against *Nilaparvatha lugens* (Stål) in paddy in the year 2017. The findings of Incremental benefit in yield and cost over control are presented in table.4 . The results are based on prevailing costs of inputs and market selling price of rice (BPT-5204 @ Rs. 920/q), the Incremental Cost Benefit Ratio (ICBR) was carried out to analyse the economics of different treatments. The data presented in (Table no. 4) indicated that, the treatment (T4) was the most economically viable treatment recording highest ICBR (3.72) due to its high yield and additional returns over the control which stands 1st rank among all the treatments. The second rank was observed in treatment (T3), the per cent increase in the yield was 16.52% over the control with incremental benefit cost ratio was 3.65. and least incremental benefit cost ratio was found in case of treatment (T6). These are two treatment found highest in yield and cost in the study period.

Comment [h48]: During the study an attempt was also made to determine the.....

Comment [h49]: Small'i

Comment [h50]: During the study the two treatments T3 and T4 recorded the highest output in yield and highest cost involved

CONCLUSION

In Asia and Pacific regions, the Brown planthopper *Nilaparvata lugens* (Stål) is major sucking pest of rice and reasoned as a major threat to rice production. There is need to know the observations on the life-history and bionomics of this pest, which is a prerequisite for developing and implementing effective chemical control measures. The control strategy that has been proven effective against brown planthopper *Nilaparvatha lugens* (Stall) using Dinotofuran 20% SG on paddy. The experimental results clearly indicates that the Dinotofuran 20% SG@ 40 g a.i./ha effectively controlled Brown Planthopper on rice. This treatment showed better performance than its lower dosages and the standard checks i.e. buprofezin 25% SC @ 187.5g a.i. and imidacloprid 17.8% SL @ 22.5g a.i. It is concluded that, Dinotofuran 20%

Comment [h51]: One of the

Comment [h52]: serious

Comment [h53]: Present study revealed the effective chemical management strategy in rice against BPH using Dinotofuran 20% SG.

SG@ 40g a.i./ha may be recommended for the control of Brown Plant hopper on rice in Northeastern Dry Zone and areas having similar geographical and climatic conditions. Further, the treatment (T4) was the most economically viable treatment recording highest ICBR (3.72) due to its high yield and additional returns over the control which stands 1st rank among all the treatments. Hence, keeping in view its cost-efficacy and effective control of BPH, the same is recommended to farmers for its suitable incorporation towards pest management of against *Nilaparvatha lugens* (Stall) in paddy.

Comment [h54]: Pl see if the bioefficacy of the highest dose is at par with the penultimate dose, it is recommended to select the penultimate dose instead of selecting higher dose for recommendation.

References

- Chung, T.C., Sun, C.N. and Hung, C.Y. 1982. Resistance of *Nilaparvatha lugens* to six insecticides in Taiwan. *J. Economic Entomology*, 75:199-200
- I.C.Pasalu, N.V. Krishnaiah, G. Katti, N.R.G. Varma., 2002. IPM in Rice, *IPM Mitr*. 45-55
- Kulshreshtha, J.P. 1974. Field problems in Brown plant hopper epidemics in Kerala (India) *Rice Entomological News letter* 1:3-4
- Liu, Z.W., Han, Z.J. and Liu, C.J. 2003. Selection for Imidacloprid resistance in *Nilaparvatha lugens*: cross resistance patterns and possible mechanism. *Pest Management Science*, 59:1355-1359
- Pathak, M.D. and Dhaliwal, G.S., 1981. Trends and Strategies for rice insect problems in Tropical Asia. *IRRI. Res. Pap. Ser.* 64, 5-6
- Patcharin, K., 2011. Brown planthopper *Nilaparvatha lugens* and pest management in Thailand, Conferences International Research on food security, *Natural Resources Management and Rural Development*. Pp, 5-7
- Saxena, R.C. and Singh, R.K., 2003. Rice research in India and the Asian perspective, Asian Biotech Dev. Rev. Neem Foundation, Gurgoan, India. Formerly with IRRI, Philippines. 203, 81-96
- Sindhu, G.S., 1970. Need for variety resistant to white backed plant hopper in Punjab. *International Rice Research News letter* 14:6-7
- Singh, S.P., 2000. Bio intensive approach helpful. *The Hindu Survey of Indian Agriculture*. 159-163
- Whalon, M.E., Mora-Sanchez, R.M. Hollingworth., 2008. Analysis of global pesticides resistance in arthropods. *Global pesticides Resistance in Arthropods*. Michigan State University, USA. 5-31
- Cheng J. Rice planthopper problems and relevant causes in China. Planthoppers: New threats to the sustainability of intensive rice production systems in Asia. 2009;157:178.
- Rahaman MM, Islam K, Jahan M, et al. Relative abundance of stem borer species and natural enemies in rice ecosystem at Madhupur, Tangail, Bangladesh. *Journal of The Bangladesh Agricultural University*. 2016;12:267-272

- Savary S, Willocquet L, Elazegui FA, et al. Rice pest constraints in tropical Asia: Quantification of yield losses due to rice pests in a range of production situations. *Plant Disease*. 2000;84:357-369
- Nirmala B, Muthuraman P. Economic and constraint analysis of rice cultivation in Kaithal district of Haryana. *Indian Research Journal of Extension Education*. 2016;9:47-49
- Rao NV, Maheshwari TV, Prashad PR, Naidu VG, Savithri. In integrated pest management. *Agrobios, India* 2003,89-90
- Kharbade SB, Chormule AJ, Tamboli ND. Bio-efficacy of granular insecticides against *Nilaparvata lugens* (Stål.) in rice under field condition. *Annals of Plant Protection Sciences*. 2015; 23:250-252.
- Wang Y, Gao C, Xu Z, Zhu YC, Zhang J, Li W et al. Buprofezin susceptibility survey, resistance selection and preliminary determination of the resistance mechanism in *Nilaparvata lugens* (Homoptera: Delphacidae). *Pest Management Science*. 2008; 64:1050-1056.
- Shashank PR, Mallikarjuna J, Chalam MSV, Madhumathi T. Efficacy of new insecticide molecules against leafhoppers and planthoppers in rice (*Oryza sativa* L.). *International Journal of Plant Protection*. 2012; 5:397-400.
- Kumar RE, Guruprasad GS, Hosamani AK, Srinivas AG, Pramesh D. Bioefficacy of novel insecticides against planthoppers in direct seeded rice. *Plant archives*. 2017; 17(2):1047-1051.
- Kendappa GN, Mallikarjunappa S, Shankar G, Mithyantha MS. Evaluation of new insecticide, Applaud 25 SC (buprofezin) against brown planthopper, *Nilaparvata lugens* (Stål.) (Family: Delphacidae, Order: Hemiptera). *Pestology*. 2005; 29:5-8.
- Hegde M, Nidagundi J. Effect of newer chemicals on plant hoppers and their mirid predator in rice. *Karnataka Journal of Agricultural Sciences*. 2009; 22:511-513.
- Suri KS, Kumar V, Brar DS. Field evaluation of insecticides for the management of rice plant hoppers, *Sogatella furcifera* and *Nilaparvata lugens*. *Indian Journal of Plant Protection*. 2012; 40(2):153-156.
- Konchada D, Chennamasetty RV, Choragudi SR. Evaluation of newer insecticides against brown planthopper, *Nilaparvata lugens* (Stål.) infesting rice. *Chemical Science Review and Letters*. 2017; 6(23):1423- 1427
- KS Matharu and PS Tanwar, Efficacy of different insecticides against brown planthopper, *Nilaparvata lugens* (Stål) in rice, *International Journal of Chemical Studies* 2020; 8(3): 870-873.
- Dhawan AK, Matharu KS, Kumar V. Impact of integrated pest management strategies on pest complex and economics of basmati rice. *Journal of Insect Science*. 2011; 24(3):219-224.
- Pasalu IC, Katti G. Advances in eco-friendly approaches in rice IPM. *Journal of Rice Research*. 2006; 1(1):83-90
- Alam MJ, Das G. Laboratory evaluation of buprofezin on the mortality of brown planthopper, *Nilaparvata lugens* (Delphacidae: Hemiptera). *Journal of Entomology and Zoology Studies*. 2017; 5(6):2172-2175.