

Original Research Article

Effect of Different Biofertilizers And Organic Manures On Growth, Yield And Quality Of Strawberry (*Fragaria × ananassa* Duch.) cv. Chandler.

ABSTRACT

The present investigation entitled “Effect of Different Biofertilizers and Organic Manures on Growth, Yield and Quality of Strawberry (*Fragaria×ananassa* Duch.) cv. Chandler.” An experiment was conducted at research field, Department of Horticulture, Allahabad School of Agriculture, Sam Higginbottom Institute of Agriculture, Technology & Sciences, Allahabad during the year 2015-2016, the experiment was laid out in a Randomized Block Design having 12 treatments and three replications of bio-fertilizers and organic manure at different levels. The result revealed that treatment T₁₂ (Phosphobacter 5kg/ha + Azatobacter 5 kg/ha + FYM 6 ton/ha + Vermicompost 3 ton/ha + Poultry Manure 3 ton/ha) was found to be best in terms of maximum plant height (19.67 cm), plant spread (26.43cm), number of leaves (18.13), petiole length (4.96 cm), day to first flowering (67.33), number of flowers per plant (16.80), fruit yield per plant (75.94 g), fruit yield per plot (683.46gm), fruit yield per hectare(6.83 t), TSS (8.54 °B), acidity (0.46%), pH of juice (4.40). Similarly the maximum gross return (1079250 rupees), net return (696022.95 rupees) and benefit cost ratio (2.77:1) for cultivation of strawberry under Allahabad agro- climatic condition. However, since these results are based on one-year experiment further trails may be needed to substantiate the same.

Key words: Strawberry, Azatobacter, Phosphobacter , FYM, Vermicompost, Poultry Manure and Yield.

INTRODUCTION

The area under fruit production in India is 7216 thousand hectares with a production of 88977 thousand million tons (NHB, 2014). Strawberry (*Fragaria x sp.*) is native of temperate regions, but varieties are available which can be cultivated in subtropical climate. Strawberry is a

delicious fruit taken fresh in several ways. It is a soft and a highly perishable fruit, often shipped in frozen condition in Western countries. Strawberry thrives best in temperate climate. It is a short day plant; the varieties grown in milder subtropical climate do not require chilling and continue to make some growth during winter. Strawberry requires a well-drained medium loam soil, rich in organic matter. The soil should be slightly acidic with pH from 5.7 to 6.5. At higher pH root formation is poor. The presence of excessive calcium in the soil causes yellowing of the leaves. In light soils and in those rich in organic matter, runner formation is better. Strawberry should not be cultivated in the same land for a number of years. It is preferable to plant it in green manure field. Thus, there is a tremendous pressure on non-renewable energy resource to meet the ever-increasing demand of quality fruits. Hence, it has become imperative to turn to more **econ-friendly** method of nutrition management in horticultural crops of which biofertilizers stand with prime importance. The beneficial effects of Azotobacter are not only due to its ability to fix nitrogen but also due to its ability to synthesize auxin vitamins, growth substance and its capacity to produce anti-fungal antibiotics which improve plant stands in inoculated fields (**Kumari et al. 1975**) by inhibiting root pathogens. Azotobacter is heterotrophic aerobic bacterium, the free living bacteria fixes nitrogen in the rhizosphere and provides it to the plant. Their inoculation is useful for horticultural crop with an increase in plant yield. The increase in yield is because in addition to adding nitrogen, these bacteria produce vitamins like biotin, folic acid also produces antifungal antibiotics which results in an early development of roots and better starts of crop plant initially. Azotobacter is free living nitrogen fixing bacteria fixing nitrogen **equivalent** to 25 to 30 kg nitrogen ha⁻¹. It is also found that crop plants inoculated with Azotobacter survive better in drought conditions.

MATERIALS AND METHODS

The present experiment was conducted at pomology section, Department of Horticulture, SHIATS, Allahabad, during 2015-2016, in a randomized block design replicated with thrice, the

experimental site is situated at a latitude of 20° and 15° North and longitude of 60° 3` East and at an altitude of 98 meters above mean sea level (MSL). Minimum temperature ranged from 4° -5° C (during Oct - Feb) and maximum temperature ranged from 45° -48° C (during March - June). One cultivar with uniform sized strawberry runners were planted during November 2015, maintaining a spacing of 30 X 30 cm. The biofertilizer and organic manures were prepared as per **the requirement** and applied to each treatments and replication at before planting and observations recorded. The recommended package of practices was followed for raising the successful crop. Data on plant growth fruit yield and quality of strawberry characters were recorded when the plants were fully grown. Treatments [T₁ Control(RDF), T₂ (Phosphobacter 5kg/ha + Azatobacter 5 Kg/ha), T₃ (Phosphobacter 5kg/ha + Azatobacter 5 Kg/ha + FYM 25 t/ha), T₄ (Phosphobacter 5kg/ha + Azatobacter 5 Kg/ha + Poultry manure 12 t/ha), T₅ (Phosphobacter 5kg/ha + Azatobacter 5 kg/ha + Vermicompost 12 t/ha), T₆ (Phosphobacter 5kg/ha + Azatobacter 5 kg/ha + FYM 13 t/ha), T₇ (Phosphobacter 5kg/ha + Azatobacter 5 kg/ha + Poultry Manure 6 t/ha), T₈ (Phosphobacter 5kg/ha + Azatobacter 5 kg/ha + Vermicompost 6 t/ha), T₉ (Phosphobacter 5kg/ha + Azatobacter 5 kg/ha + FYM 13 t/ha + Poultry manure 6 t/ha), T₁₀ (Phosphobacter 5kg/ha + Azatobacter 5 kg/ha + FYM 13 t/ha + Vermicompost 6 t/ha), T₁₁ (Phosphobacter 5kg/ha + Azatobacter 5 kg/ha + Vermicompost 6 t/ha + Poultry Manure 6 t/ha), T₁₂ (Phosphobacter 5kg/ha + Azatobacter 5 kg/ha + FYM 6 t/ha + Vermicompost 3 t/ha + Poultry Manure 3 t/ha)]

RESULTS AND DISCUSSION

Growth parameters:

The mean performance of the treatments for growth parameters like plant height, plant spread, numbers of leaves per plant and petiole length have been presented in table no.1 showing significant differences for growth parameters. Vigorous maximum plant height was observed with T₁₂ (19.67cm) followed by T₁₀ (19.14cm). Minimum plant height was observed with T₁ (15.27cm), other treatments showed moderate plant height. Maximum plant spread in treatment T₁₂ (26.43 cm), followed by T₁₀ (25.00 cm). The plant spread was lowest in the T₁ (20.27 cm). The remaining treatments recorded medium plant spread with significantly. Number of leaves per plant was treatment T₁₂ had the maximum number of leaves (18.13), followed by treatment T₁₀ (17.60). The least number of leaves was observed in the treatment T₁ (12.47). Maximum Petiole length in treatment T₁₂ (4.96 cm), followed by T₁₀ (4.77 cm). The Petiole length was lowest in

the T₁ (3.41 cm). These increases may be due to of application of biofertilizer and organic manure. Similar results reported by **Wang *et al.*, (1997)**.

Flowering and fruiting characters:

The data pertaining to the flowering and fruiting characters like **Days taken to first flowering**, number of flowers per plant, number of fruits per plant and fruit yield per plant was recorded under one cultivar of strawberry and 12 treatments presented in table no. Maximum **Days taken to first flowering** (67.33) followed by T₁₀ (69.20). Minimum (70.47). **Maximum fruit yield per plant (75.94 g) was observed in treatment T₁₂ followed by treatment T₁₀ (68.70 g). Minimum fruit yield per plant (39.95 g) was observed in treatment T₁. Maximum fruit yield per plot (683.46 g) was observed in treatment T₁₂ followed by treatment T₁₀ (618.30 g). Minimum fruit yield per plant (359.55 g) was observed in treatment T₁. Maximum fruit yield per hectare (6.83 t) was observed in treatment T₁₂ followed by treatment T₁₀ (6.18 t). Minimum fruit yield per plant (3.59 t) was observed in treatment T₁. Similar results observed by (Chavez and Ferrera, 1990).**

Fruit quality parameters

The outcome of the investigations revealed that TSS content, acidity, pH of the fruit juice and benefit cost ratio (table no. 1) Significantly maximum TSS content (8.54 °Brix) was noticed with T₁₂ followed by treatment T₁₀ (8.46 °Brix). The minimum TSS content (6.90 °Brix) was recorded in T₁. Minimum acidity score (0.46%) was noticed with T₁₂ which was statistically at par with (0.49 %) followed by treatment T₁₀. Maximum acidity score (0.75 %) was recorded in T₁. Maximum pH of the fruit juice (4.40) was noticed with T₁₂ followed by treatment T₁₀ (4.30). The minimum pH of the fruit juice (3.9) was recorded in T₁. Maximum benefit cost ratio (2.27:1) was recorded under treatments T₁₂. Minimum benefit cost ratio (1.39:1) was recorded in treatment T₁ (control). However, better qualities of fruits obtained from the cultivar during winter season may be attributed due to application of different bio fertilizers and organic manures. Similar results observed by **Umar *et al.*, (2010)**

CONCLUSION

Considering the result of the presented investigation. It is concluded that treatment combination of Phosphobacter 5kg/ha + Azatobacter 5kg/ha + FYM 6 t/ha + Vermicompost

3 t/ha + Poultry Manure 3 t/ha was the best treatment for growth, yield and fruit quality of strawberry and **2.27:1** maximum benefit cost ratio. The experiment may be repeated to substantiate the findings which are on the basis of one year trial.

Reference

- Abuzahara, T.R. and Tahboub.A.B.(2008)** Effect of organic matter sources on chemical properties of soil and yield of strawberry under organic farming condition. *J. of World applied sciences*. **5**(3):383-388.
- Akath Singh and Singh, J. N. (2009)** Effect of biofertilizers and bioregulators on growth, yield and nutrient status of strawberry cv. Sweet Charlie. *Indian Journal of Horticulture*. **66**(2): 220-224.
- Cabilovski1,RankoManojlovic,MajaBogdanovic,DarinkaMagazin,NenadKeserovic,Ran**
(2014) Mulch type and application of manure and compost in strawberry (*Fragaria × ananassa* Duch.) production: impact on soil fertility and yield. ISSN 1392-3196 / e-ISSN 2335-8947 *Zemdirbyste-Agriculture*, **(1)**101: 67–74.
- Chavez, M. G. and Ferrera, R. (1990)**. Effect of vesicular arbuscular **mycorrhizae** on tissue culture-driven plantlets of strawberry. *Hort. Science*. 25:903-905.
- Geddeda, Y. I., Trappe J. M. and Stebbins R. L., (1984)**. Effect of vasicular arbuscular mycorrhizae and phosphorus on apple seedlings. *J. of the American Society of Horticultural Sciences*. **109**(1):24-27
- Iqbal, U., Wall, V. K., Ravi K., Mahital J., (2009)**. Effect of fym, urea and *Azotobacter* on growth, yield and quality of strawberry cv. Chandler. *Notulae Botanicae, Horti Agrobotanici, Cluj-Napoca* **(1)**37:139-143.
- Karlidag, H.; Yildirim, E.; Turan, M.; Donmez, M. F. (2010)** Effect of plant growth-promoting bacteria on mineral-organic fertilizer use efficiency, plant growth and mineral contents of strawberry (*Fragaria x ananassa* L. Duch.). Hohenheim, Germany, **(22)**: 218-226.

Kumari L., Kavimadanan M. S. and Rao K. N. S. (1975). Occurance of nitrogen fixing spirillum in root of sorghum, maize and other plants. *Indian J. of Exp. Biol.* 14:638-639.

Macit, I., Koc, A., Guler, S., Deligoz, L. (2007) Yield, quality and nutritional status of organically and conventionally-grown strawberry cultivars. *Asian J. of Plant Sciences.* 7(6): 1131-1136.

Rana, R. K. and Chandel, J. S. (2003) Effect of biofertilizers and nitrogen on growth, yield and fruit quality of strawberry. *Progressive Horticulture.* 3(5)1:25-30.

Uddin M R, M F Hossain, S M Zaman, M Islam and N Ara. (2013). Effect of organic manure on growth and yield of strawberry. *Wudpecker Journal of Agricultural Research* ISSN 2315-7259 Vol. 3(1):35 – 38.

Umar, I.; Wali, V. K.; Rehman, M. U.; Mir, M. M.; Banday, S. A.; Bisati, I. A. (2010) Effect of subabul (*Leucaena leucocephala*), *Applied Biological Research.* (2)12:50-54.

Wang, S. S., Patil, M. T. and Singh, B. R. (1997). Cultivar x Biofertilizer interaction study in strawberry *Recent Horticulture.* 4: 43-44.

Yavari, S., Eshghi, S.; Tafazoli, E., and Yavari, S. (2008). Effects of various organic substrates and nutrient solution on productivity and fruit quality of strawberry 'Selva' (*Fragaria x ananassa* Duch.). *Journal of Fruit and Ornamental Plant Research.* (16)34:167-178.

Table 1 : Effect of Different Biofertilizers And Organic Manures On Growth, Yield And Quality Of Strawberry (*Fragaria × ananassa* Duch.) cv. Chandler.

Treatment	Plant height (cm)	plant spread (cm)	number of leaves	petiole length (cm)	Days taken to first flowering	Number of flower per plant.	fruit yield per plant	Fruit yield per plot (g)	Fruit yield per ha (tonns)	TSS of fruit	Fruits acidity	pH of the fruit juice	Cost benefit ratio
T ₁	15.27	20.27	12.47	3.41	70.47	11.33	39.95	359.55	3.59	6.90	0.75	3.9	1.39
T ₂	16.29	22.10	15.33	3.84	70.27	12.73	46.68	420.09	4.19	7.70	0.56	4.0	1.45
T ₃	16.35	21.33	15.13	3.77	69.33	12.60	44.64	431.76	4.31	7.78	0.62	3.92	1.79
T ₄	16.29	22.37	14.47	3.80	69.53	12.80	48.14	433.29	4.33	7.61	0.61	4.0	2.13
T ₅	16.77	22.23	14.93	3.93	69.60	12.53	46.37	417.36	4.17	7.69	0.57	4.05	2.05
T ₆	16.61	22.63	14.53	3.87	69.87	12.67	46.31	416.76	4.16	7.68	0.63	4.10	1.10
T ₇	16.91	22.07	15.20	3.85	69.27	12.00	50.65	455.88	4.55	7.83	0.54	4.10	1.69
T ₈	16.90	22.00	14.80	4.11	69.47	13.20	55.86	502.74	5.02	7.80	0.70	4.10	1.72
T ₉	16.79	22.73	14.87	4.11	69.80	12.80	52.51	472.56	4.72	7.67	0.60	4.15	1.98
T ₁₀	19.14	25.00	17.60	4.77	69.20	16.07	68.70	618.30	6.18	8.46	0.49	4.30	1.49
T ₁₁	18.74	23.80	16.07	4.32	69.80	15.20	60.60	545.37	5.45	8.34	0.51	4.34	1.67
T ₁₂	19.67	26.43	18.13	4.96	67.33	16.80	75.94	683.46	6.83	8.54	0.46	4.40	2.27
F - test	S	S	S	S	S	S	S	S	S	S	S	S	
CD (0.05%) =	0.76	1.27	0.83	0.20	3.61	0.68	3.34	21.95	0.22	0.41	0.03	0.28	
CV (%) =	3.32	3.31	4.06	3.67	3.88	3.81	4.70	3.42	3.42	3.93	4.45	5.03	
S Ed (±)	0.37	0.61	0.41	0.10	1.76	0.33	1.63	10.71	0.11	0.20	0.02	0.14	

UNDER PEER REVIEW