

## Original Research Article

### **Comparative Morphometric studies of European honey bee (*Apis mellifera* L.) at different altitudes of Kashmir Region**

#### **Abstract**

European honey bee (*Apis mellifera* L.) was introduced in Kashmir during late 19<sup>th</sup> century and since, then its domestication is continued in diverse conditions of Kashmir region. It appears well adapted to unique climatic conditions of the valley. In order to analyse diversification in its characters during ecological adaptations to different altitudes and climatic conditions, morphological characterization of *Apis mellifera* was employed at different altitudes of Kashmir region. Samples of honey bees were obtained from nine locations representing three different altitudes viz., high, medium and low of Kashmir. Total of 17 characters were measured at three different altitudes. Highly significant difference ( $p \leq 0.01$ ) in 15 characters were observed among three altitudes. The three main characters viz., proboscis length, fore wing length and cubital index were compared with earlier data as reported by Ruttner (1988). The proboscis length ( $6.35 \pm 0.06$ mm) at high altitude, forewing length ( $9.21 \pm 0.01$ mm) at mid altitude, cubital index at low altitude ( $2.55 \pm 0.08$ mm) were found same or close to *A. mellifera ligustica*. However, other parameters did not support presence of this sub-species, therefore, it can be conclude that *A. mellifera* might have undergone degradations in its certain characters after its introduction in Kashmir region due to diverse climate conditions

**KEYWORDS:** Morphometrics, *Apis mellifera*, Diversification, Proboscis, Degradation, Cubical Index, Kashmir.

#### **Introduction**

Honey bees are of huge economic importance, plays a vital role in pollination of fruit crops, vegetables and wild plants. They are capable of producing world's amazing sweetener, "honey" along with other important product of commercial value such as royal jelly, wax, propolis, etc. A honey bee colony comprises a queen, hundreds of drones and thousands of worker bees. The workers are always firm and perform different duties including honey, pollen collection, etc. Honeybee (*Apis mellifera* L.) is widespread in Africa, Europe and Western Asia (Miguel *et al* 2007) and comprises of 26 subspecies according to morphometric analyses (Miguel *et al* 2007, Engel M 1999). The state of Jammu and Kashmir is one of the most important beekeeping areas in India with at least four agro climatic zones ranging from low altitude subtropical, intermediate, temperate and cold alpine. Kashmir lies in the Northern Hemisphere, has a moderate climate, which is largely defined by its geographic location. As a large valley with significant differences in Geo-location among various districts, the weather is often cooler in the hilly areas compared to the flat lower part. *Apis mellifera* L., was introduced in Kashmir during late 19<sup>th</sup> century and since, then its domestication is continued in diverse conditions of state. *Apis mellifera* appears well adapted to unique climatic conditions of Kashmir region. For example, foraging activities, worker and drone brood

rearing, pollen storage and colony population size during the summer was significantly higher than those of other honeybee species, resulted in higher pollen and honey production. Higher pollen and honey production does not only depend upon the floral abundance of the area but also on the morphological characters of workers such as body size, proboscis length, corbícula, etc. (Alburaki and Alburaki, 2008). Factors that affect worker body characteristics include pollen and nectar resources (Hela R M *et al*, 2003), content of dietary protein (Roulston T H 2000, Zheng B 2014), geographical location (Al-Kahtani S N, 2014, Charistos L *et al* 2014 and Kulici M and Kume K 2014). Furthermore, the length of proboscis of a particular species of worker bee plays a vital role for honey production as it has positive correlation with foraging (Mostajeran *et al.*, 2006). Besides these features, wing morphometry plays important role in classification of races and their size affects the flight ability of worker honey bees (Horowitz, 1983). However, the honey bee species in different climates may possess varied phenotypic characters and is suspected to have different ecological races. There is often a positive correlation in some characteristics, for instance, the length of the wing and altitude, the size of pollen basket and hind-leg and the size of honey. Such correlation can indicate the importance of the morphologic characteristics in the adaptation of individuals to the environment (Ajao, 2012) which can lead to the evolution of many subspecies (Hepburn and Radloff, 1998). The study of morphometry of honey bees assumes significance for its application in species and subspecies characterization and discrimination. The honey bee races differ in their morphology, physiology and behavioural aspects in relation to their environment (Abou-Shaara *et al* 2011) and the use of their morphological characters is highly important in discriminating among the honey bee subspecies and in preserving the honey bee biodiversity. Worldwide, several workers (Bucó *et al* 1987, Ruttner 1988, Tofliski 2008; Bouga *et al* 2011) have used morphometric techniques to resolve the intraspecific taxonomy of honey bees, specifically for *A. mellifera*.

Morphometric tools along with sophisticated statistical analysis for studies of phenotypes and of the diversification of these bees made it possible to describe changes in the bee body dimensions depending on latitude (Apatov 1925). Originally, discrimination of subspecies was based on descriptive methods but they proved insufficient and were replaced by morphometric methods (Ruttner, 1988). Morphometry is therefore, a powerful tool which can discriminate populations (Cornuet *et al* 1975, Ruttner *et al* 1978). This method uses numeric data resulting from exact measurements of morphological characters from which means of colony characters are obtained for statistical analyses (Ruttner 1988).

In order to analyse the present morphometric status of honey bee *Apis mellifera* in Kashmir region, the present studies were undertaken. Presently, no study has specified morphometric characterization of *Apis mellifera* found throughout Kashmir at three different altitudes. The objective of this study was to determine difference in characters of *Apis mellifera* at different altitude under diverse climatic conditions since its introduction.

## **MATERIALS AND METHODS**

### **Sampling of *A. mellifera* bees from different altitudes**

The samples of *A. mellifera* (worker bees) were obtained from nine locations viz., Narastan, Yarmukam Ganderba, Laam Tral, Damhal Anantnag, Bonpora Shopian Kalaros Kupwara, Bijihama Uri, Anchiodora Anantnag and Dara Srinagar representing three different altitudes viz., high, medium and low of Kashmir region. Before the collection of bees, hive entrance was closed and the worker bees were handpicked gently one by one. Foraging bees

were placed in killing bottles and accordingly anaesthetised with ethyl acetate. The bees were preserved in 70 per cent ethyl alcohol until morphometric analysis.

**Sample size:** Nine stationary apiaries at different locations representing three altitudes of Kashmir region were selected for sampling. Five honey bee colonies per location were taken and from each colony ten bees were collected. Total number of bees per location were fifty. One hundred fifty bees per altitude were analysed for studying different morphometric traits.

### **Methodology for measurement of different morphological traits of honey bee *Apis mellifera* L.**

17 morphological parts of *A. mellifera* L. were dissected out. The dissected parts were mounted on glass slides covered with cover slips. A stereo zoom microscope (Magnus INVI) equipped with Mag-vision software calibrated with micrometre lens, was used for measurement. Some parts like head, thorax, abdomen, hind wing and forewing were measured with the help of measuring scale (mm).

#### **Morphometric characters of *Apis mellifera* studied:**

After carefully detaching the head of *A. mellifera*, it was placed on the slide. The height was measured from the anterior most part to the origin of various appendages and the maximum head width was measured from one side of compound eye to the other. Proboscis of *A. mellifera* was detached carefully with the help of forceps, placed on slide and measured from cardo with sub-mentum up to the end of labellum. The length of flagellum and total antennal length of *A. mellifera* was measured. The sum of length of scape, pedicel and flagellum was considered as total antennal length. The thorax length was measured from cervical portion up to the first abdominal segment dorsally, whereas abdomen length were measured from anterior to posterior abdominal segment dorsally. Slides of the dissected fore wings from selected samples of bees were measured with the help of measuring scale. Length of wing was taken from the humeral angle to the apex and width was measured from costal margin to anal angle. The cubital index of *A. mellifera* was calculated after measuring the length of two wing vein segments (A to B and B to C) of honey bees (Goetze, 1930).  $CI = BC/AB$ . BC = Distance between points B to C, AB = Distance between points A to B. The hind wing length was measured from humeral angle to the apex while the width from costal margin to the anal angle. The number of hamuli were counted under microscope at magnification of 4x. The detached hind legs were analysed by measuring the length of femur, tibia and metatarsus. The breadth of metatarsus was also measured. The morphometric characters studied during the present investigations were based on the previous work of Ruttner *et al* (1978), Sakagami *et al* (1980) and Verma *et al* (1990) on adult worker bees of *Apis mellifera* L., *A. cerana cerana* F. and *Apis dorsata* F. Statistical analysis of the data was carried out by calculating colony means, standard deviation and covariance's of the morphometric characters. Further one way analysis of variance (one-way ANOVA) by SPSS software was applied for testing the significance of results. Significant results were compared on the basis of critical differences.

## **RESULTS**

### **Head height and head width**

The data on head height and head width of *Apis mellifera* at different altitudes of Kashmir region is presented in Table 1. The perusal of data on head height revealed highest mean ( $3.27\pm 0.01$  mm) at Narastan of high altitude and lowest mean ( $3.14\pm 0.07$ mm) was recorded at Achidora at low altitude. Highest mean head width ( $3.71\pm 0.08$  mm) was recorded at Damhal at mid altitude and lowest mean ( $3.61\pm 0.01$ mm) was at Anchidora at low altitudes. At three altitudes the head height and head width ranged from 3.10-3.15mm and 3.28-4.03mm respectively. The data revealed highly significant difference in head height ( $CD=0.0101$ ,  $p\leq 0.01$ ) and significant difference in head width ( $CD=0.0101$ ,  $p\leq 0.05$ ) among three altitude. Non- significant variation in head height and head width was observed within the locations at each altitude.

**Proboscis length, flagellum length and antennal length:** Same trend was followed by proboscis length, flagellum length and antennal length of *Apis mellifera* where highest proboscis length ( $6.37\pm 0.02$  mm), flagelleum length ( $3.81\pm 0.01$  mm), antennal length ( $6.11\pm 0.01$  mm) was recorded at Narastan at high altitude, respectively (table 2). Lowest mean proboscis length ( $6.22\pm 0.07$ mm), flagellum length ( $5.63\pm 0.03$ mm) antennal length ( $3.31\pm 0.03$ mm) was measured at Uri Bijihama, Dara and Anchidora at low altitude, respectively. At three altitudes, the proboscis length ranged from 6.20-6.76mm, flagellum length from 3.22-3.89mm and antennal length ranged from 5.44-6.12mm. The perusal of data revealed highly significant difference in proboscis length ( $CD=0.0101$ ,  $p\leq 0.01$ ), flagellum length ( $CD=0.0305$ ,  $p\leq 0.01$ ) and antennal length ( $CD=0.0143$ ,  $p\leq 0.01$ ) among three altitudes.

#### **Thorax and Abdomen**

The data on thorax length and abdomen length of *Apis mellifera* is presented in Table 3. Highest thorax length ( $4.30\pm 0.01$  mm) was measured at Narastan at high altitude. The lowest mean ( $4.15\pm 0.07$ mm) was recorded at Anchidora. Highest abdomen length ( $6.08\pm 0.04$  mm) was measured at yarmukaam at high altitude and lowest mean ( $5.55\pm 0.02$ mm) was recorded at Anchidora. At three altitudes, thorax length and abdomen length ranged from 3.89-4.56mm and 5.26-6.89mm, respectively. Highly significant difference in thorax length ( $CD=0.0143$ ,  $p\leq 0.01$ ) and abdomen length ( $CD=0.0305$ ,  $p\leq 0.01$ ) among three altitude was observed.

#### **Fore wing length, fore wing width, cubital index, hind wing length, hind wing width, number of hamuli**

The data on forewing length, fore wing width, cubital index, hind wing length, hind wing width, number of hamuli of *Apis mellifera* at different altitudes in Kashmir region is presented in Table 4. The data revealed highest average forewing length ( $9.32\pm 0.02$  mm) of *A. mellifera* at Laam followed by Yarmuqam and Narastan ( $9.29\pm 0.07$  and  $9.23\pm 0.07$  mm) at high altitude. The lowest mean ( $9.17\pm 0.06$ mm) was recorded at Dara at low altitude. At three altitudes, the forewing length ranged from 8.24-9.67.15mm. Forewing width was recorded highest ( $3.06\pm 0.02$  mm) at Laam at high altitude and lowest mean ( $2.96\pm 0.28$ mm) was recorded at Bonpora at mid altitude. The forewing width ranged from 2.09-3.31mm. Cubital index, highest mean ( $2.55\pm 0.08$  mm) was recorded at Uri bijihama at low altitude and lowest mean ( $2.19\pm 0.02$ mm) at Bonpora at mid altitude. At three altitudes, the cubital index ranged from 2.05-2.55mm. Highest hind wing length ( $6.59\pm 0.03$  mm) and hind wing width ( $1.82\pm 0.08$  mm) was recorded at Narastan at high altitude and lowest mean ( $6.19\pm 0.06$ mm) and ( $1.71\pm 0.02$ mm) was recorded at Uri bijihama and Anchidora, respectively. At three altitudes, the hind wing length ranged from 6.01-6.99mm and hind wing width ranged from 1.34-1.98mm. Number of

hamuli of *Apis mellifera* at different altitudes of Kashmir region revealed highest mean ( $23.88 \pm 0.06$ ) at Dara at low altitude and lowest mean ( $18.64 \pm 0.28$ ) was recorded at Damhal. At three altitudes, the number of hamuli ranged from 15-23. The perusal of data revealed highly significant difference in fore wing length ( $CD=0.0203$ ,  $p \leq 0.01$ ), significant difference in fore wing width ( $CD=0.0305$ ,  $p \leq 0.05$ ), highly significant difference in cubital index ( $CD=0.0380$ ,  $p \leq 0.01$ ), hind wing length ( $CD=0.0635$ ,  $p \leq 0.01$ ), hind wing width ( $CD=0.0176$ ,  $p \leq 0.01$ ) and number of hamuli ( $CD=0.04016$ ,  $p \leq 0.01$ ) among three altitude.

### **Femur length, tibia length, metatarsus length, metatarsus breadth**

The data on Femur length, tibia length, metatarsus length, metatarsus breadth of *Apis mellifera* at different altitudes in Kashmir region is presented in Table 5. Femur length of *Apis mellifera* at different altitudes of Kashmir region revealed highest mean ( $2.55 \pm 0.09$  mm) at Narastan. The lowest mean ( $2.33 \pm 0.08$  mm) was recorded at Uri bijihama at low altitude. At three altitudes, the femur length ranged from (2.21-3.35 mm). Tibia length was measured highest ( $2.94 \pm 0.01$  mm) at Laam at high altitude. The lowest mean ( $2.80 \pm 0.08$  mm) was recorded at Bonpora. At three altitudes, the tibia length ranged from (2.70-3.00 mm). Metatarsus length of *Apis mellifera* was recorded highest ( $1.96 \pm 0.02$  mm) at Narastan. The lowest mean ( $1.78 \pm 0.08$  mm) was recorded at Anchidora at low altitude. At three altitudes, the Metatarsus length ranged from (1.34-1.98 mm). Metatarsus breadth of *Apis mellifera* was also found highest mean ( $1.13 \pm 0.07$  mm) at Narastan. The lowest mean ( $1.03 \pm 0.02$  mm) was recorded at Dara at low altitude. Damhal, Bonpora and kalaroos of mid altitude recorded ( $1.07 \pm 0.04$ ,  $1.07 \pm 0.03$ , and  $1.08 \pm 0.03$  mm), respectively. At three altitudes, the metatarsus breadth ranged from (1.02-1.23 mm). The perusal of data revealed highly significant difference in femur length ( $CD=0.0203$ ,  $p \leq 0.01$ ), tibia length ( $CD=0.0143$ ,  $p \leq 0.01$ ), metatarsus length ( $CD=0.0176$ ,  $P \leq 0.01$ ) and metatarsus breadth ( $CD=0.0101$ ,  $p \leq 0.01$ ) among three altitude. Non-significant variation in femur length, tibia length, metatarsus length and breadth was observed within the locations at each altitude.

## DISCUSSION

Earlier too, morphometric studies were carried out by different Indian workers. Kshirsagar (1981) on *Apis cerana* (Pune), Ananda (2000) on *A. cerana* (Bangalore), Mujumdar and Kshirsagar (1986) on *A. dorsata* (Pune), Makhmoor (2001) on four honey bee species (Jammu), Ibrahim *et al* (2017) on *Apis mellifera* (Himachal Pradesh) and Makkar *et al* (2020) on *Apis mellifera jemenitica* (Yemen). Our findings are in conformity with all these above mentioned studies. Sharma (1990), measured head height of *A. mellifera* as 3.19 mm which is near to average head height at mid altitude in Kashmir region as per the present findings. The proboscis length of various geographical races of honey bees has shown considerable variations. Ruttner (1988) worked on proboscis length of different races of *A. mellifera* viz., Australia (6.40 mm), Hungary (6.43 mm), Romania (6.39 mm) and Greece (6.57 mm) which substantiate the present findings on proboscis length wherein slight variations might be due to different geographical races. On subspecies level, the proboscis length was reported to be  $6.07\pm 1.3$ ,  $6.41\pm 0.7$ ,  $6.35\pm 0.79$ ,  $7.06\pm 1.5$ ,  $6.64\pm 0.70$ ,  $6.28\pm 0.89$ ,  $6.43\pm 1.7$  and  $6.41\pm 1.80$  mm in *A. mellifera mellifera*, *A. mellifera carnica*, *A. mellifera ligustica*, *A. mellifera caucasica*, *A. mellifera armeniaca*, *A. mellifera meda*, *A. mellifera anatoliaca* and *A. mellifera pomonella*, respectively (Sheppard and Meixner, 2003). The proboscis length observed at High altitude as  $6.35\pm 0.06$  in the present investigations is in close proximity of *A. mellifera ligustica*. In India, different workers have reported the proboscis length of *A. mellifera* as  $6.37\pm 0.05$  (Sharma, 1990) and 7.00 mm (Jagannadham and Goyal, 1983). Ajao *et al*, (2014) reported the average antenna length of *A. mellifera* as  $5.48\pm 0.015$  which is near to average antenna length at mid altitude in Kashmir region. Dyer and Seeley (1987) reported the fore wing length in *A. mellifera* of different attitudes between 7.64 to 9.70 mm. Jevtic *et al* (2007) reported fore wing length ranged from 7.81 to 10.37 mm and width from 2.12 to 3.75 mm from different parts of Serbia. Alattal *et al*, (2014) reported the length of forewing of 3 clusters of local Saudi honey bees as  $8.79 \pm 0.22$ ,  $8.31 \pm 0.19$  and  $8.04 \pm 0.12$  and width of fore wing as  $3.05 \pm 0.22$ ,  $2.86 \pm 0.06$  and  $2.76 \pm 0.05$  mm. Our findings are in congruence with the report of Bouzeraa *et al*, (2016) who measured cubital index of *Apis mellifera* at three different sites of Algeria as  $2.08\pm 0.20$ ,  $2.50\pm 0.34$  and  $2.53\pm 0.33$ . The mean cubital index  $2.55\pm 0.08$  mm of low altitude in Kashmir region was found same as *A. mellifera ligustica*. AL-Kahtani S N and Taha E-KA (2021) reported that with the exception of the number of hamuli, worker Yemeni bee body size and morphometric parameters related to the colony productivity were smaller than Carniolan bees under environmental conditions of the study region. Earlier, Sharma (1990) reported the femur length in *A. mellifera* as  $2.41\pm 0.05$  mm which is same to the average femur length at mid altitude of the present findings ( $2.41\pm 0.05$  mm). Bouzeraa *et al*. (2016) measured femur length at three different sites of Algeria as  $3.87\pm 0.16$ ,  $3.71\pm 0.09$  and  $3.86\pm 0.12$  mm which is higher than present findings. According to their results, values obtained are closer to giant bee (*Apis dorsata*). However, El-Aw *et al*. (2012) found smaller length (2.22 mm) of femur in Egypt in same bee species. The higher length of femur ranging from 2.60– 2.73 mm has also been reported for different races of *A. mellifera* (Gencer *et al* 2004; Adl *et al* 2007). The tibial length in *A. mellifera* has been reported by different workers as  $3.01 \pm 0.10$  for Indian population (Sharma, 1990),  $2.83\pm 0.08$  mm for Egyptian population and 3.03- 3.22 mm for Iranian population (Adl *et al* 2007) and 3.16 to 3.18 mm for bees population in Turkey (Gencer *et al* 2004). In other studies the dimension of metatarsus were also observed to be (length 2.00 to 2.44 mm, width 1.08 to 1.19 mm) for the same bee species (Alqarni *et al* 2011, Adl *et al* 2007, El-Aw *et al* 2012). Intraspecific variation in body size may reflect quality of the environment

Banaszak C W *et al* (2018) and worker body weight is altered by the availability of nectar and pollen resources (Helal *et al* 2003, Szentgyorgyi H *et al* 2016) and season (Kunert K and Crailsheim K 1988). Ahmad and Bilal (2017) who found positive correlation of bee size with increase in altitude. Narayanan *et al* (1960) also observed a definite trend of variation in *A. cerana* from plains to hills. These observations lend support to present investigation that altitude has impact. Fernando (1979) observed tongue length of bees significantly longer in mountainous districts than lowlands. Present findings also draw the support from Kshirsagar (1976) who reported the bigger size of antenna at higher elevations. This is because of the fact that bees found at the higher elevations require longer antenna in order to have better sense perception, so as to cope with the larger environmental disturbances at these altitudes. Mattu and Verma (2015) reported significant and positive correlation of length of postmentum with altitude. Narayanan *et al.* (1961a, b) reported smaller number of hooks in bees at higher altitudes. Hepburn *et al* (2001) also found smaller number of hooks from Northeast Himalayan region. Size of hind leg especially metatarsus affect the pollen carrying capacity of honey bee (Ruttner, 1979). In Kashmir region, significant differences were observed only in length of femur and breadth of metatarsus of *Apis cerana*. Bees of mountainous zone also showed significantly higher values for these characters as compared to those of sub mountainous zone of Kashmir region (Ahmad and Bilal, 2017). Bouzeraa *et al* (2016) reported highly significant effect concerning the length of femur, tibia, metatarsus ( $p \leq 0.001$ ) and significant effect concerning to the width of metatarsus ( $p \leq 0.05$ ) on bees coming from three different sites of Algeria. Verma *et al* (1994) also found bigger size of hind leg at higher elevations as compared to the plain area.

## Conclusion

The morphometric variations found in *Apis mellifera* L. in Kashmir with respect to altitude particularly, body size, tongue length, length of antenna, number of hamuli, length of femur could have a significant impact in future bee breeding programmes. The mean proboscis length ( $6.35 \pm 0.06$ mm) of high altitude, mean forewing length ( $9.21 \pm 0.01$ mm) of mid altitude, mean cubital index of low altitude ( $2.55 \pm 0.08$ mm) were found same to *Apis mellifera ligustica* when compared with data as reported by Ruttner (1988). The other characters, however, were different in measurement which imply presence of subspecies other than *A. mellifera ligustica* at different altitudes. Hence, *Apis mellifera* L. differing in their morphology in relation to their environment and the use of their morphological characters is highly important in discriminating among the sub-species and in preserving the honey bee biodiversity.

## References

- Abou-Shaara, H.F., Draz, K.A., Al-Aw, M and Eid, K. 2011. Simple method in measuring honey bee morphological characters. In: Proceedings of the 42nd international apiculture congress - Apimondia, pp 222, Buenos Aires
- AdI, M.B.F., Gencer, H.V., Firatii, C. and Bahreini, R. 2007. Morphometric characterization of Iranian (*Apis mellifera meda*) Central Anatolian (*Apis mellifera anatoliaca*) and Caucasian (*Apis mellifera caucasica*) honeybees populations. *Journal of Apicultural Research and Bee World* **46**(4): 225-231.

- Ahmad, S and Bilal, S. 2017. Morphometric variations and expression of body colour pattern of honey bee, *Apis cerana* F. in Kashmir. *Journal of Entomology and Zoology Studies* **5**(4): 364-371.
- Ajao, A.M. 2012. Comparative Studies on Ecology and Morphometrics of Reared and Feral Honeybees in Geological Zones of Kwara State, Nigeria. Phd Thesis submitted to Federal University of Agriculture, *Apidologie*. pp. 1-57.
- Ajao, M.A., Oladimeji, U., Idowu, B.A., Babatunde, K.S and Obembe, A. 2014. Morphological characteristics of *Apis mellifera* L. (Hymenoptera: Apidae) in Kwara State, Nigeria. *International Journal of Agricultural Science* **4**: 171-175.
- Alqarni, A.S., Hannan, M.A., Owayss, A.A and Engel, M.S. 2011. The indigenous honey bees of Saudi Arabia (Hymenoptera, Apidae, *Apis mellifera jemenitica* Ruttner): Their natural history and role in beekeeping. *ZooKeys* 134: 83-98.
- Alattal, Y., Ghamdi, A.H., Sharhi, A.M and Fuchs, S. 2014. Morphometric characterisation of the native Honey bee, *Apis mellifera* L.1758, of Saudi Arabia. *Zoology in the Middle East*. **60**:226-235
- Alpatov, W.W. 1929. Biometric studies on variation and races of the honey bee *Apis mellifera* L. *Quarterly Review of Biology* **4**: 1-57.
- Alburaki, M and Alburaki, A. 2008. Morphometrical study on Syrian honey bees (*Apis mellifera syriaca*). *Emirate Journal of Food and Agriculture* **20**(1): 89-93.
- Al-Kahtani, S.N, Taha, E.K.A. 2014. Morphometric studies on dwarf honeybee *Apis florea* F. workers in Saudi Arabia. *J Apic Sci*. **58**: 127–134.
- AL-Kahtani, S.N, Taha, E-K.A. 2021. Morphometric study of Yemeni (*Apis mellifera jemenitica*) and Carniolan (*A. m. carnica*) honeybee workers in Saudi Arabia. *PLoS ONE* **16** (2): e0247262. <https://doi.org/10.1371/journal>.
- Ananda, M.R. 2000. Studies on morphological characters and behavioural traits of Indian honeybee, *Apis cerana indica* F. from different parts of Karnataka. M.Sc. Thesis, UAS, Bangalore
- Banaszak-Cibicka W., Fliszkiewicz, M., Langowska, A and Żmihorski, M. 2018. Body size and wing asymmetry in bees along an urbanization gradient. *Apidologie* ;**49**: 297–306.
- Bouzeaa, H., Achou, M., Sellami, H. And Slotani. 2016. Study of the morphometric diversity of the population of honeybees (*Apis mellifera*) in North-East Algeria. *European Journal of Experimental Biology* **6**: 6-16.
- Bouga, M., Alaux, C., Bienkowska, M., Büchler, R., Carreck, N.L., Cauia, E., Chlebo, R and Wilde, J. 2011. A review of methods for discrimination of honey bee populations as applied to European beekeeping. *J Apicult Res* **50**(1):51–84

- Buco, S.M., Rinderer, T.E and Sylvester, H.A. 1987. Morphometric differences between south American Africanized and South African (*Apis mellifera scutellata*) honey bees. *Apidologie* 18:217–222.
- Cornuet, J. M., Fresnaye, J. and Tassencourt, L. 1975. Discrimination and classification of honeybee populations by means of biometric characters. *Apidologie* 6: 145-187.
- Charistos, L., Hatjina, F., Bouga, M., Mladenovic, M and Maistros, A.D. 2014. Morphological discrimination of greek honeybee populations based on geometric morphometrics analysis of wing shape. *J Apic Sci.* 58: 75–84
- Dyer, F.C and Seeley, T.D. 1987. Interspecific comparisons of endothermy in honey bees (*Apis*): deviations from the expected size related patterns. *Journal of Experimental Biology* 127: 1-26.
- Engel, M.S., 1999. The taxonomy of recent and fossil honey bees (Hymenoptera: Apidae; *Apis*). *Journal of Hymenoptera Research* 8: 165 – 196.
- El-Aw, M.A., Draz, K.A., Bid, K.S.A and Abo-Shara, H. 2012. Measuring the Morphological characters of honey bee (*Apis mellifera* L.) using a simple Semi-automatic technique. *Journal of American Science* 8(3): 558-564.
- Fernando, E. F. W. 1979. Some biometrical features of *Apis cerana* F. from Sri Lanka. *Indian Bee Journal.* 41: 5-8.
- Gencer, H.V., Baspinar, E and Fira, I .C. 2004. The Graphic evaluations of morphological charactes in hoenybees (*Apis mellifera* L.) by Chernoff Faces. *Tarim Bilimleri Dergisi Journal of Agricultural Sciences* 10(3): 245- 249.
- Goetze, G. 1930. Variability of the ants and honey bees with special consideration of longitivity. *Archive for bees customers* 11: 135-274.
- Hepburn, H.R, Radloff, S.E and Verma, S. 2001. Morphometric analysis of *Apis cerana* population in the Southern Himalayan region. *Apidologie* 32: 435- 444.
- Hepburn, H.R and Radloff, S.E. 1998 - *Honey bees of Africa*. Springer, Berlin, 312
- Helal, R. M., El-Dakhkhni, T. N., Shawer, M. B., Taha, E. K. A. 2003. Effect of moving the apiaries on activity of honeybee colonies. 2- flight activity, gathering of nectar and sugar concentration contents and honey. *J Agric Res Tanta Univ* 29: 268–282.
- Horomitz, R.F.A., 1983. Inbreeding effects in flight muscle mitochondria of *Apis mellifera*. *Rev. Brasil. Genet.* 6: 59-70.
- Jevtic, G., Mladenovic, M., Lusic, Z and Sokolovic, D. 2007. Morphological and production characteristics of Cariolan honey bee (*Apis mellifera carnica* Poll.) from different parts of Serbia. *Biotechnology in Animal Husbandry* 23(5-6): 609-617
- Kshirsagar, K. K. 1973. Comparative biometric studies on Indian honeybees. III. Preliminary observations on biometry of *Apis cerana* F. queen. *Indian Bee Journal* 35: 21–26

- Kunert K, Crailsheim K. Seasonal changes in carbohydrate, lipid and protein content in emerging worker honeybees and their mortality. *J Apic Res.* 1988;27: 13–21
- Kuliçi M, Kume K. Study on morphometric traits of the Albanian bees. *Albanian J Agric Sci.* 2014;(Special edition): 419–425.
- Kshirsagar KK (1981) Morphometric studies on *Apis cerana indica* worker. III Effect of geographical position on morphometric characters. *Indian Bee J* 43(1):1–5
- Mujumdar, S and Kshirsagar, K. K. 1986. Morphometric characterization of *Apis dorsata* Fabr. worker. *Indian Bee J* 48:25–29
- Makhmoor, H.D and Ahmad, H. 2001. Biometric studies on four species of honey bees in Jammu region, *India.* *Indian Bee J* 60:141–142.
- Mattu, V.K and Verma, L.R. 1984a. Comparative morphometric studies on the Indian honey bee of the North-West Himalayas 2 wings. *Journal of Apicultural Research* 23(1): 1 -10.
- Mattu, V.K and Verma, L.R. 1984b. Comparative morphometric studies on the Indian honey bee *Apis cerana indica* F., effect of seasonal variations. *Apidologie* 15: 63-74.
- Makkar, G. S., Chhuneja, P. K and Singh, J. 2020. Morphometric characterization of *Apis* species (Hymenoptera: Apoidea) Vegetos <https://doi.org/10.1007/s42535-020-00136-3>
- Meixner, M.D., Worobik, M., Wilde, J., Fuchs, S. and Koeniger, N. 2007. *Apis mellifera mellifera* in Eastern European morphometric variation. *Apidologie* 38(1): 1-7.
- Miguel I, Baylac M, Iriondo M, Manzano C, Garnery L, Estonba A. 2011. Both geometric morphometric and microsatellite data consistently support the differentiation of the *Apis mellifera* M evolutionary branch. *Apidologie*;42: 150–161.
- Mostajeran, M., Mohammad, A.E and Mohammad, R.B. 2006. Analysis of colony and morphological characters in honey bees *Apis mellifera meda*. *Pakistan Journal of Biological Sciences* 9(14): 2685-2688
- Narayanan, E. S., Sharma, P. L. and Phadke, K. G. 1960. Tongue length and number of hooks on the hind wings of *Apis indica* F. *Indian Bee Journal* 22:81-88.
- Narayanan, E. S., Sharma, P. L. and Phadke, K. G. 1961a. Studies on biometry of the Indian bees. III. Tongue length and number of hooks on the hind wings of *Apis indica* F. collected from Madras State. *Indian Bee Journal* 23:3–9.
- Narayanan, E. S., Sharma, P. L. and Phadke, K. G. 1961b. Studies on biometry of the Indian bees. IV. Tongue length and number of hooks on the hind wings of *Apis indica* F. collected from Uttar Pradesh. *Indian Bee Journal* 23: 69–74
- Oldroyd, B.P, Wongsiri, S. 2006. Asian honey bees: biology, conservation and human interactions. Harvard University Press, Harvard, p 360

- Roulston TH, Cane JH. Pollen nutritional content and digestibility for animals. *Plant Syst Evol.* 2000; 222: 187–209.
- Ruttner F. 1988. *Biogeography and Taxonomy of Honey bees*. Springer-Verlag .p 251
- Ruttner,F., Tassencourt, L and Louveaux,J. 1978. Biometrical statistical analysis of the geographic variability of *Apis mellifera* L. *Apidologie* 9(4): 363-381.
- Sakagami, S.F., Matsumura, T and Ito, K. 1980.*Apis laboriosa* in the Himalaya, the little known world largest honey bee (Hymenoptera:Apidea).*Insecta Matsumurana* 19:47-77
- Sharma,S.K. 1990. Biometric and developmental biology of *Apis mellifera* L. workers, M.Sc thesis submitted to Department of Entomology, Himachal Pradesh Krishi Vishvavidyalaya, Palampur India. pp. 1-56.
- Sheppard,W.S and Meixner, M.D. 2003. *Apis mellifera pomonella*, a new honey bees subspecies from Central Asia. *Apidologie* 34: 367-375.
- Szabo, T.I. 1990. Morphometric characteristics on *Apis cerana* from Sri Lanka. *Apidologie* 2: 505-509.
- Szentgyörgyi H, Czekońska K, Tofilski A. Influence of pollen deprivation on the fore wing asymmetry of honeybee workers and drones. *Apidologie.* 2016;47: 653–662.
- Tofilski, A. 2008. Using geometric morphometries and standard morphometry to discriminate three honey bee subspecies. *Apidologie* 39(5): 558-563.
- Verma ,L.R., Kafle, G.P., Sharma, A., Mattu V.K and Daly, H.V. 1990. Multivariate morphometries of the Indian honey bee in the North West Himalayan region. *Apidologie* 25: 203- 223.
- Zheng B, Wu Z, Xu B. The effects of dietary protein levels on the population growth, performance, and physiology of honeybee workers during early spring. *J Insect Sci.* 2014;14: 1–

**Table 1: Variability in dimensions of head of *Apis mellifera* L. at different altitudes in Kashmir region**

Parameters	Altitudes	Locations	Mean±S.E (mm)	Range	Over all mean	C.V	C.D (P=0.05)	P value
Head height	High	Narastan	3.27±0.01	3.21-3.35	<b>3.23±0.01</b>	1.70	0.0101	0.005** <0.01
		Laaam	3.21±0.04	3.21-3.34		6.81		
		Yarmukaaam	3.22±0.09	3.13-3.34		1.48		
	Mid	Damhal	3.18±0.05	3.12-3.23	<b>3.18±0.04</b>	0.82		
		Bonpora	3.18±0.04	3.12-3.23		0.70		
		Kalaroos	3.18±0.04	3.12-3.23		0.72		
	Low	Uri Bijihama	3.15±0.07	3.10-3.23	<b>3.14±0.03</b>	1.15		
		Anchidora	3.14±0.07	3.10-3.23		1.26		
		Dara	3.15±0.01	3.10-3.31		1.66		
Head width	High	Narastan	3.64±0.02	3.28-3.81	<b>3.67±0.20</b>	3.33	0.0101	0.023* <0.05
		Laaam	3.71±0.02	3.50-4.03		3.05		
		Yarmukaaam	3.68±0.01	3.50-4.00		2.39		
	Mid	Damhal	3.71±0.08	3.60-3.76	<b>3.70±0.03</b>	1.19		
		Bonpora	3.70±0.08	3.60-3.76		1.10		
		Kalaroos	3.70±0.09	3.60-3.76		1.25		
	Low	Uri Bijihama	3.62±0.01	3.45-3.78	<b>3.62±0.08</b>	2.68		
		Anchidora	3.61±0.01	3.45-3.76		2.58		
		Dara	3.64±0.02	3.45-3.91		3.46		

**Table 2: Variability in dimensions of proboscis, flagellum and antenna of *Apis mellifera* L. at different altitudes in Kashmir region**

Parameters	Altitudes	Locations	Mean±S.E (mm)	Range	Overall mean	C.V	C.D (P=0.05)	P value
<b>Proboscis length</b>	High altitude	Narastan	6.37±0.01	6.23-6.67	<b>6.35±0.06</b>	1.46	0.0101	0.001** <0.01
		Laaam	6.34±0.02	6.21-6.74		2.12		
		Yarmukaaam	6.35±0.02	6.23-6.76		1.93		
	Mid altitude	Damhal	6.27±0.08	6.24-6.45	<b>6.26±0.03</b>	0.71		
		Bonpora	6.26±0.06	6.21-6.34		0.52		
		Kalaroos	6.26±0.09	6.23-6.43		0.74		
	Low altitude	Uri Bijihama	6.22±0.07	6.20-6.34	<b>6.22±0.03</b>	0.56		
		Anchidora	6.23±0.07	6.21-6.34		0.57		
		Dara	6.23±0.07	6.32-6.43		0.76		
<b>Flagellum length</b>	High altitude	Narastan	3.81±0.02	3.67-3.87	<b>3.79±0.08</b>	1.66	0.0394	0.001** <0.01
		Laaam	3.78±0.01	3.71-3.89		1.55		
		Yarmukaaam	3.79±0.01	3.71-3.87		1.44		
	Mid altitude	Damhal	3.60±0.02	3.56-3.76	<b>3.60±0.01</b>	1.83		
		Bonpora	3.58±0.01	3.54-3.66		0.94		
		Kalaroos	3.64±0.03	3.55-3.87		2.71		
	Low altitude	Uri Bijihama	3.64±0.03	3.22-3.45	<b>3.42±0.01</b>	3.62		
		Anchidora	3.33±0.02	3.22-3.45		3.20		
		Dara	3.31±0.02	3.45-3.91		2.61		
<b>Antennal length</b>	High altitude	Narastan	6.11±0.01	6.01-6.12	<b>6.09±0.03</b>	0.52	0.0305	0.001** <0.01
		Laaam	6.09±0.01	6.01-6.12		0.52		
		Yarmukaaam	6.09±0.01	6.01-6.23		0.90		
	Mid altitude	Damhal	5.95±0.01	5.87-5.98	<b>5.94±0.05</b>	0.78		
		Bonpora	5.93±0.01	5.87-5.98		0.80		
		Kalaroos	5.94±0.01	5.87-5.99		0.80		
	Low altitude	Uri Bijihama	5.67±0.03	5.45-5.87	<b>6.09±0.03</b>	1.95		
		Anchidora	5.63±0.03	5.44-5.78		1.97		
		Dara	5.64±0.02	5.45-5.68		1.27		

**Table 3: Variability in dimensions of thorax and abdomen of *Apis mellifera* L. at different altitudes in Kashmir region**

Parameters	Altitudes	Locations	Mean±S.E (mm)	Range	Over all mean	C.V	C.D P=0.05	P value
Thorax length	High	Narastan	4.30±0.01	4.2-4.42	<b>4.28±0.08</b>	1.51	0.0143	0.000** <0.01
		Laaam	4.27±0.01	4.2-4.44		1.48		
		Yarmukaaam	4.28±0.01	4.2-4.56		1.80		
	Mid	Damhal	4.24±0.02	3.89-4.44	<b>4.23±0.03</b>	2.54		
		Bonpora	4.24±0.09	4.12-4.33		1.15		
		Kalaroos	4.23±0.02	3.98-4.44		2.47		
	Low	Uri Bijihama	4.16±0.05	4.14-4.23	<b>4.15±0.05</b>	0.69		
		Anchidora	4.15±0.04	4.14-4.23		0.57		
		Dara	4.16±0.06	4.12-4.3		0.80		
Abdomen length	High	Narastan	6.03±0.03	5.89-6.92	<b>6.02±0.03</b>	3.09	0.0431	0.001** <0.01
		Laaam	5.96±0.05	5.26-6.92		4.79		
		Yarmukaaam	6.08±0.04	5.89-6.8		3.75		
	Mid	Damhal	5.81±0.04	5.45-6.04	<b>5.88±0.04</b>	3.44		
		Bonpora	5.90±0.05	5.45-6.78		4.74		
		Kalaroos	5.95±0.04	5.67-6.98		3.86		
	Low	Uri Bijihama	5.58±0.02	5.45-5.98	<b>5.56±0.08</b>	2.53		
		Anchidora	5.55±0.02	5.45-5.98		2.25		
		Dara	5.57±0.02	5.45-5.98		2.29		

**Table 4: Variability in dimensions of fore wing and hind wing of *Apis mellifera* L. at different altitudes in Kashmir region**

Parameters	Altitudes	Locations	Mean±S.E (mm)	Range	over all mean	C.V	C.D (P=0.05)	P value
Fore wing length	High	Narastan	9.23±0.07	8.24-9.83	<b>9.28±0.02</b>	4.29	0.0203	0.003** <0.01
		Laaam	9.32±0.02	9.01-9.67		1.44		
		Yarmukaaam	9.29±0.07	8.24-9.91		4.27		
	Mid	Damhal	9.20±0.02	9.12-9.19	<b>9.21±0.01</b>	0.15		
		Bonpora	9.21±0.03	8.24-9.19		2.02		
		Kalaroos	9.21±0.01	8.78-9.34		0.98		
	Low	Uri Bijihama	9.18±0.06	9.12-9.23	<b>9.17±0.05</b>	0.36		
		Anchidora	9.17±0.08	9.10-9.25		0.44		
		Dara	9.17±0.06	9.23-9.21		0.35		
Forewing width	High	Narastan	3.03±0.02	2.90-3.50	<b>3.04±0.08</b>	3.36	0.0305	0.047* <0.05
		Laaam	3.06±0.02	2.99-3.50		4.36		
		Yarmukaaam	3.04±0.01	2.99-3.50		3.27		
	Mid	Damhal	2.97±0.05	2.55-3.09	<b>2.97±0.01</b>	2.75		
		Bonpora	2.96±0.28	2.34-3.09		4.74		
		Kalaroos	3.00±0.07	2.94-3.09		1.22		
	Low	Uri Bijihama	3.04±0.04	2.90-3.89	<b>3.00±0.02</b>	7.20		
		Anchidora	3.00±0.04	2.09-3.31		7.01		
		Dara	2.97±0.01	2.67-3.10		2.41		
Cubital index	High	Narastan	2.30±0.01	2.20-2.55	<b>2.29±0.03</b>	3.58	0.0380	0.001** <0.01
		Laaam	2.29±0.01	2.16-2.55		4.17		
		Yarmukaaam	2.29±0.01	2.20-2.55		3.52		
	Mid	Damhal	2.19±0.08	2.05-2.23	<b>2.19±0.03</b>	2.01		
		Bonpora	2.19±0.02	2.08-2.34		2.33		
		Kalaroos	2.20±0.06	2.21-2.25		1.50		
	Low	Uri Bijihama	2.55±0.08	2.42-2.55	<b>2.55±0.08</b>	1.71		
		Anchidora	2.55±0.01	2.08-2.55		3.95		
		Dara	2.55±0.02	2.21-2.55		4.70		
Hind wing length	High	Narastan	6.59±0.03	6.11-6.99	<b>6.55±0.01</b>	3.00	0.0635	0.001** <0.01
		Laaam	6.55±0.03	6.20-6.98		3.00		
		Yarmukaaam	6.53±0.01	6.33-6.78		1.95		
	Mid	Damhal	6.35±0.02	6.06-6.55	<b>6.35±0.05</b>	1.81		
		Bonpora	6.36±0.02	6.06-6.56		1.66		
		Kalaroos	6.35±0.02	6.06-6.55		1.88		
	Low	Uri Bijihama	6.19±0.06	6.01-6.23	<b>6.20±0.06</b>	0.52		
		Anchidora	6.21±0.09	6.01-6.24		0.75		
		Dara	6.21±0.09	6.01-6.24		0.78		
Hind wing width	High	Narastan	1.82±0.08	1.72-1.92	<b>1.81±0.06</b>	2.40	0.0176	0.001** <0.01
		Laaam	1.80±0.04	1.67-1.88		2.89		
		Yarmukaaam	1.82±0.01	1.56-1.98		4.80		
	Mid	Damhal	1.75±0.08	1.67-1.87	<b>1.75±0.03</b>	2.34		
		Bonpora	1.76±0.07	1.68-1.87		2.08		
		Kalaroos	1.75±0.01	1.56-1.98		4.03		
	Low	Uri Bijihama	1.73±0.01	1.56-1.89	<b>1.72±0.05</b>	3.52		
		Anchidora	1.71±0.02	1.34-1.87		7.79		
		Dara	1.72±0.01	1.56-1.73		3.03		
No. of Hamuli	High	Narastan	20.64±0.18	20-23	<b>20.66±0.02</b>	4.39	0.4016	0.001** <0.01
		Laaam	20.64±0.18	20-23		4.39		
		Yarmukaaam	20.72±0.18	20-23		4.51		
	Mid	Damhal	18.64±0.28	15-21	<b>18.86±0.2</b>	7.72		
		Bonpora	18.68±0.24	15-21		6.68		
		Kalaroos	19.28±0.23	15-21		6.08		
	Low	Uri Bijihama	23.44±0.19	21-24	<b>23.58±0.14</b>	7.72		
		Anchidora	23.44±0.15	22-24		3.27		
		Dara	23.88±0.06	23-24		1.38		

**Table 5: Variability in dimensions of hind leg (femur, tibia, metatarsus length and breadth) of *Apis mellifera* L. at different altitudes in Kashmir region**

Parameters	Altitudes	Locations	Mean±S.E (mm)	Range	Over all mean	C.V	C.D (P=0.05)	P value
Femur length	High	Narastan	2.55±0.09	2.50-2.67	<b>2.53±0.01</b>	1.92	0.0203	.001** <0.01
		Laaam	2.53±0.01	2.34-2.66		2.46		
		Yarmukaaam	2.51±0.01	2.34-2.66		2.76		
	Mid	Damhal	2.38±0.01	2.22-2.45	<b>2.41±0.05</b>	2.81		
		Bonpora	2.42±0.01	2.22-2.52		3.01		
		Kalaroos	2.43±0.01	2.22-2.56		3.10		
	Low	Uri Bijihama	2.33±0.08	2.21-2.45	<b>2.35±0.05</b>	1.91		
		Anchidora	2.38±0.04	2.21-3.35		8.7		
		Dara	2.34±0.01	2.21-2.45		2.16		
Tibia length	High	Narastan	2.93±0.06	2.87-2.98	<b>2.93±0.03</b>	1.03	0.0143	.001** <0.01
		Laaam	2.94±0.01	2.78-3.00		1.70		
		Yarmukaaam	2.93±0.07	2.87-2.99		1.26		
	Mid	Damhal	2.82±0.01	2.75-2.98	<b>2.81±0.05</b>	1.83		
		Bonpora	2.80±0.01	2.71-2.94		1.93		
		Kalaroos	2.81±0.08	2.75-2.89		1.47		
	Low	Uri Bijihama	2.86±0.01	2.76-2.94	<b>2.87±0.03</b>	2.15		
		Anchidora	2.87±0.01	2.70-2.99		2.80		
		Dara	2.88±0.01	2.76-2.98		1.74		
Metatarsus length	High	Narastan	1.96±0.02	1.95-1.98	<b>1.95±0.05</b>	0.52	0.0176	.001** <0.01
		Laaam	1.95±0.05	1.87-1.98		1.34		
		Yarmukaaam	1.94±0.05	1.86-1.98		1.51		
	Mid	Damhal	1.83±0.08	1.67-1.89	<b>1.83±0.01</b>	2.31		
		Bonpora	1.83±0.01	1.67-1.98		3.77		
		Kalaroos	1.82±0.08	1.67-1.88		2.37		
	Low	Uri Bijihama	1.78±0.09	1.67-1.89	<b>1.78±0.03</b>	2.52		
		Anchidora	1.78±0.08	1.74-1.98		2.23		
		Dara	1.79±0.08	1.74-1.88		2.30		
Metatarsus breadth	High	Narastan	1.13±0.07	1.1-1.23	<b>1.12±0.05</b>	3.20	0.0101	0.001** <0.01
		Laaam	1.12±0.08	1.05-1.23		3.64		
		Yarmukaaam	1.11±0.04	1.08-1.23		2.38		
	Mid	Damhal	1.07±0.04	1.2-1.09	<b>1.07±0.03</b>	2.01		
		Bonpora	1.07±0.03	1.04-1.10		1.41		
		Kalaroos	1.08±0.03	1.04-1.12		1.51		
	Low	Uri Bijihama	1.04±0.03	1.02-1.09	<b>1.04±0.00</b>	1.66		
		Anchidora	1.04±0.03	1.02-1.09		1.70		
		Dara	1.03±0.02	1.02-1.08		1.29		

S.E = Standard error, C.V= Coefficient of variation, C.D = Critical Difference, P=threshold of Significance, P≤0.01(\*\*) difference highly significant, P≤ 0.05(\*) difference significant, P≥0.05 ns (Non-Significant)