

Original Research Article

THE USE OF CROP RESIDUES AND HIGH ACTIVITY CLAYS FOR THE MANAGEMENT OF SANDY SOILS FOR SUSTAINABLE RICE PRODUCTION IN SOKOTO, NORTH-WESTERN NIGERIA

Comment [tb1]: In title section the mentioned words 'sustainable rice production' are not properly justified throughout the manuscript. Include relevant literature in introduction or literature review section and correlating it with the factors in results and discussion section.

Abstract

A field experiment was carried out to assess the effects of various crop residues (rice husk, rice straw, groundnut husk, and millet husk as well as high activity clay (HAC) on some soil properties, growth and yield parameters of rice (*Oryza sativa* L). Sixteen treatment combinations, replicated thrice, were applied at the rates of 5tha⁻¹ and 10tha⁻¹ each for rice husk, rice straw, millet husk, groundnut husks and high activity clay. Results obtained from the field trial as presented in this paper, showed significant effect of some of the treatments on soil chemical parameters especially organic Carbon, available P and exchangeable K. Similarly, soil physical parameters such as texture were also significantly influenced by the treatments. The texture of the soil, which was initially loamy sandy before the incorporation of the amendments, was altered to new textural classes in all the treatments after the experiment. The treatments had significant effects on all the growth parameters except for plant height at four (4) weeks after transplanting. The treatments also had significant effects on all the yield parameters of the test crop (rice).

Key Words: Sandy soils, Crop Residues and High activity clay.

Comment [tb2]: Try to avoid a lengthy terminology adjoining with the words 'and' or 'but'. It will be better to use single word like Sandy soils, Crop Residues, High actively clay (separated by commas).

Introduction

Sandy soils are characterized by possession of less than 18% clay and more than 68% sand in the first 100 cm of the column (Brady and Weil, 1999). These soils are often considered as soils with physical properties that are easy to define such as weak structure or no structure, poor water retention properties, low nutrient retention highly sensitive to compaction with many adverse consequences (Bell *et al.*, 2004; Wada, 1996 and Ishida *et al.*, 1993).

Crop residues are plant materials remaining after harvest including leaves, stalks and roots. Their presences on the soil surface decrease erosion, increases organic matter, improve soil quality, increase water holding capacity and infiltration (Lal, 2014). High activity clays have a high cation exchange capacity due to their large surface area. These soils have great capacity

Comment [tb3]: Introduction section is too short. Maintain the proper conceptual development, review of literature and justification of the study in this section.

to retain and supply large quantities of nutrients such as calcium, magnesium and potassium. They generate high CEC under all soil conditions regardless of soil pH (Masters, 2010).

MATERIALS AND METHODS

Experimental site and materials used

The experiment was carried out at the Fadama Teaching and Research Farm (Lat. 13^o.092250N' Long. 5^o.208746E, altitude 265m a.s.l.), Faculty of Agriculture, Usmanu Danfodiyo University Sokoto. The soil of the study site belongs to the order Entisols in the USDA Soil Taxonomy system. The soils of the farm are mostly loamy sand in texture. The land was divided into 48 plots each measuring 3m X 3m. Three replicates were used with 1m as spacing between and within rows and laid out in a Randomized Complete Block Design (RCBD) with 16 treatment combinations. The gross plot was 9m² (3m x 3m) with a net plot area of 7.56m² (2.75 X 2.75) and total experimented area of 591.25m² (27.5 X 21.5).

The variety of rice used was Faro 44, which was obtained from National Cereal Research Institute (NCRI) Outstation, Birnin Kebbi. The high activity clay was obtained from the Inland Valley and flood plains along Illela Road, Sokoto State.

Experimental Layout

The total number of plots was 48 with each measuring 9m² in size and labeled for the various treatments combination i.e., mixture of crop residues and high activity clay at difference rates. The treatments consisted of a factorial combination of two levels of high activity clay (HAC); 10tha⁻¹ and 15tha⁻¹ each of which was combined with two level each, 5 tha⁻¹ and 10 tha⁻¹, of groundnut shell, rice straw, rice husk and millet husk respectively giving a total of 16 treatment combinations laid out in a Randomized Complete Block Design (RCBD) replicated three times.

Soil analysis

The soil samples were analysis for physical and chemical properties before and after the experiment. Organic carbon was determined by Walkley and Black (1934) method. Total nitrogen was determined by the Microkhjeldal method. Available phosphorus extraction was determined calorimetrically using spectrophotometer as described by Olsen and Sommers (1982). Textural triangle was used for soil texture determinations. The cation exchange

Comment [tb4]: .The total number of cited words have been exceeded '40'. So, use of 'Block citation style' is required. Link: <https://research.wou.edu/apa/apa-block-quote#-:text=APA%20Style%20Guide%3A%20Block%20Quote&text=Block%20quotes%20are%20used%20for.quote%20may%20be%20single%20spaced.>

Comment [tb5]:

Comment [tb6]: Mention the justification of selection of the farm.

Comment [tb7]: The degree symbols are not properly adjusted with the value. Adjust the symbols.

Comment [tb8]: Same as the previous.

Comment [tb9]: Use the term Mean Sea Level (M.S.L.) instead of Average Sea Level (A.S.L.).

Comment [tb10]: Use the full forms. Instead of Lat. it will be Latitude and Long. – Longitude. Convert the values of the coordinates into degree and minutes.

Comment [tb11]: Use the full form of the abbreviations when it is used first. Then use the abbreviations subsequently.

Comment [tb12]: Reference is not same as cited.

Comment [tb13]: Include reference.

analysis (CEC) of the soil was determined by using pH 7.0 buffer solution of calcium ammonium acetate while EDTA titration was used to measure the Ca^{2+} , Mg^{2+} and K^+ . Soil pH was measured in both water (1:1) and 0.01m CaCl_2 (1:2.5). Particle size distribution was carried out using the hydrometer method.

Comment [tb14]: Use the full form of the abbreviations when it is used first. Then use the abbreviations subsequently.

TREATMENT COMBINATIONS	ACRONYMS
------------------------	----------

Groundnut shell 5tha ⁻¹ + HAC, 10tha ⁻¹	G ₁ HAC ₁
Groundnut shell 10 tha ⁻¹ + HAC 15tha ⁻¹	G ₂ HAC ₂
Rice straw 5 tha ⁻¹ + HAC 10tha ⁻¹	RS ₁ HAC ₁
Rice straw 10tha ⁻¹ + HAC 15tha ⁻¹	RS ₂ HAC ₂
Rice husk 5tha ⁻¹ + HAC 10tha ⁻¹	RH ₁ HAC ₁
Rice husk 10tha ⁻¹ + HAC 15tha ⁻¹	RH ₂ HAC ₂
Millet husk 10tha ⁻¹ + HAC 10tha ⁻¹	MH ₁ HAC ₁
Millet husk 10tha ⁻¹ + HAC 15tha ⁻¹	MH ₂ HAC ₂
Groundnut shell 5tha ⁻¹ + HAC 15tha ⁻¹	G ₁ HAC ₂
Groundnut shell 10 tha ⁻¹ + HAC 10tha ⁻¹	G ₂ HAC ₁
Rice straw 5tha ⁻¹ HAC + 15tha ⁻¹	RS ₁ HAC ₂
Rice straw 10tha ⁻¹ + HAC 10tha ⁻¹	RS ₂ HAC ₁
Rice husk 5tha ⁻¹ + HAC 15tha ⁻¹	RH ₁ HAC ₂
Rice husk 10 tha ⁻¹ + HAC 10tha ⁻¹	RH ₂ HAC ₁
Millet husk 5tha ⁻¹ + HAC 15tha ⁻¹	MH ₁ HAC ₂
Millet husk 10tha ⁻¹ + HAC 10tha ⁻¹	MH ₂ HAC ₁

Comment [tb15]: Mention the justification that why these treatment combinations have been used in experimental design and soil analysis.

Comment [tb16]: Bracket is incomplete.

Comment [tb17]: Mention the source(s).

RESULTS AND DISCUSSION

Comment [tb18]: If possible, try to separate the section as Results, Discussion. In Results section mention the analytical outputs as mentioned in the tables; and in Discussion section properly discuss the results.

The results of soil analysis before the experiment are presented in table 1. The particle size analysis revealed that the soil is loamy sand in texture, slightly acidic and low in organic matter, CEC and available phosphorus. The results indicate that; pH values based on the ratings by Chudeet *al.* (2011), ranged from acidic with mean values of 5.2 (1:1 soil water ratio) and pH measured in CaCl₂ (1:1 soil solution ratio) was 4.5. Exchangeable cations (Ca, Mg, K, Na) were low.

Comment [tb19]: Use the full form of the abbreviations when it is used first. Then use the abbreviations subsequently.

The treatments had significant ($P < 0.05$) on soil pH both in water and CaCl₂. The initial pH recorded before the experiment was 5.3 and 4.5 for pH (H₂O) and pH (CaCl₂) respectively. (Table 3) This shows the pH was strongly acidic (Chudeet *al.*, 2011). The pH value recorded on MH₂HAC₂ (7.3) which is slightly alkaline for pH (H₂O) and 7.0 neutral for pH (CaCl₂). The significant increase in the soil pH on all treatments might be because of the release of basic cations by the crop residues (Vanlauwe *et al.*, 1994).

The results from table 3 shows that treatments significantly ($P < 0.05$) influenced soil organic matter. Treatment G₂HAC₂ recorded the highest value. Ashrif and Thornton (1965) found out that the incorporation of groundnut residue will provide organic matter and other plant nutrients in the soil. Groundnut being a legume has the ability to fix atmospheric nitrogen which is directly related to increased organic matter in the soil.

Results in table 3 showed that the treatments had significant effect on nitrogen. Treatment G₂HAC₁ recorded the highest total nitrogen. Anderson and Peterson (1973) found out that the groundnut residues can be incorporated to provide nitrogen for non-leguminous crops such as rice. The water holding capacity of the clay might have aided the soil in conserving and releasing nutrients (Ashrif and Thornton, 1965).

All the treatments (Table 3) had significant effect on CEC. Treatment RH₁HAC₂ had the highest CEC (11.83 Cmolkg⁻¹) Singh *et al.* (1980) reported that application of rice residues and manure resulted in increased CEC.

The treatments had significant effect on available phosphorus. The initial value of available P before the experiment (4.6mgkg⁻¹) was low (Esu, 1991). Treatment RH₁HAC₂ recorded the highest value of Available Phosphorus. Brady and Weil (1999) reported that, clays that possess greater anion exchange capacity (due to positive surface charges) have a greater affinity for phosphorus ions.

The results in Table 3 indicated that all the treatments had significant effects on exchangeable calcium. Treatment RH₁HAC₂ gave the highest value (3.6cmolk⁻¹). Generally there was an increase in exchangeable calcium in all treatments. This is supported by the findings of Ahmad (2015).

Comment [tb20]: Grammatical and punctuational checkup is required for the whole document.

All the treatments had significant effects on exchangeable Magnesium. Treatment (RH₂HAC₂) recorded the highest value. It was observed that exchangeable magnesium increased with increase in clay content (Roychand and Marschner, 2013).

The treatments had significant effect on exchangeable potassium. The highest value was recorded by treatment RH₂HAC₂. Kaur and Benepal (2006) reported an increase in available K and water soluble K as a result of incorporation of rice residue and farm yard manure.

The exchangeable sodium was significantly ($P < 0.05$) affected by the treatments. Treatment RS₂HAC₁ recorded the highest value (0.26cmolk⁻¹). This value is low according to the rating of Esu (1991).

Electrical conductivity (EC) was significantly affected by the treatments. The EC value of the soil before the experiment was 0.27ds/m. Treatment RS₂HAC₁ recorded the highest value of EC (1.7 dS/m). The soils of the study area were generally non-saline, based on the rating of Landon (1991) who reported that soils of EC levels of 0-2dS/m are salt free soils.

The data on textural classes of the treatments is presented in table 2. The initial texture of the soil before the experiment (Table 1) was Loamy sand. As can be seen from Table 2, all the textural classes of the 16 treatments were altered. All treatments where 10tha⁻¹ high activity clay was incorporated had their textural class altered to Loamy while all treatments where 15tha⁻¹ high activity clay was incorporated changed to clay loam. This is irrespective of the type of crop residue that was combined with the high activity clay. This confirmed the assertion that soil texture is a fixed property that can only be changed by the addition of one soil separate or the other (Hallet *et al.*, 2010).

The tiller count of rice in Table4 was significantly affected by the treatments. Treatment RH₂HAC₂ recorded the highest tiller per stand. The result obtained may be attributed to an improved soil moisture retention condition as a result of higher rate of rice husk and clay. This assertion was earlier stated by Muktaret *et al.* (2015).The treatments did not significantly ($P > 0.05$) affect plant height at 4 weeks after transplanting (WAT). However, plant height was significantly ($P < 0.05$) affected at 8 and 12 weeks after transplanting.

Treatment RH₁HAC₂ recorded the highest value both at 8 and 12 weeks after transplanting. This is in agreement with the work of Lal (1997) who found that application of rice husk combined residues produced the highest plant height.

The effect of crop residue and HAC on the number of spikes in rice is presented in Table 4. The treatments had significant (P<0.05) effect on the number of spikelet per spike. Treatment RH₂HAC₂ recorded the highest value with a mean of 15.00. This was attested by Ogboghodo *et al.* (2018), whose results showed that nutrient uptake increased as the rate of application of rice husk and clay increased. The number of grains per spike (Table 4) was significantly (P<0.05) influenced by the treatments. Treatment RH₂HAC₂ recorded the highest number of grains per spike. This was earlier confirmed by Hall *et al.* (2010) who found that the addition of 15 t/ha clay to sandy top soil increased number of grains per spike in rice. The treatment had significant effect on grain yield. RH₁HAC₂ recorded the highest grain yield. This may be attributed to the high quality of N, K, Mg and Ca as attested by Muktareta *et al.* (2015).

Comment [tb21]: Clarify it based on reference.

Roychand and Marschner (2013) also found that the amending sandy soil with rice husk and clay rich subsoil could increase grain yield in wheat. The treatments had significant (P<0.05) effect on straw yield of rice. Gupta (2002) reported high straw yield in rice where rice husk amended was combined with 8 t/ha clay. The treatments had significant (P<0.05) effect on one thousand (1000) grain weight. Treatment RH₂HAC₂ recorded the highest value of one thousand (1,000) grain weight. This was supported by findings of Sanderman and Baldock (2010) who recorded a similar result. They reported that the high magnesium (0.36 cmol kg⁻¹) and Calcium (0.61 cmol kg⁻¹) as being responsible for high yield in one thousand (1000) grain weight. Schweizer *et al.* (1999) reported increased soil organic carbon retention and a high one thousand grain weight where rice residues amendments were used in rice cultivation. This finding is in confirmity with Oghoghodo *et al.* (1995) who found out that the incorporation of rice crop residues as being superior over removal or burning of crop residue practices possibly due to improved physical, chemical and biological properties of soils due to increase organic carbon content in the soil.

Comment [tb22]: Include reference.

Comment [tb23]: Correct the word.

CONCLUSION

Comment [tb24]: Elaborate the conclusion.

The results of the study showed that all the treatments had a significant effect on chemical and physical parameters of the soil. Also most of the chemical and physical parameters such as organic carbon, phosphorus, potassium EC, CEC, pH increased at the end of the

experiment. However sodium level in the soil decreased. Also, all the textures of the soils before the experiment (Loamy sand) were completely altered in the 16 treatments. This was a result of addition of various rates of clay to the soil. This was irrespective of the quantity of crop residue applied. The treatments had significant effects on the plant height (8 and 12 weeks after transplanting), number of tillers, number of spikelets per spike, number of grains per spikelet, straw yield grain yield and one thousand 1000 grain. Treatment RH₂HAC₂ proved to be the best among the 16 treatments.

UNDER PEER REVIEW

Table 1: Physical and chemical properties of the soil before the experiment

Soil Property	
Sand (%)	70
Silt (%)	22
Clay (%)	8
Textural class	Loamy sand
pH (H ₂ O) 1:1	5.3
pH (CaCl ₂) 1:1	4.5
EC(dS/m)	0.27
Org.matter (g kg ⁻¹)	12.7
Total N (g kg ⁻¹)	0.95
Org.C (g kg ⁻¹)	5.02
Av. P (mg kg ⁻¹)	4.6
CEC (Cmol kg ⁻¹)	9.2
ExchCa (Cmol kg ⁻¹)	1.55
Exch Mg (Cmol kg ⁻¹)	1.36
Exch K (Cmol kg ⁻¹)	0.80
Exch Na (Cmol kg ⁻¹)	1.6

Comment [tb25]: Mention the source(s).

Comment [tb26]: Include the tables in Appendix section.

UNDER PEER REVIEW

Table 2: Particle Size Distribution at the end of the Experiment

Comment [tb27]: Same.

Treatment	% Sand	% Silt	% Clay	% Textural Class
G ₁ HAC ₁	50	30	20	Loam
G ₁ HAC ₂	36.4	33.4	30.2	Clay Loam
G ₂ HAC ₁	49	31	20	Loam
G ₂ HAC ₂	40.4	24.4	35.3	Clay Loam
MH ₁ HAC ₁	49.4	31.1	20.2	Loam
MH ₁ HAC ₂	42.4	22.4	35.2	Clay Loam
MH ₂ HAC ₁	44.3	35.3	20.4	Loam
MH ₂ HAC ₂	36.4	33.3	30.3	Clay Loam
RH ₁ HAC ₁	50	29	21	Loam
RH ₁ HAC ₂	30.2	39.4	30.4	Clay Loam
RH ₂ HAC ₁	49.4	31.1	20.2	Loam
RH ₂ HAC ₂	39	31	30	Clay Loam
RS ₁ HAC ₁	51	29	20	Loam
RS ₁ HAC ₂	24	40	36	Clay Loam
RS ₂ HAC ₁	50	30	20	Loam
RS ₂ HAC ₁	32.4	25.3	33.3	Clay Loam

Table 3: Effect of crop residues and high activity clay on soil physical and chemical properties of the soil

Comment [tb28]: Same.

TRT	pH (H ₂ O 1:1)	pH (CaCl ₂ 1:1)	CEC (Cmol kg ⁻¹)	EC (dS/m)	Ca (cmol kg ⁻¹)	Mg (cmol kg ⁻¹)	K (cmol kg ⁻¹)	Na (cmol kg ⁻¹)	O.C g kg ⁻¹	O.M g kg ⁻¹	N (g kg ⁻¹)	Avail P (mg/kg ⁻¹)
RH ₂ HC ₂	6.933 ^{hi}	6.867 ^{ab}	11.83 ^a	0.5733 ^h	2.143 ^c	12.200 ^a	0.9700 ^c	0.1160 ^e	24.10 ^h	68.70 ^h	0.6667 ^d	7.633 ^a
RH ₁ HC ₂	6.833	5.700 ^f	11.80 ^a	1.1867 ^{bc}	3.633 ^a	4.567 ^b	0.5500 ^g	0.1740 ^e	40.33 ^a	41.93 ⁿ	0.6300 ^{de}	8.000 ^a
RS ₂ HC ₂	6.933 ^{hi}	6.867 ^{ab}	11.07 ^b	0.5733 ^h	2.143 ^c	2.333 ^j	0.5600 ^g	0.2167 ^b	26.67 ^g	46.13 ^l	0.6667 ^d	5.567 ^{cde}
RH ₂ HC ₁	6.967 ^{hi}	6.867 ^{ab}	10.77 ^c	0.9367 ^e	2.210 ^c	3.667 ⁱ	0.8133 ^{de}	0.1737 ^c	33.13 ^e	47.37 ^k	0.6067 ^e	6.667 ^b
G ₂ HC ₂	7.267 ^a	6.967 ^a	10.27 ^d	0.8300 ^f	2.400 ^{bc}	3.727 ^{ef}	0.7500 ^e	0.1343 ^d	39.20 ^{ab}	69.20 ^a	0.7333 ^c	4.800 ^{ef}
G ₂ HC ₁	7.153 ^{bcd}	6.833 ^{ab}	10.22 ^d	1.0267 ^d	2.133 ^c	4.233 ^c	0.8333 ^d	0.1313 ^d	36.23 ^d	62.77 ^e	0.9667 ^a	51.133 ^{def}
RS ₂ HC ₁	7.167 ^{abcd}	6.867 ^{ab}	10.17 ^d	0.5667 ^h	2.117 ^c	2.860 ^h	0.8133 ^{de}	0.2673 ^a	27.27 ^g	47.37 ^k	0.7667 ^c	5.433 ^{ef}
G ₁ HC ₂	7.137 ^{cd}	6.433 ^{cde}	9.97 ^e	0.9033 ^e	2.627 ^b	3.700 ^f	1.1000 ^b	0.1723 ^c	38.47 ^{bc}	67.37 ^c	0.8433 ^c	47.33 ^{ef}
RH ₁ HC ₁	6.867 ^{ij}	6.067 ^{ef}	9.83 ^f	0.7133 ^g	2.227 ^c	3.830 ^{de}	0.5500 ^g	0.1740 ^c	30.20 ^f	52.23 ^h	0.6767 ^d	6.667 ^b
G ₁ HC ₁	7.067 ^{def}	6.533 ^{bcd}	9.77 ^f	1.1333 ^c	2.133 ^c	3.500 ^g	0.6600 ^f	0.0863 ^f	35.80 ^d	63.63 ^d	0.8467 ^c	4.867 ^{ef}
RS ₁ HC ₁	7.000 ^{efgh}	6.867 ^{ab}	9.71 ^f	1.6800 ^a	3.470 ^a	3.750 ^{def}	0.5500 ^g	1737 ^c	27.87 ^g	48.13 ^j	0.7667 ^c	5.267 ^{def}
RS ₁ HC ₂	7.067 ^{delg}	6.800 ^{abc}	9.71 ^f	0.3200 ⁱ	3.633 ^a	2.610 ^j	0.6533 ^f	0.2647 ^a	24.10 ^h	41.93 ⁿ	0.7367 ^c	5.133 ^{def}
MH ₂ HC ₂	7.233 ^{abc}	7.000 ^a	9.57 ^g	0.5467 ^h	2.217 ^c	3.853 ^d	0.5433 ^g	0.1743 ^c	26.70 ^g	43.13 ^m	0.6400 ^{de}	5.733 ^{cd}
MH ₁ HC ₁	7.067 ^{def}	6.900 ^{ab}	9.24 ^h	1.1733 ^c	2.133 ^c	2.667 ⁱ	0.5133 ^g	0.1747 ^c	37.67 ^c	68.50 ^b	0.7333 ^c	6.233 ^{bc}
MH ₂ HC ₁	7.253 ^{ab}	7.033 ^a	9.17 ^h	0.6800 ^g	2.243 ^c	3.683 ^f	0.5667 ^g	0.1750 ^c	29.90 ^f	51.53 ^j	0.6733 ^d	5.700 ^{cd}
MH ₁ HC ₂	7.077 ^{de}	6.333 ^{de}	8.21 ⁱ	1.2433 ^b	2.233 ^c	2.583 ^j	0.6400 ^f	0.1743 ^c	34.27 ^e	59.83 ^f	0.6400 ^{de}	6.200 ^{bc}

P	0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	0.001	<0.001	<0.001	<0.001	<0.001
S. E	0.03240	0.0324>	0.0458	0.03073	0.1083	0.03985	0.02243	0.001113	0.435	0.001	0.01555	0.2165

Means followed by the same letter (s) within the same column are statistically the same at 5% level of probability using Duncan's New Multiple Range Test (DNMRT).

Comment [tb29]: Clarify the footnote style.

Comment [tb30]: Need proper citation and reference.

UNDER PEER REVIEW

KEY:**TREATMENT COMBINATIONS****ACRONYMS)**

Groundnut shell 5tha ⁻¹ + HAC, 10tha ⁻¹	G ₁ HAC ₁
Groundnut shell 10 tha ⁻¹ + HAC 15tha ⁻¹	G ₂ HAC ₂
Rice straw 5 tha ⁻¹ + HAC 10tha ⁻¹	RS ₁ HAC ₁
Rice straw 10tha ⁻¹ + HAC 15tha ⁻¹	RS ₂ HAC ₂
Rice husk 5tha ⁻¹ + HAC 10tha ⁻¹	RH ₁ HAC ₁
Rice husk 10tha ⁻¹ + HAC 15tha ⁻¹	RH ₂ HAC ₂
Millet husk 10tha ⁻¹ + HAC 10tha ⁻¹	MH ₁ HAC ₁
Millet husk 10tha ⁻¹ + HAC 15tha ⁻¹	MH ₂ HAC ₂
Groundnut shell 5tha ⁻¹ + HAC 15tha ⁻¹	G ₁ HAC ₂
Groundnut shell 10 tha ⁻¹ + HAC 10tha ⁻¹	G ₂ HAC ₁
Rice straw 5tha ⁻¹ HAC + 15tha ⁻¹	RS ₁ HAC ₂
Rice straw 10tha ⁻¹ + HAC 10tha ⁻¹	RS ₂ HAC ₁
Rice husk 5tha ⁻¹ + HAC 15tha ⁻¹	RH ₁ HAC ₂
Rice husk 10 tha ⁻¹ + HAC 10tha ⁻¹	RH ₂ HAC ₁
Millet husk 5tha ⁻¹ + HAC 15tha ⁻¹	MH ₁ HAC ₂
Millet husk 10tha ⁻¹ + HAC 10tha ⁻¹	MH ₂ HAC ₁

Comment [tb31]: Exclude this table here as it is previously included.

UNDER PEER REVIEW

Table 4: Effect of crop residues and high activity clay on growth parameters of rice in

Comment [tb32]: Data/Table source?

Sokoto

Treatment	Tiller Number (M ²)	Plant height (cm) 4WATP	Plant height (cm) 8 WATP	Plant height (cm) 12 WATP
G ₁ HAC ₁	39.67 ^{bcde}	31.00 ^a	58.67 ^{cd}	70.33 ^{gh}
G ₂ HAC ₁	39.33 ^{bcde}	32.33 ^a	62.33 ^{bcd}	70.67 ^g
RS ₁ HAC ₁	38.33 ^{bcde}	34.00 ^a	64.67 ^{abcd}	82.33 ^{ef}
RS ₂ HAC ₁	37.00 ^{cde}	33.67 ^a	67.33 ^{abc}	87.00 ^{de}
RH ₁ HAC ₁	31.33 ^e	31.67 ^a	70.33 ^{ab}	94.00 ^{bcd}
RH ₂ HAC ₁	41.00 ^{abcde}	32.33 ^a	71.00 ^{ab}	98 ^{abc}
MH ₁ HAC ₁	35.67 ^{cde}	33.33 ^a	55.67 ^d	68.67 ^{gh}
MH ₂ HAC ₁	40.00 ^{bcde}	31.67 ^a	59.00 ^{cd}	71.33 ^{gh}
G ₁ HAC ₂	40.67 ^{abcde}	32.33 ^a	56.33 ^{cd}	69.67 ^{gh}
G ₂ HAC ₂	39.67 ^{bcde}	32.33 ^a	56.33 ^{cd}	73.00 ^{gh}
RS ₁ HAC ₂	32.67 ^{de}	32.67 ^a	56.33 ^{cd}	89.00 ^{de}
RS ₂ HAC ₂	34.67 ^{cde}	34.33 ^a	62.33 ^{bcd}	91.00 ^{cd}
RH ₁ HAC ₂	48.33 ^{ab}	33.67 ^a	73.67 ^a	99.00 ^{ab}
RH ₂ HAC ₂	50.67 ^a	33.67 ^a	71.00 ^{ab}	102.67 ^a
MH ₁ HAC ₂	42.33 ^{abcd}	32.33 ^a	56.00 ^{cd}	66.33 ^h
MH ₂ HAC ₂	43.33 ^{abc}	32.00 ^a	58.67 ^{cd}	75.67 ^{fg}
Significance	*	NS	*	*
SE	3.101	2.231	3.351	2.46
P – Value	0.010	0.832	0.002	0.001

Means followed by the same letter (s) within the same column are statistically the same at 5% level of probability using Dunca's New Multiple Range Test (DNMRT).

* = Significant at 5% level of probability

NS = Not significant at 5% level probability

Comment [tb33]: Need proper citation and reference.

UNDER PEER REVIEW




Table 5: Effect of crop residues and high activity clay on yield parameters of rice in

Comment [tb34]: Data/Table source?

Sokoto

Treatment	Number of Spikes	Number of grains perspike	Grain yield Kg ha^{-1}	Straw yield Kg ha^{-1}	One thousand grain weight (g)
G ₁ HAC ₁	7.667 ^{def}	9.00 ^{de}	3.883 ^{cd}	8.623 ^{cd}	23.07 ^{efg}
G ₂ HAC ₁	6.33 ^f	11.00 ^{bcde}	3.330 ^{cd}	11.330 ^{bc}	23.73 ^{ef}
RS ₁ HAC ₁	6.667 ^{ef}	9.33 ^{cde}	4.700 ^{cd}	8.440 ^{cd}	22.80 ^{efg}
RS ₂ HAC ₁	9.00 ^{bcde}	8.33 ^e	4.623 ^{cd}	8.293 ^{ed}	27.33 ^{cd}
RH ₁ HAC ₁	8.667 ^{bcd}	8.67 ^e	8.143 ^b	10.923 ^{bc}	30.50 ^{bc}
RH ₂ HAC ₁	9.667 ^{bcd}	12.33 ^{abcd}	8.477 ^b	13.367 ^b	32.50 ^b
MH ₁ HAC ₁	6.667 ^f	9.33 ^{cde}	2.623 ^{cd}	7.997 ^{cd}	19.87 ^g
MH ₂ HAC ₁	8.33 ^{cdef}	11.00 ^{bcde}	2.070 ^d	7.030 ^d	20.83 ^{fg}
G ₁ HAC ₂	9.333 ^{bcd}	10.00 ^{cde}	4.147 ^{de}	6.923 ^d	23.30 ^{efg}
G ₂ HAC ₂	9.333 ^{bcd}	12.00 ^{abcd}	4.183 ^{cd}	6.477 ^d	25.43 ^{de}
RS ₁ HAC ₂	10.333 ^{bc}	8.33 ^e	3.810 ^{cd}	8.589 ^{cd}	27.43 ^{cd}
RS ₂ HAC ₂	11.000 ^b	10.33 ^{bcde}	4.993 ^c	8.107 ^{cd}	28.53 ^{cd}
RH ₁ HAC ₂	13.667 ^a	13.33 ^{ab}	9.587 ^{ab}	13.513 ^b	30.67 ^{bc}
RH ₂ HAC ₂	15.00 ^a	15.00 ^a	11.683 ^a	17.773 ^a	38.17 ^a
MH ₁ HAC ₂	9.667 ^{bcd}	11.00 ^{bcde}	3.290 ^{cd}	8.067 ^{cd}	21.40 ^{fg}
MH ₂ HAC ₂	10.000 ^{bcd}	11.00 ^{bcde}	2.827 ^{cd}	7.140 ^d	21.72 ^{fg}
Significance	*	*	*	*	*
SE	0.726	0.982	0.798	1.057	1.130
P - Value	0.001	0.001	0.001	0.001	0.001

Means followed by the same letter (s) within the same column are statistically the same at 5% level of probability using Dunca's New Multiple Range Test (DNMRT).

* = Significant at 5% level of probability


Comment [tb35]: Need proper citation and reference.

| NS = Not significant at 5% level probability

Comment [tb36]: Clarify the footnote style.

UNDER PEER REVIEW

UNDER PEER REVIEW



REFERENCES

- Ahmad, N. (2015). *Fertilizer use efficiency in Rice*. Paper presented. Paper presented on rice research and production, 29th – 31st March, 1988Islamabed, Pakistan.
- Anderson, F.N. and Peterson, G.A. (1973).Effect of continuous corn (*Zeamays L.*) manuring and N fertilization on yield and protein content.*Agronomy Journal* 65:697-700
- Ashrif, M.I., and Thornton, I. (1965).Effect on grass mulch on groundnut in Gambia.*Journal of Agriculture*, 1:145-152.
- Bell, R.W., Dell, B. and Huang, L. (2004).Importance of micro nutrients in crop production. In: International Fertilizer Association symposium on Micro-nutrients Delhi, 23-25 Feb, 2004. Published at <http://www.fertilizer>.
- Brady, N. and Weil, R.R. (1999).*The nature and properties of soils*, 12th Edition. Prentice Hall Incorporation, London, 881 pp.
- Bray, R.H. and Kurtz, L.T. (1945) Determination of Total Organic and Available Forms of Phosphorus in Soils. *Soil Science*, 59, 39-45.
- Chude, V.O., Jayeoba, O.J. and Berdin, F. (2011). A soil fertility map of Nigeria showing different P,N,Zn, B and K classes of the top soil. National Programme for Food Security, AdekumboAdemolacresent, Wusez, Abuja, Nigeria. 8pp.
- Esu, I.E. (1991). Detailed soil survey of NIHORT Farm at Bunkure, Kano State, Nigeria. Institute for Agricultural Research, Ahmad Bello University, Zaria, Nigeria. 72 pp.
- Gupta, R. (2002). Study on agroclimatic condition and productivity pattern of sugarcane in India. *Sugar Technology*, 6 (3): 142-149.
- Hall, D.J.M., Jones, H.R.H., Crabtree, W.L., and Daniels, T.L. (2010). Claying and dip ripping can increase crop yields and profits on water repellent sands with marginal fertility in Southern Western Australia. *Australian Journal of Soil Research*, 48:178-187.
- Ishida, H., Chairaj, P., Morakul, P, and wongwiwachai, C. (1993). Techniques to improve productivity of sandy upland soil in Northeast Thailand. ADRC technical paper No. 10.(Agricultural Development Research Centre) 45p.

Comment [tb37]: This is not included in citation.

- Kaur, N. and Benipal, D.S. (2006).Effect of crop residue and farmyard manure on K forms in soils of long-term fertility experiment. *Indian Journal of Crop Science*, 1(2): 161-164.
- Lal, R. (1997). Residue Management, conservation tillage and soil restoration for mitigating greenhouse effect by CO₂ – enrichment.*Soil and Tillage Research*,43: 81-107.
- Lal, R. (2014). Farm Management: encyclopedia of Agriculture and Food systems, ACADEMIC Press, 100-112.
- Landon, J.R. (1991). Booker Tropical Soil Manual.A hand-book for soil survey and agricultural land evaluation in the tropics and subtropics.John Wiley and sons Publishers, New York. 841pp.
- Masters, B. (2010). “Best Practice Clay Spreading and delving- a summary of responses to a survey of clay spreading and delving contractions”.Finalreport.Rural solutions SA.*Agronomy*,6:86-92.
- Muktar, S., Rahman, M.M. and Morrtuza, M.G. (2005).Yield and nutrient content of tomato as influenced by the application of vermicompost and chemical fertilizers. *Journal of Environmental Sciences and Natural Resources*, 8 (2): 115-122.
- Oghoghodo, I.A. and Hegar, P.I. (1995). The use of cassava Peels and Rice Husk as Soil Ammendments. *Nigerian Agriculture Journal*, 28:96-100.
- Oghoghodo, I.A., Akpowhere, M.S. and Chokor, J.U. (2008). The effects of rice husk and abbatoir effluent on soil properties, nutrient uptake, microbial species and some maize (*Zea mays* L.) growth parameters II: field experience. *Nigerian Journal of Soil Science*,18:39-47.
- Roychand, P. and Marschner, P. (2013). Respiration in a sand amended with clay Effect of residue type and rate. *European Journal of Soil Biology*, 58:19-23.
- Schweizer, M., Fear, J. and Cadisch, G. (1999).Isotopic (13C) fractionation during plant residue decomposition and its implications for soil organic matter studies. *Rapid Communications in Mass Spectrometry* 13:1284-1290.
- Singh, L., Verma, R.N.S. and Lohia, S.S. (1980).Effect of continuous application of FYM and chemical fertilizer on some soil properties.*Journal of the Indian Society of soil science*, 28:170-172.

Vanlauwe, B., Dendoven, L., Merckx, R. (1994). Residue fractionation and decomposition: the significance of the active fraction. *Plant and soil*, 158:263-274.

Wada, H. (1996). Dissimilarities between Japanese and Thai paddy soils. In: Attanadana, T. (ed). proceedings of international symposium on maximizing sustainable rice yield through improved Soil and Environmental Management. Nov. 11-17, 1996, Khonkae Thailand. 743-758.

Walkey, A. and Black, C.A. (1934). An examination of the Degtjareff method for determining soil organic matter and a proposal modification or chronic acid titration method. *Soil Sci.* 37: 29-38.

UNDER PEER REVIEW