

Estimation of genetic variability, direct and indirect effects of yield contributing characters on grain yield of upland rice (*Oryza sativa* L.) germplasm

ABSTRACT

A study was carried out with the experimental materials consisting of 24 Upland Rice genotypes grown in Randomized block design with three replications to estimate genetic variability, correlation and path coefficient analysis. Based on the mean performance, high grain yield per hill was identified for the genotype SHUATS UPR-13 followed by SHUATS UPR-11. High phenotypic and genotypic coefficient of variation were observed for grain yield per hill, biological yield per hill and number of panicles per hill. High heritability coupled with high genetic advance as percent of mean was observed for grain yield per hill, biological yield per hill, number of panicles per hill, number of tillers per hill indicating that genotypic variation present in the genetic material studied is probably due to additive genetic variance, which can be effectively exploited in crop improvement programme. The traits biological yield per hill, number of tillers per hill, number of panicles per hill, plant height, harvest index were found to have positive significant association with grain yield per hill at both genotypic and phenotypic level. Hence, selection of these traits will help in selecting the genotypes with high grain yield per hill. Number of tillers per hill, panicle length, biological yield per hill, harvest index had positively direct effect on grain yield per hill at both genotypic and phenotypic level. Therefore, effective selection must be attempted for these traits which would help in improvement of grain yield in rice genotypes suitable for different environments.

Key Words : Rice, Genetic variability, Heritability, Genetic advance, Correlation coefficient analysis and Path coefficient analysis

INTRODUCTION

Rice (*Oryza sativa* L.) belongs to the family Poaceae (Graminae) with a Chromosome no. of $2n = 24$. It is the second most important cereal crop after Wheat. It is the staple food in many parts of the world including many developing countries. (Akinola *et al.*, 2019). Approximately 90% of the world's rice is grown and consumed in Asia, whereas 50% of the population depends on rice for food. (Tenorio *et al.*, 2013). It is one of the world's largest cereal crop providing the caloric need for millions of people. It plays a pivotal role in Indian economy also is the staple food for two third of the population. Thus, to meet the challenge of producing more rice from suitable lands need of rice varieties with higher yield potential and greater yield stability to ensure food and nutritional security of India's ever-increasing population.

In India, most of the upland rice is grown in the eastern and north central states and along the south western coast. Upland rice is grown in around 6.0 million hectares of well-drained soil where the moisture stress and blast are the major constraints and productivity is around 1t/ha (ICAR-NRRI, Cuttack 2019). Principal upland rice growing states are Madhya Pradesh (1.3 million ha), West Bengal (0.88 million ha), Uttar Pradesh (0.7 million ha), Orissa (0.7 million ha), and Bihar (0.53

million ha). Globally rice is cultivated in 162 million hectares with annual production of around **503.17 million metric tonnes (United States, Department of Agriculture, Annual Report 2020).**

Grain yield is a complex polygenic character controlled by many genes interacting with the environment and is the product of many factors called yield components. The selection of parents based on yield alone is often misleading (Selvaraj *et al.*, 2011). Knowledge regarding the relative contribution of individual traits to yield may be accomplished by correlation does not provide adequate information about the contribution of each factor towards the yield. Therefore a techniques of path coefficient analysis is utilized to have an idea of direct and indirect contribution of traits on yield. The present paper deals with genetic variability, character association and path analysis in a set of 24 rice genotypes.

MATERIALS AND METHODS

Twenty four upland rice genotypes collected from SHUATS were sowing in the randomized block design with three replications, at the Experimental Farm of the Department of the Genetics and Plant breeding, Sam Higginbottom Institute of Agriculture technology and Sciences, Allahabad, during Kharif 2021. The university is situated on the left side of Allahabd Rewa National Highway, about 5km from Prayagraj city. All types of facilities necessary for cultivation of successful crop including field preparation inputs, irrigation facilities were provided from tht Department of Genetics and Plant Breeding, Sam Higginbottom Institute of Agriculture, Technology and Sciences, Prayagraj (Allahabad), U.P. In each replication, two to three seedling was sowed per hill with 20x10 cm spacing. The observations were recorded on five randomly selected plants from each plot for days to 50 percent flowering, plant height, number of tillers per hill, number of panicles per hill, panicle length, number of spikelets per panicle, flag leaf lenth, flag leaf width, test weight, days to maturity, biological yield per hill, harvest index and grain yield per hill. The statistical and biometrical analysis of GCV and PCV were calculated by formula given by Burton (1952), heritability in broad sense (h^2) by Burton and de Vane (1953) and genetic advance i.e., the expected genetic gain was calculated using the procedure given by Johnson et al., (1955). The correlation coefficient and path coefficient were worked out as for the method recommended by Al-Jibouri et al., (1958) and Deway and Lu (1959) respectievely, the estimated values were compared with table values of the correlation coefficient to test the significanceof the corellation coefficient prescribed by Fisher and Yates (1967).

Table 1 : Experimental material:

| Sr. No. | Advanced breeding lines | Cross combination | Grain type |
|----------------|--------------------------------|---------------------------------|-------------------|
| 1 | SHUATS UPR-2 | IR82589-B-B-95-2 X Vandana (P5) | Long slender |

| | | | |
|-----|---------------|-------------------------------------|----------------|
| 2 | SHUATS UPR-3 | IR82589-B-B-95-2 X Vandana (P6) | Long slender |
| 3. | SHUATS UPR-4 | IR82589-B-B-95-2 X Vandana (P7) | Long slender |
| 4. | SHUATS UPR-5 | IR82589-B-B-95-2 X NDR-97 (P1) | Long slender |
| 5. | SHUATS UPR-6 | IR82589-B-B-95-2 X NDR-97 (P3) | Long slender |
| 6. | SHUATS UPR-7 | IR82589-B-B-95-2 X NDR-97 (P5) | Long slender |
| 7. | SHUATS UPR-8 | IR82589-B-B-95-2 X NDR-97 (P6) | Long slender |
| 8. | SHUATS UPR-9 | IR82589-B-B-95-2 X NDR-97 (P7) | Long slender |
| 9. | SHUATS UPR-10 | IR82589-B-B-95-2 X NDR-97 (P8) | Long slender |
| 10. | SHUATS UPR-11 | IR82589-B-B-95-2 X NDR-97 (P9) | Long slender |
| 11. | SHUATS UPR-12 | IR82589-B-B-2-3 X NDR-97 (P1) | Medium bold |
| 12. | SHUATS UPR-13 | IR82589-B-B-2-3 X NDR-97 (P2) | Medium bold |
| 13. | SHUATS UPR-14 | IR82589-B-B-2-3 X NDR-97 (P3) | Medium bold |
| 14. | SHUATS UPR-15 | IR82589-B-B-2-3 X NDR-97 (P4) | Medium bold |
| 15. | SHUATS UPR-16 | IR82589-B-B-2-3 X NDR-97 (P5) | Medium bold |
| 16. | SHUATS UPR-17 | IR82589-B-B-2-3 X NDR-97 (P6) | Medium bold |
| 17. | SHUATS UPR-18 | IR82589-B-B-2-3 X NDR-97 (P7) | Medium bold |
| 18. | SHUATS UPR-19 | IR82589-B-B-2-3 X NDR-97 (P8) | Medium bold |
| 19. | SHUATS UPR-20 | IR82589-B-B-2-3 X NDR-97 (P9) | Medium bold |
| 20. | SHUATS UPR-21 | IR82589-B-B-2-3 X NDR-97 (P10) | Medium bold |
| 21. | SHUATS UPR-22 | IR82589-B-B-2-3 X Sahbhagidhan (P1) | Medium slender |
| 22. | SHUATS UPR-23 | IR82589-B-B-2-3 X Sahbhagidhan (P2) | Medium slender |
| 23. | SHUATS UPR-24 | IR82589-B-B-2-3 X Sahbhagidhan (P5) | Medium slender |
| 24. | SAHBHAGI DHAN | | Medium slender |

RESULTS AND DISCUSSION

Analysis of variance indicated highly significant differences among the genotypes for all the traits. This indicates that there was an ample scope for selection of promising lines from the present gene pool for yield and its components. All the characters shown significant differences at 1% level of significance.

Mean performance of grain yield per plant was high for the Rice genotypes **SHUATS UPR-13 (28.53g)** followed by **SHUATS UPR-11 (20.00g)**, **SHUATS UPR-3 (19.53g)**, **SHUATS UPR-5 (18.46g)** and **SHUATS UPR-16 (18.33g)**.

In the present investigation, it is depicted that estimates of the phenotypic coefficient of variance was found higher than their corresponding genotypic coefficient of variations, indicating that the apparent variation is not only due to genotypes but, also due to influence of environment. On an average high phenotypic and genotypic coefficient of variation were recorded for grain yield per hill, number of tillers per hill, number of panicles per hill and flag leaf length indicated the presence of variation for all the characters under study.

HERITABILITY

Heritability varied from 30.74 (days to 50% flowering) to 94.84 (grain yield per hill). High magnitude of heritability was recorded for grain yield per hill (94.84), biological yield per hill (92.78), number of tillers per hill (85.38), number of panicles per hill (84.14), flag leaf length (82.62), flag leaf width (75.49), plant height (69.70) and spikelet per panicle (69.41). Whereas, moderate heritability was recorded for Panicle length (54.11), test weight (47.09), harvest index (42.77), days to maturity (34.08), days to 50% flowering (30.74).

In the present study estimates of broad sense heritability are computed which includes both additive and non-additive gene effects. Higher values of broad sense heritability for the traits grain yield per hill, biological yield per hill, number of tillers per hill, number of panicles per hill, flag leaf length, flag leaf width, plant height and spikelet per panicle indicates the scope of genetic improvement of these characters through selection. High to moderate heritability for most of the traits in the present study indicated considerable potential for the development of high yielding varieties through desirable selection in succeeding generations.

GENETIC ADVANCE AS A PERCENTAGE MEAN (5%)

The estimation of genetic advance as percent mean helps to understand the type of gene action involved in the expression of a trait. It is classified as low (less than 10%), moderate (10-20%) and high (more than 20%) as proposed by **Johnson *et al.* (1955)**.

Genetic advance as percent mean varied from 3.26 (days to maturity) to 59.00 (grain yield per hill). high genetic advance as percent mean was recorded for grain yield per hill (59.00) followed by biological yield per hill (52.68), number of panicles per hill (36.77), number of tillers per hill (35.90), flag leaf length (28.41). Moderate genetic advance as percent mean was observed for flag leaf width (19.56), spikelets per panicle (18.62), plant height (14.98), panicle length (12.49) and harvest index

(11.08). whereas test weight (8.78), days to 50% flowering (4.27), days to maturity (3.26) recorded low genetic advance as percent mean.

In the present investigation, high genetic advance as percent of mean coupled with high heritability was recorded for grain yield per hill, biological yield per hill, number of panicles per hill, number of tillers per hill indicating most likely the heritability might be due to additive gene effect and selection may be effective in segregating generations improvement of these traits.

CORRELATION and PATH COEFFICIENT

In the present investigation, the path coefficient analysis at genotypic level revealed that the characters number of tillers per hill, panicle length, biological yield per hill, harvest index had positive direct effect on grain yield per hill. Therefore, selection programme should be based on these traits for obtaining high yielding genotypes of upland rice using pureline method of breeding.

Correlation and path coefficient studies suggest that selection based on characters Days to 50% flowering, biological yield per hill and harvest index had positive correlation and direct effects with grain yield per hill and would bring out desired improvement in yield and crop improvement programme based on these traits in upland rice.

Table 2 : Correaltion Coefficient Analysis

| Traits | | PH | NT/H | NP/H | PL | FLL | FLW | S/P | DM | BY/H | HI | TW | GY/H |
|--------|---|--------|--------|---------|---------|---------|---------|---------|---------|---------|---------|----------|---------|
| DF | P | 0.0022 | 0.0993 | 0.0817 | 0.281* | -0.0760 | 0.0854 | 0.0638 | 0.471* | -0.1148 | 0.0355 | -0.2285 | -0.0870 |
| | G | 0.0857 | 0.0752 | 0.1363 | 0.700* | -0.0697 | 0.1470 | 0.265* | 0.822** | -0.1693 | 0.371* | -0.663** | -0.0100 |
| PH | P | 1 | 0.2261 | 0.1633 | 0.286* | 0.640** | 0.313* | 0.399** | -0.0649 | 0.322* | 0.2315 | 0.1628 | 0.406** |
| | G | 1 | 0.289* | 0.2302 | 0.415** | 0.830** | 0.483** | 0.533** | -0.1722 | 0.428** | 0.397** | 0.1453 | 0.508** |
| NT/H | P | | 1 | 0.833** | 0.1296 | 0.0236 | 0.0715 | 0.316* | -0.0039 | 0.525** | 0.1927 | 0.0655 | 0.594** |
| | G | | 1 | 0.986** | 0.1697 | 0.0731 | 0.1304 | 0.402** | -0.2072 | 0.570** | 0.404** | -0.0052 | 0.664** |
| NP/H | P | | | 1 | 0.1846 | -0.0054 | 0.0979 | 0.2127 | -0.0586 | 0.469** | 0.309* | 0.0576 | 0.592** |
| | G | | | 1 | 0.1708 | -0.0309 | 0.1640 | 0.335* | -0.1737 | 0.527** | 0.473** | 0.0912 | 0.650** |
| PL | P | | | | 1 | 0.1293 | 0.531** | 0.0480 | 0.2118 | -0.0730 | 0.1171 | -0.1862 | 0.0290 |
| | G | | | | 1 | 0.289* | 0.825** | 0.1504 | 0.460** | -0.0785 | 0.407** | -0.0975 | 0.0977 |
| FLL | P | | | | | 1 | 0.355* | 0.0515 | -0.1831 | 0.1440 | 0.0385 | 0.0473 | 0.1710 |
| | G | | | | | 1 | 0.456** | 0.0736 | -0.2001 | 0.1599 | -0.0217 | 0.0083 | 0.1598 |
| FLW | P | | | | | | 1 | 0.1537 | -0.0993 | -0.0795 | 0.2217 | -0.0511 | 0.0467 |
| | G | | | | | | 1 | 0.1850 | -0.0867 | -0.0663 | 0.383** | -0.1619 | 0.0733 |
| S/P | P | | | | | | | 1 | 0.0499 | 0.366* | 0.469** | -0.1018 | 0.513** |
| | G | | | | | | | 1 | 0.0408 | 0.502** | 0.594** | -0.263* | 0.613** |
| DM | P | | | | | | | | 1 | -0.2154 | 0.249* | -0.1900 | -0.1087 |
| | G | | | | | | | | 1 | -0.303* | -0.0332 | -0.701** | -0.266* |
| BY/H | P | | | | | | | | | 1 | -0.0361 | 0.266** | 0.901** |
| | G | | | | | | | | | 1 | 0.1562 | 0.411** | 0.945** |
| HI | P | | | | | | | | | | 1 | -0.1064 | 0.382** |
| | G | | | | | | | | | | 1 | -0.288* | 0.460** |
| TW | P | | | | | | | | | | | 1 | 0.1940 |
| | G | | | | | | | | | | | 1 | 0.268* |

DF50: Days to 50% Flowering, **DM:**Days to Maturity, **PH:** Plant Height, **NT/H:** Number of Tillers per Hill, **NP/H:** Number of Panicles per Hill, **PL:** Panicle Length, **FLL:** Flag Leaf Length, **FLW:** Flag leaf width, **S/P:** Spikelets per Panicle, **DM:** Days to Maturity, **BY/H:** Biological Yield per Hill, **HI:** Harvest index, **TW:** Test weight, **P:** Phenotypic, **G:** Genotypic




Fig.1 Genotypic Correlation Diagram




fig.2 Phenotypic Correlation Diagram

UNDER REVIEW

Table 3 : Path Coefficient Analysis

| Traits | | DF50 | PH | NT/H | NP/H | PL | FLL | FLW | S/P | DM | BY/H | HI | TW | GY/H |
|-------------|---|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| DF50 | P | -0.0036 | 0 | -0.0004 | -0.0003 | -0.0010 | 0.0003 | -0.0003 | -0.0002 | -0.0017 | 0.0004 | -0.0001 | 0.0008 | -0.0870 |
| | G | -0.0081 | -0.0007 | -0.0006 | -0.0011 | -0.0057 | 0.0006 | -0.0012 | -0.0022 | -0.0120 | 0.0014 | -0.0030 | 0.0054 | -0.0100 |
| PH | P | 0.0000 | -0.0168 | -0.0038 | -0.0027 | -0.0048 | -0.0108 | -0.0053 | -0.0067 | 0.0011 | -0.0054 | -0.0039 | -0.0027 | 0.406** |
| | G | 0.0148 | 0.1729 | 0.0499 | 0.0398 | 0.0717 | 0.1435 | 0.0835 | 0.0922 | -0.0298 | 0.0740 | 0.0687 | 0.0251 | 0.508** |
| NT/H | P | 0.0038 | 0.0087 | 0.0384 | 0.0320 | 0.0050 | 0.0009 | 0.0027 | 0.0121 | -0.0001 | 0.0202 | 0.0074 | 0.0025 | 0.594** |
| | G | 0.0243 | 0.0931 | 0.3227 | 0.3181 | 0.0548 | 0.0236 | 0.0421 | 0.1298 | -0.0669 | 0.1840 | 0.1340 | -0.0017 | 0.664** |
| NP/H | P | -0.0004 | -0.0007 | -0.0037 | -0.0044 | -0.0008 | 0.0000 | -0.0004 | -0.0009 | 0.0003 | -0.0021 | -0.0014 | -0.0003 | 0.592** |
| | G | -0.0422 | -0.0713 | -0.3053 | -0.3097 | -0.0529 | 0.0096 | -0.0508 | -0.1037 | 0.0538 | -0.1633 | -0.1466 | -0.0282 | 0.650** |
| PL | P | 0.0123 | 0.0125 | 0.0057 | 0.0081 | 0.0438 | 0.0057 | 0.0233 | 0.0021 | 0.0093 | -0.0032 | 0.0051 | -0.0082 | 0.0290 |
| | G | 0.0269 | 0.0159 | 0.0065 | 0.0066 | 0.0384 | 0.0111 | 0.0316 | 0.0058 | 0.0177 | -0.0030 | 0.0156 | -0.0037 | 0.0977 |
| FLL | P | -0.0022 | 0.0182 | 0.0007 | -0.0002 | 0.0037 | 0.0284 | 0.0101 | 0.0015 | -0.0052 | 0.0041 | 0.0011 | 0.0013 | 0.1710 |
| | G | 0.0105 | -0.1253 | -0.0110 | 0.0047 | -0.0437 | -0.1510 | -0.0688 | -0.0111 | 0.0302 | -0.0241 | 0.0033 | -0.0013 | 0.1598 |
| FLW | P | -0.0009 | -0.0034 | -0.0008 | -0.0011 | -0.0057 | -0.0038 | -0.0108 | -0.0017 | 0.0011 | 0.0009 | -0.0024 | 0.0006 | 0.0467 |
| | G | -0.0012 | -0.0039 | -0.0010 | -0.0013 | -0.0066 | -0.0036 | -0.0080 | -0.0015 | 0.0007 | 0.0005 | -0.0031 | 0.0013 | 0.0733 |
| S/P | P | -0.0015 | -0.0097 | -0.0077 | -0.0052 | -0.0012 | -0.0013 | -0.0037 | -0.0243 | -0.0012 | -0.0089 | -0.0114 | 0.0025 | 0.513** |
| | G | -0.0566 | -0.1137 | -0.0858 | -0.0714 | -0.0321 | -0.0157 | -0.0395 | 0.2133 | -0.0087 | -0.1071 | -0.1267 | 0.0562 | 0.613** |
| DM | P | -0.0089 | 0.0012 | 0.0001 | 0.0011 | -0.0040 | 0.0035 | 0.0019 | -0.0009 | -0.0189 | 0.0041 | -0.0047 | 0.0036 | -0.1087 |
| | G | 0.0010 | -0.0001 | -0.0001 | -0.0001 | 0.0003 | -0.0001 | -0.0001 | 0.0000 | 0.0007 | -0.0002 | 0.0000 | -0.0005 | -0.266* |
| BY/H | P | -0.0907 | 0.2544 | 0.4150 | 0.3709 | -0.0577 | 0.1138 | -0.0628 | 0.2888 | -0.1701 | 0.7901 | -0.0285 | 0.2103 | 0.901** |
| | G | -0.1793 | 0.4529 | 0.6040 | 0.5584 | -0.0831 | 0.1693 | -0.0703 | 0.5319 | -0.3212 | 0.9240 | 0.1654 | 0.4350 | 0.945** |
| HI | P | 0.0134 | 0.0873 | 0.0727 | 0.1164 | 0.0441 | 0.0145 | 0.0836 | 0.1767 | 0.0940 | -0.0136 | 0.3770 | -0.0401 | 0.382** |
| | G | 0.1434 | 0.1534 | 0.1561 | 0.1829 | 0.1572 | -0.0084 | 0.1481 | 0.2296 | -0.0128 | 0.0604 | 0.3864 | -0.1113 | 0.460** |
| TW | P | 0.0008 | -0.0005 | -0.0002 | -0.0002 | 0.0006 | -0.0002 | 0.0002 | 0.0003 | 0.0006 | -0.0009 | 0.0004 | -0.0033 | 0.1940 |
| | G | 0.0537 | -0.0118 | 0.0004 | -0.0074 | 0.0079 | -0.0007 | 0.0131 | 0.0213 | 0.0568 | -0.0333 | 0.0233 | -0.0810 | 0.268* |
| GY/H | P | -0.0870 | 0.406* | 0.594** | 0.592** | 0.0290 | 0.1710 | 0.0467 | 0.513** | -0.1087 | 0.901** | 0.382** | 0.1940 | 1.0000 |
| | G | -0.0100 | 0.508** | 0.664** | 0.650** | 0.0977 | 0.1598 | 0.0733 | 0.613** | -0.266* | 0.945** | 0.460** | 0.268* | 1.0000 |

DF: Days to 50% flowering, **PH:** Plant Height, **NT/H:** Number of Tillers per hill, **NP/H:** Number of Panicles per hill, **PL:** Panicle Length, **FLL:** Flag Leaf Length, **FLW:** Flag leaf width, **S/P:** Spikelets per Panicle, **DM:** Days to Maturity, **BY/H:** Biological Yield per Hill, **HI:** Harvest index, **TW:** Test weight, **P:** Phenotypic, **G:** Genotypic




Fig.3 Genotypic Path Coefficient




fig.4 Phenotypic Path Coefficient

CONCLUSION

From the present investigation, it is concluded that SHUATS UPR – 13 (9.41 g) was found superior in grain yield per hectare followed by SHUATS UPR – 11 (6.6 g) over the check variety SAHBHAGI DHAN (4.37g). Among 24 genotypes of upland rice Analysis of variance revealed that mean sum of squares due to genotypes were significant for all the characters studied, suggesting the existence of high genetic variability among the genotypes for all the traits. High GCV and PCV were recorded for grain yield per hill followed by biological yield per hill and number of panicles per hill. High heritability coupled with high genetic advance as percent mean was recorded for characters like grain yield per hill, biological yield per hill, number of panicles per hill and number of tillers per hill. The traits biological yield per hill, number of tillers per hill and number of panicles per hill recorded positive significant association with grain yield per hill at both genotypic and phenotypic level. Hence, these characters emerged as most important associate of grain yield per hill and selection of these traits will help in selecting genotypes with high grain yield. Number of tillers per hill, panicle length and biological yield per hill had high positive direct effect on grain yield per hill. Therefore, direct selection for these traits will be rewarding for yield improvement in Upland rice.

REFERENCES

- Ahamed, S., Chowdhury, R., Atik, Ahmed, S., Hossain, S. and Matin, M.N. (2021).** Analysis of Genetic Variability, Heritability and Correlation among Bangladeshi Local Rice Cultivars (*Oryza sativa* L.). *Pakistan Journal of Agricultural Research*, **34**(2) :356-361
- Agrawal, K. B. (2003).** Variability studies in segregating population of rice. *Anna Agriculture Research*, **24**(4): 707-709.
- Barhate, K.K., Jadhav, M.S. and Bhavsar, V. (2021).** Genetic variability, heritability and genetic advance in aromatic lines of rice (*Oryza sativa* L.) *Journal of Pharmacognosy and Phytochemistry*, **10**(3): 360-362
- Chamar, J.K., Joshi, R.P., Katkani, D., Sahu D. and Patel, V. (2021).** Genetic variability, heritability and correlation coefficient analysis in rice (*Oryza sativa* L.) germplasm. *The Pharma Innovation Journal*, **10**(5): 763-769
- Ekka, R., Sarawgi, A. K. and Raja, R. (2011).** Correlation and path analysis in traditional rice accessions of Chhattisgarh. *Journal of Rice Research*, **4**(1&2): 11-17.
- Elsheik, P.K., S. Chakraborty., Barman. and K.K. Sarmah. (2007).** Genetic variation for harvest index grain yield and yield components in boro rice. *Oryza*, **38**(3&4): 149-150.
- Fisher, R.A. (1936).** The use of multiple measurements in taxonomic problems. *Annals of Eugenics*, **7**(2).
- Fukrei P., Amit, K., Tyagi, W., Rai M. and Pattanayak, A. (2011).** Genetic variability in yield and its components in upland rice grown in acid soils of North East India. *Journal of Rice Research*, **4**(1& 2): 4-7.
- Hossain, S., Haque, M. and Rahman, J. (2015).** Genetic variability, correlation and path coefficient analysis of morphological traits in some extinct local Aman rice (*Oryza sativa* L.). *Journal of Rice Research*, **4**(1).
- Islam M.Z., Khalequzzaman M., Bashir M. K., Ivy N. A., Haque M.M. and Mian M.A.K. (2016).** Variability Assessment of Aromatic and Fine Rice Germplasm in Bangladesh Based on Quantitative Traits. *The Scientific World Journal*, **pp: 14**.
- Jayasudha, S. and Deepak, S. (2010).** Genetic parameters of variability, correlation and path-coefficient for grain yield and physiological traits in rice (*Oryza sativa* L.) under shallow lowland situation. *Electronic Journal of Plant Breeding*, **1**(5): **1332-1338**.
- Kishor, C., yogendra, P., Haider, Z. A., Kumar, R., and Kamleshwar. K. (2008).** Quantitative analysis of upland rice. *Oryza*, **45**(4): 268-272.
- Lingaiah, N. (2015).** Genetic variability, heritability and genetic advance in rice (*Oryza sativa* L.). *Asian Journal of environmental Sciences*, **10**(1): 110-112.

- Marekar, R.V. and Siddiqui, M. A. (1996).** Genetic variability and correlation studies in rice. *Journal of Maharashtra Agricultural Universities*, **21**(2): 249-251.
- Prasad, G.S.V., Prasad, A.S.R., Sastry, M.V.S. and Srinivasan, T.E. (2013).** Genetic relationship among yield components in rice. *Indian Journal of Agriculture Sciences*, **58**(6): 470-472.
- Rahman, S.S., Mishra, S. B. and Ali, F. (2012).** Estimation of grain yield and inter-relationship with yield components in upland rice. *Mysore Journal of Agriculture Sciences*, **34**: 142-146.
- Saidaiah, P., Sudheer, S. and Ramesha, M. S. (2010).** Variability for yield and yield component attributes in rice. *Crop Research*, **39**(1, 2 & 3): 91-93.
- Tiwari, G. C. (2015).** Variability, heritability and genetic advance analysis for grain yield in rice. *Journal of Engineering Research and Application*. **Vol 5: pp 46-49.**
- Verma O. P., Santoshi, U., Dwivedi, J. L. and Singh, P. P. (2000).** Genetic variability, heritability and genetic advance for quantitative traits in rice. *Oryza*, **37** (2): 38-40.
- Yadav, R. K. (2000).** Genetic variability in some varieties of upland rice in Chattisgarh State. *Indian journal of Dryland Agricultural Research & Development*, **15**(2): 166-167.