

Original Research Article

Crop Phenology, Growth Indices and Yield of Wheat as Influenced by Different Establishment and Moisture Conservation Practices under Limited Irrigated Conditions

Abstract

The field experiment was conducted at the Research Farm of CCS Haryana Agricultural University, Hisar during the *Rabi* seasons of 2019-20 and 2020-21. The experiment consisted of twenty four treatment combinations comprising three establishment methods *i.e.* Conventional Tillage (CT), Zero Tillage (ZT) and Bed Planting (BP) and two irrigation levels *i.e.* I₁-one irrigation applied at crown root initiation stage (I₁) and I₂-two irrigation applied at CRI and booting stages as main plot treatments and four moisture conservation practices *i.e.* M₀-no mulch, M₁-mulch (pearl millet straw @ 4 tonnes ha⁻¹), M₂- antitranspirant (kaolin @ 6% w/v) and M₃- mulch (pearl millet straw @ 4 tonnes ha⁻¹) + antitranspirant (kaolin @ 6% w/v) as sub-plots treatment, were tested in split split-plot design with three replications. The results revealed that days taken to booting, heading and physiological maturity did not influence significantly by crop establishment methods. However, among the irrigation levels, maximum days taken for booting (79.64 and 77.89), heading (90.33 and 88.08) and physiological maturity (138.86 and 137.42) stages under two irrigation (I₂) and minimum *i.e.* 78.81 and 75.64; 88.83 and 86.11; 136.83 and 135.14 were taken in one irrigation during 2019-20 and 2020-21, respectively. Among moisture conservation practices significantly higher days taken for booting (80.24 and 79.06), heading (91.00 and 88.39) and physiological maturity (140.39 and 138.72) stages in M₃ treatment over M₀ 78.41 and 74.50; 87.78 and 85.67; 135.33 and 133.72 but at par with M₁ and M₂. The growth indices *i.e.* crop growth rate and relative growth rate are significantly higher in BP, two irrigation levels and M₃ during 2019-20 and 2020-21, respectively. The maximum grain yield was recorded in BP 4873 and 4423; two irrigation levels 4,880 and 4,465 and M₃ 4,879 and 4,468 which was significantly higher as compared to other treatments during 2019-20 and 2020-21, respectively.

Keywords: Wheat, Establishment, Irrigation Levels, Moisture Conservation, Phenology, Growth Indices and Grain Yield

Introduction

Wheat is one of the most important cereal ~~erocrops~~ crops of the world on account of its wide adaptability to cultivation under different agro-climatic and soil conditions. In India, it is the second most important source of staple food after rice. In India, area, production and average yield of wheat is 31.76 m ha, 108.75 m t and 34.24 q ha⁻¹, respectively during 2020-21. In Haryana, it is grown over an area of 2.52 m ha with a production of 12.15 m t and productivity of 48.22 q ha⁻¹ during 2020-21 (ICAR-IIWBR, 2020-21). Wheat is usually planted either by drilling closely spaced rows apart on the flat bed or by broadcasting the seed on a levelled soil surface and then incorporating it by means of shallow tillage operations. Real challenges for today's Indian agricultural system are resource fatigue with stagnating productivity and profitability, decreasing human resources and their rising costs and socioeconomic changes (Kumar *et al.*, 2013). The new tillage and crop establishment technologies like minimum, zero till and furrow irrigated raised broad bed could answer the saving energy, resources and increasing yield of different wheat-based cropping systems (Singh *et al.*, 2013). In this sense, alternative best crop management options like zero tillage and beds have demonstrated potential benefits on crop yield and profits while saving water, energy and restoring soil degradation across diverse ecologies (Das *et al.*, 2014).

Wheat is highly sensitive to water stress during the CRI and flowering stage and ultimately decrease the yield. Thus, timing the length of irrigation interval with the stages of crop growth might bring about a reduction in the number of irrigations and results in an economic crop yield. In principle, irrigation should take place while the soil water potential is still high enough to enable soil supply water fast enough to meet the local atmospheric demands without placing the plants under stress that would reduce yield and quality of crop. Singh *et al.* (2017) reported that the application of two irrigations at tillering and flowering stage (I₂) resulted in significant increase in CGR (g⁻¹ m²-day⁻¹), RGR (g⁻¹ g⁻¹ day⁻¹) and LAI than no irrigation (I₀) and one irrigation (I₁) treatments during both the years.

Mulching has been proved to be useful in conserving moisture and increasing productivity of wheat. Straw mulch also provides benefit in terms of increasing infiltration rate, lowers the temperature, ~~improvesand improves~~ improves fertilizer availability and increase crop yield (Singh *et al.*, 2011).

Mulches enhanced soil water status and improved growth and yield, which subsequently reducing runoff and evaporation losses. Angbabu *et al.* (2007) reported that the combined application of straw mulch at @ 6 t ha⁻¹ + kaolin spray at @ 6.0% w/v significantly influenced LAI, ~~Cerop gGrowth #Rate~~ CGR and ~~N#et aAssimilation #Rate~~ NAR during both the years of investigation. With these leads,

the present investigation was formulated to study the effects of crop establishment methods, irrigation schedule and moisture conservation practices on the performance of wheat in terms of phenology, growth indices and grain yield under semi-arid conditions.

Materials and Methods

The field experiments were conducted at the Research Farm of Chaudhary Charan Singh Haryana Agricultural University, Hisar during the *Rabi* seasons of 2019-20 and 2020-21. The experiment consisted of twenty four treatment combinations comprising three establishments *i.e.* Conventional Tillage (CT), Zero Tillage (ZT) and bed planting (BP) and two irrigation levels *i.e.* one irrigation applied at Crown Root Initiation (CRI) (I_1) and two irrigation applied at CRI and booting stages (I_2) as main plot treatment and four moisture conservation practices *i.e.* no mulch (M_0), mulch (pearl millet straw @ 4 tonnes ha^{-1}) (M_1), antitranspirant (kaolin @ 6% w/v) (M_2) and mulch (pearl millet straw @ 4 tonnes ha^{-1}) + antitranspirant (kaolin @ 6% w/v) as sub-plots treatment, were tested in a split split-plot design with three replications. The crop was sown in the second week of November and the last week of November during 2019-20 and 2020-21, respectively. Fertilizer doses of 150 kg N, 60 kg P_2O_5 and 40 kg K_2O ha^{-1} was applied as per the recommendation. Half dose of N and full dose of P and K were applied at the time of sowing while the remaining half dose of N applied to the crop at first irrigation. Agronomic practices, *i.e.* weeding, hoeing and plant protection measures were carried out as per recommendations at appropriate times. Data were recorded on days taken to emergence, days taken to booting, days taken to heading, days taken to physiological maturity, crop growth rate, relative growth rate and grain yield as per the standard procedure. Data collected during the study were statistically analyzed by using the technique of analysis of variance (ANOVA) described by Cochran and Cox (1959).

Results and Discussion

The data on days taken to the emergence, days taken to booting, days taken to heading and days taken to physiological maturity stages of wheat as influenced by crop establishment methods, irrigation schedules and moisture conservation practices are presented in Table 1 and illustrated in figures 1 and 2. The data indicate that the crop establishment methods, irrigation schedules and moisture conservation practices did not influence emergence of wheat during both the years. Days taken to the booting, heading and physiological maturity stages of wheat did not influence by crop establishment methods during both the years. The data indicate that the days taken to the booting, heading and physiological maturity stages of wheat differed significantly by irrigation levels. At booting stage, significantly more days were taken by the crop in I_2 treatment (79.64 and 78.61) as compared to I_1 (78.81 and 75.64) during 2019-20 and 2020-21, respectively. More days were taken

for heading by the crop in I₂ treatment (90.33 and 88.08) which were significantly higher than I₁ (88.83 and 86.11) during 2019-20 and 2020-21, respectively. As well as significantly more days were taken by the crop in I₂ treatment (138.86 and 137.42) to attain the physiological stage as compared to days taken in I₁ (136.83 and 135.14) during both the years, respectively. Under higher levels of irrigation, the longer reproductive phase of crop growth and longer crop duration has been also reported by many workers (Zabihullah, 2020; Kumar *et al.*, 2018; Ngwako and Mashiqa, 2013; Dhaka *et al.*, 2006). Under higher levels of irrigation, the longer reproductive phase of crop growth and longer crop duration has been also reported by Kumar *et al.* (2018).

Among the moisture conservation practices, data indicate that treatments M₃ (80.24 and 79.06) and M₁ (79.33 and 76.89) significantly took more days for the booting stage as compared to M₂ (78.91 and 76.61 days) and M₀ (78.41 and 74.50 days) during 2019-20 and 2020-21, respectively. Also, among the moisture conservation treatment M₃ (91.00 and 88.39) and M₁ (90.00 and 87.44) significantly took more days for the heading stage than M₂ (89.56 and 86.89 days) and M₀ (87.78 and 85.67 days) during 2019-20 and 2020-21, respectively. As well as, M₃ (140.39 and 138.72) and M₁ (138.33 and 136.94) significantly took more days to attain the physiological maturity stage than M₂ (137.33 and 135.72 days) and M₀ (135.33 and 133.72 days) during both the years, respectively. Whereas, M₀ (no mulch) took less days for the booting, heading and physiological maturity stages during both the years.

The data on Crop Growth Rate (CGR) are presented in Table 1 showed that there was no significant difference in CGR due to crop establishment methods at 0-30 and 31-60 DAS during both the years. At 61-90 DAS, maximum CGR was recorded with BP (4.30 and 4.05 g dm⁻² day⁻¹) which was significantly greater than ZT (4.22 and 3.96 g dm⁻² day⁻¹) but it was at par with CT (4.26 and 4.00 g dm⁻² day⁻¹) during 2019-20 and 2020-21, respectively. A similar trend was also observed by Singh (2017). Among the irrigation levels, CGR differed significantly due to irrigation levels at 91-120 DAS and 121 DAS and at harvest. At 91-120 DAS, maximum CGR (2.44 and 1.99 g dm⁻² day⁻¹) was recorded in the treatment I₂ (two irrigation were applied to the crop) which was significantly higher over one irrigation (I₁) (2.24 and 1.93 g dm⁻² day⁻¹) during 2019-20 and 2020-21, respectively. Similar trends was also noticed at 120 DAS- at harvest stage during both the years These results are in conformity with the findings of Vishuddha *et al.* (2014) and Kumar *et al.* (2015) who also reported the maximum values of growth indices under more number of irrigation. Among the moisture conservation practices there was no significant differences in CGR at 0-30 DAS during both the years. At 61-90 DAS, application of straw mulch and antitranspirant (M₃) recorded significantly higher CGR (4.33 and 4.07 g dm⁻² day⁻¹) as compared to all other treatments, *i.e.* M₀ (4.19 and 3.92 g dm⁻² day⁻¹), M₂ (4.26 and 4.00 g dm⁻² day⁻¹) and M₁ (4.28 and 4.01 g dm⁻² day⁻¹) during 2019-20 and

2020-21, respectively. Similar trends was also recorded at 91-120 and 121 DAS to harvest. Brahma *et al.* (2007) obtained similar results and reported that kaolin contributed to enhanced growth indices. Wairagade *et al.* (2020) recorded significantly higher CGR and RGR under the application of wheat straw mulch over control.

Data pertaining to the RGR at various stages was analyzed statistically and the data as influenced by different treatments are given in Table 1. The data revealed that RGR attained maximum value between 0-30 and 31-60 DAS stages and then declined consistently till crop maturity during both the years of study. However, there was no significant differences in RGR at 0-30 and 31-60 DAS due to crop establishment methods during both the years. Among methods of crop establishment, at 61-90 DAS stage, maximum RGR was recorded with BP (1.44 and 1.38 g/g/day) which was significantly higher than ZT (1.42 and 1.36 g/g/day) but was at par with CT (1.43 and 1.37 g/g/day) during 2019-20 and 2020-21, respectively. Similar trend was also reported by Singh (2017). The data on RGR presented in Table 3 showed that RGR not influence RGR by irrigation level significantly up to 90 DAS stage during both the years of experimentation. However, at 91-120 DAS, application of two irrigations (I_2) recorded significantly higher values of RGR (0.84 and 0.64 g/g/day) than one irrigation applied at CRI stage (I_1) (0.75 and 0.61 g/g/day) during 2019-20 and 2020-21, respectively. These results are in conformity with the findings of Vishuddha *et al.* (2014) and Kumar *et al.* (2015) who also reported the maximum values of growth indices under higher levels of irrigation. Among the moisture conservation practices, there was no significant differences in RGR at 0-30 DAS during both the years. At 61-90 DAS, M_3 recorded significantly higher CGR (1.44 and 1.38 g/g/day) over all other treatments *i.e.* M_0 (1.39 and 1.34 g/g day), M_2 (1.41 and 1.36 g/g day) and M_1 (1.42 and 1.36 g/g/day) during 2019-20 and 2020-21, respectively. Similar trends was also recorded at 91-120 and 121 DAS to harvesting stage during both the years. Among moisture stress management practices, SH + mycorrhizae + KCl spray + kaolin spray recorded significantly highest growth indices, viz. LAI, CGR and RCG over on-management treatment (Patil *et al.*, 2014). Brahma *et al.* (2007) also reported that kaolin contributed to enhanced growth indices. Wairagade *et al.* (2020) found significantly higher CGR and RGR under the application of wheat straw mulch treatment over control. In general, the grain yield was recorded higher during 2019-20 as compared that of 2020-21. Bed planting (BP) produced significantly higher grain yield of wheat (4,873 and 4,423 Kg ha⁻¹) over ZT (4,645 and 4,220 Kg ha⁻¹) but statistically at par with CT (4,751 and 4,330 Kg ha⁻¹) during 2019-20 and 2020-21, respectively. Whereas, lower grain yield of 4,645 during 2019-20 and 4,220 Kg ha⁻¹ during 2020-21 was recorded in ZT. Similar trends have been observed by Singha *et al.* (2018), Kumar *et al.* (2018) and Singh *et al.* (2019). Sagar *et al.* (2017) reported higher grain in bed planting

than another crop establishment. Higher grain yield with bed planting of wheat has been also reported by Thind *et al.* (2010).

Grain yield was significantly influenced by different levels of irrigations. The higher grain yield of wheat (4,880 and 4,465 Kg ha⁻¹) was obtained in I₂ treatment (two irrigation) which was significantly higher than I₁ treatment (4,633 and 4,184 Kg ha⁻¹) during 2019-20 and 2020-21, respectively. Similar results were reported by Singh and Katiyar, 2014; Sepat *et al.* (2015), Singh *et al.*, 2018; Kumar *et al.*, 2018. Also, Kaur *et al.* (2018) noticed that the higher yield is due to the strong effect of irrigation frequency and mulch on grain yield. As well as that moisture conservation practices significantly affected the grain yield of wheat. The maximum grain yield (4,879 and 4,468 Kg ha⁻¹) was produced in the treatments when mulch and antitranspirant were applied to the crop (M₃) which was significantly higher over M₀ (4,589 and 4,154 Kg ha⁻¹) but statistically at par with M₁ (4,785 and 4,341 Kg ha⁻¹) and M₂ (4,772 and 4,333 Kg ha⁻¹) during both the years of investigation respectively. Similar findings were also been described by (Ahmed *et al.*, 2009; Sarwar *et al.*, 2013; Patil *et al.*, 2014; Singh *et al.*, 2018) Also, Al-Amin *et al.* (2017) noticed that the higher grain yield in straw mulch @ 6 t ha⁻¹ while the lowest grain yield was recorded in control. This may occur due to the different environmental factors and cultural management practices.

Table 1: Effect of Different Crop Establishment Methods and Moisture Conservation Practices on Phenology Stages and Grain Yield of Wheat Under Limited Irrigated Conditions.

Treatments	Days taken to								Grain Yield (kg ha ⁻¹)	
	Emergence		Booting		Heading		Physiological Maturity		2019-20	2020-21
	2019-20	2020-21	2019-20	2020-21	2019-20	2020-21	2019-20	2020-21		
Crop Establishment Methods										
CT	6.35	7.76	79.71	77.29	90.13	87.92	138.63	137.08	4,751	4,330
ZT	6.24	6.99	78.63	76.00	89.00	86.08	136.50	135.00	4,645	4,220
BP	6.25	7.53	79.33	77.00	89.63	87.29	138.42	136.75	4,873	4,423
SEm±	0.10	0.14	0.30	0.84	0.49	0.65	0.70	0.77	46	49
CD at 5 %	NS	NS	NS	NS	NS	NS	NS	NS	145	154
Irrigation Schedules										
I ₁	6.27	7.41	78.81	75.64	88.83	86.11	136.83	135.14	4,633	4,184
I ₂	6.29	7.45	79.64	77.89	90.33	88.08	138.86	137.42	4,880	4,465
SEm±	0.05	0.09	0.21	0.69	0.40	0.53	0.57	0.63	37.51	40
CD at 5 %	NS	Ns	0.73	2.17	1.25	1.67	1.81	1.98	118	126
Moisture Conservation Practices										
M ₀	6.00	7.17	78.41	74.50	87.78	85.67	135.33	133.72	4,589	4,154
M ₁	6.50	7.41	79.33	76.89	90.00	87.44	138.33	136.94	4,785	4,341

M ₂	6.02	7.32	78.91	76.61	89.56	86.89	137.33	135.72	4,772	4,333
M ₃	6.19	7.02	80.24	79.06	91.00	88.39	140.39	138.72	4,879	4,468
SEm±	0.10	0.13	0.44	0.66	0.24	0.27	0.13	0.15	48	41
CD at 5 %	NS	NS	1.27	1.88	0.70	0.77	0.39	0.44	136	119




Fig. 1. Phenology of Wheat Crop As Influenced By Crop Establishment Methods, Irrigation Schedules and Moisture Conservation Practices in Wheat Crop during 2019-20




Fig. 2. Phenology of Wheat Crop As Influenced By Crop Establishment Methods, Irrigation Schedules and Moisture Conservation Practices in Wheat Crop during 2020-21

UNDER PEER REVIEW

Table 2: Effect of Different Crop Establishment Methods and Moisture Conservation Practices on Crop Growth Rate (Cgr) (g/m²/day) at Different Stages of Wheat Under Limited Irrigated Conditions


Treatments	Crop Growth Rate (CGR) (g/m ² /day)									
	At 0-30 DAS period		At 31-60 DAS period		At 61-90 DAS period		At 91- 120 DAS period		At 121 DAS – harvest period	
	2019-20	2020-21	2019-20	2020-21	2019-20	2020-21	2019-20	2020-21	2019-20	2020-21
Crop Establishment Methods										
CT	0.134	0.116	2.08	1.92	4.25	3.97	2.34	1.96	1.99	1.86
ZT	0.131	0.105	2.06	1.89	4.21	3.96	2.19	1.93	1.96	1.82
BP	0.139	0.117	2.13	1.94	4.29	4.04	2.55	2.01	2.14	1.91
SEm±	0.006	0.006	0.01	0.01	0.02	0.02	0.06	0.02	0.04	0.02
CD at 5 %	NS	NS	NS	NS	0.05	0.07	0.20	0.05	0.12	0.06
Irrigation Schedules										
I ₁	0.136	0.113	2.10	1.94	4.24	3.96	2.26	1.94	1.93	1.83
I ₂	0.133	0.112	2.10	1.90	4.26	4.02	2.46	2.00	2.08	1.90
SEm±	0.005	0.005	0.01	0.01	0.01	0.02	0.05	0.01	0.03	0.02
CD at 5 %	NS	NS	NS	NS	NS	NS	0.16	0.04	0.10	0.05
Moisture Conservation Practices										
M ₀	0.131	0.113	1.99	1.84	4.19	3.91	2.14	1.90	1.85	1.74
M ₁	0.135	0.117	2.11	1.93	4.26	4.00	2.36	1.97	2.04	1.88
M ₂	0.133	0.110	2.10	1.91	4.25	3.99	2.38	1.96	2.03	1.87
M ₃	0.139	0.110	2.15	2.00	4.31	4.06	2.56	2.04	2.19	1.97
SEm±	0.004	0.004	0.02	0.02	0.02	0.02	0.06	0.02	0.04	0.03
CD at 5 %	NS	NS	0.06	0.06	0.04	0.05	0.17	0.06	0.12	0.08

Table 3: Effect of Different Crop Establishment Methods and Moisture Conservation Practices on Relative Growth Rate (Rgr) (g/g/day) at Different Stages of Wheat Under Limited Irrigated Conditions.

Treatments	Relative Growth Rate (RGR) (g/g/day)									
	At 0- 30 DAS Period		At 31- 60 DAS Period		At 61-90 DAS Period		At 91- 120 DAS Period		At 121 DAS- harvest Period	
	2019-20	2020-21	2019-20	2020-21	2019-20	2020-21	2019-20	2020-21	2019-20	2020-21
Crop Establishment Methods										
CT	0.071	0.062	0.81	0.72	1.41	1.36	0.80	0.62	0.66	0.60
ZT	0.070	0.062	0.80	0.71	1.40	1.35	0.73	0.61	0.63	0.58
BP	0.070	0.067	0.81	0.74	1.42	1.38	0.87	0.64	0.72	0.64
SEm±	0.001	0.002	0.005	0.01	0.004	0.01	0.03	0.01	0.02	0.01
CD at 5 %	NS	NS	NS	NS	0.014	0.02	0.08	0.03	0.05	0.03
Irrigation schedules										
I ₁	0.070	0.063	0.81	0.73	1.41	1.36	0.76	0.61	0.63	0.58
I ₂	0.070	0.064	0.81	0.71	1.42	1.37	0.84	0.64	0.71	0.62
SEm±	0.001	0.001	0.004	0.01	0.004	0.01	0.02	0.01	0.01	0.01
CD at 5 %	NS	NS	NS	NS	NS	NS	0.06	0.02	0.04	0.03
Moisture conservation Practices										
M ₀	0.070	0.063	0.78	0.67	1.38	1.34	0.71	0.58	0.58	0.53
M ₁	0.070	0.064	0.81	0.73	1.42	1.36	0.80	0.62	0.68	0.61
M ₂	0.068	0.062	0.80	0.71	1.41	1.36	0.81	0.63	0.67	0.60
M ₃	0.071	0.065	0.83	0.77	1.44	1.38	0.89	0.67	0.75	0.66
SEm±	0.002	0.001	0.005	0.01	0.004	0.01	0.02	0.01	0.02	0.02
CD at 5 %	NS	NS	0.014	0.03	0.011	0.01	0.07	0.03	0.06	0.04

Put the full meaning of abbreviations as foot note e.g. **DAS** below each table

UNDER PEER REVIEW



Conclusion

BP sowing was found suitable in terms of phenology, growth indices and yield among crop establishment methods and wheat crop perform better with the application of two irrigations as well as the application of kaolin as anti-transpiration under limited irrigation conditions.

References

- Ahmed, Z.I., Ansar, M., Iqbal, M. and Minhas, N.M. Effect of planting geometry and wheat straw mulch on performance of wheat under rainfed conditions of Pothohar. *Pak. J. Agric. Res.* 2009; **22**: 112-115.
- A-Amin, M.A., Hasan, A.K., Ali, M.H., Nessa, S. and Islam M.N. Effect of mulching and organic manure on growth and yield performance of wheat. *Archives of Agriculture and Environmental Science* 2017; **2**(3): 134-140.
- Angbabu, S., Pravat, P., Sarkar, A.K. and Sinha, A.C. Growth and yield of wheat as influenced by evapotranspiration control measures and levels of fertilizer under rain fed condition. *Indian Journal of Plant Physiology* 2007; **12**(2): 194-198
- Bahma, R., Janawade, A.D. and Palled, Y.B. Effect of irrigation schedules, mulch and antitranspirant on growth, yield and economics of Wheat. *Karnataka Journal of Agricultural Science* 2007; **20**(1): 6-9.
- Cochran, W.C. and Cox, G.M. *Experimental Designs* (2nd Edn.) Wiley Publisher, 1992; 640.
- Das, T.K., Bhattacharyya, R., Sudhishri, S., Sharma, A.R., Saharawat, Y.S., Bandyopadhyay, K.K., Sepat, S., Bana, R.S., Aggarwal, P., Sharma, R.K., Bhatia, A., Singh, G., Datta, S.P., Kar A., Singh B., Singh P., Pathak, H., Vyas, A.K. and Jat, M.L. Conservation agriculture in an irrigated cotton-wheat system of the western Indo-Gangetic Plains: crop and water productivity and economic profitability. *Field Crops Research*. 2014; **58**: 24-33.
- Dhaka, A.K., Bangarwa, A.S., Pannu, R.K., Malik, R.K and Garg, R. Phenological development, yield and yield attributes of different wheat genotypes as influenced by sowing time and irrigation. *Agricultural Science Digest* 2006; **26**(3):174-177.
- Idhani, L.K. and Kumar, A. Relative efficiency of different irrigation schedules for conventional, ridge and raised bed seeding of wheat (*Triticum aestivum*). *Indian Journal of Agronomy* 2012; **57** (2): 148-151.
- Indian Council of Agricultural Research-Indian Institute of wheat and barley research (ICAR-IIWBR). Directors report of AICRP on wheat and barley, Ed: G. P. Singh. ICAR- India Institute of wheat and barley research, Karnal, Haryana, India. 2021; P72.

Comment [B1]: Write in full

- Kaur, R., Kumar, S., Singh, G. and Kaur, R. Effect of irrigation scheduling and mulches on growth and yield of wheat (*Triticum aestivum* L.) in Central Punjab. *International Journal of Chemical Studies* 2018; (5): 723-726.
- Kumar, B., Dhar, S., Vyas, A.K. and Paramesh, V. Impact of irrigation schedules and nutrient management on growth, yield and root traits of wheat (*Triticum aestivum*) varieties. *Indian Journal of Agronomy* 2015; **60**(1):87- 91.
- Kumar, M., Pannu, R.K. and Singh, B. Effect of irrigation regimes and nitrogen levels on phenology and grain yield of late sown wheat. *Wheat and Barley Research*. 2018; **10**(1): 15- 19.
- Kumar, V., Saharawat, Y.S., Gathala, M.K., Jat, A.S., Singh, S.K. and Chaudhary, N. Effect of different tillage and seeding methods on energy use efficiency and productivity of wheat in the Indo-Gangetic Plains. *Field Crops Research* 2013; **142**: 1–8.
- Ngwako S and Mashaqa PK. The effect of irrigation on growth and yield of winter wheat (*Triticum Aestivum* L.), *International Journal of Agriculture Crop Science*, 2013; **5**: pp 976-982.
- Patil, M.D., Dhindwal, A.S. and Rajanna, G.A. Integrated moisture stress management in wheat (*Triticum aestivum*). *Indian Journal of Agronomy* 2014; **59**(4): 629-633
- Sarwar, M.A., Akbar, N., Javeed, H.M.R., Shehzad, M.A., Mehmood, A. and Abbas, H.T. Response of zero tilled wheat crop to different mulching techniques in a semi-arid environment. *International Journal of Advanced Research* 2013; **1**(9): 768-776.
- Sapat, S., Sharma, A.R., Dinesh, K. and DAS, T.K. Effect of conservation agriculture practices on productivity and sustainability of pigeonpea (*Cajanus cajan*)-wheat (*Triticum aestivum*) cropping system in Indo-Gangetic plains of India. *Indian Journal of Agricultural Sciences*; 2015; **85**(2): 212-218.
- Singh, B., Gaydon, D.S., Humphreys, E. and Eberbach, P.L. The effects of mulch and irrigation management on wheat in Punjab, India-Evaluation of the APSIM model. *Field Crops Research* 2011; **124**(1):1-13.
- Singh, V. Evaluation of tillage-cum-crop establishment methods under irrigation scheduling on soil properties, water productivity and yield of wheat (*Triticum aestivum* l.). Ph.D Thesis. Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut-250110 (U P), India 2017. .
- Singh, Y.K. and Katiyar, T.P.S. Effect of tillage system, irrigation scheduling and depth on growth, yield attribute, yield and water use in wheat (*Triticum aestivum*). *Int J Tropical Agri*. 2014; **32** (1/2): 512-521.
- Singh, Y.P., Singh Dharvendra, Tomar, S.S. and Gupta, R.K. Effect of time of pre-irrigation and tillage practices on wheat (*Triticum aestivum* L.) under pigeon pea (*Cajanus cajan*) - wheat cropping sequence. *Indian Journal of Agricultural Sciences* 2013; **83**(12): 1317-1321.

Comment [B2]: Full meaning

- Singh YP, Singh S, Singh AK and Panwar B. Influence of wheat establishment techniques and previous *Kharif* season crops on productivity, profitability, water use efficiency, energy indices and soil properties in central India. *Agricultural Research* 2019; DOI: 10.1007/s40003-019-00420-y.
- Singha P, Mondal T, Patra K, Mitra B. Straw mulch and restricted irrigation effect on productivity, profitability and water use in wheat under various crop establishment technique in Eastern Sub-Himalayan Plains of India. *Int. J Curr. Microbi. App. Sci.* 2018; 7(2):1521-1533.
- Thind HS, Buttar GS and Aujla MS. Yield and water use efficiency of wheat and cotton under alternate furrow and check basin irrigation with canal and tube well water in Punjab, India. *Irrig. Sci.*, 2010; 28(6): 489–96.
- Vishuddha N, Singh GR, Kumar R, Raj S and Yadav B. Effect of irrigation levels and nutrient sources on growth and yield of wheat (*Triticum aestivum* L.). *Annals of Agricultural Research* 2014; 35(1):14-20.
- Wairagade MN, Choudhary AA, Mairan NR and Kirnapure VS Effect of moisture conservation practices under limited irrigation on yield and economics of wheat (*Triticum aestivum* L.). *Journal of Pharmacognosy and Photochemistry*. 2020; 9(5): 1853-1856.
- Zabihullah. Assessment of nitrogen and phosphorus requirement of timely sown bed planted wheat under varying irrigation levels. MSc Thesis CCS Haryana Agricultural University, Hisar, India. 2020.