"THE EFFECT OF ORGANIC MANURES AND BIOFERTILIZERS ON GROWTH, FLOWERING AND YIELD OF GLADIOLUS *(Gladiolus grandiflorus)* cv. PusaSrijana"

ABSTRACT

The experiment was conducted on "The effect of organic manures and biofertilizers on growth, flowering and yield of gladiolus *(Gladiolus grandiflorus)* cv. PusaSrijana" conducted during the period of October,2021 to March, 2022 at Horticulture Research Farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom, University of Agriculture, Technology and Sciences, Prayagraj (Uttar Pradesh). The experiment was planned in randomized block design with three replications and eight treatments. The observations were recorded on various growth, flowering and yield contributing characters. Based on the results obtained from the present investigation, it is concluded that the T_7(RDF + Vermicompost + Azotobacter @ 5Kg/ha+ PSB @ 5Kg/ha) was found superior followed by T_6 (RDF + Vermicompost + Azotobacter @ 4.5Kg/ha+ PSB @ 4.5Kg/ha) and T_5(RDF + Vermicompost + Azotobacter @ 4Kg/ha+ PSB @ 4Kg/ha) in growth and flowering with higher yield factors. In this investigation the T_7 was found most suitable for cultivation for better yield per hectare of corms, cormels and spike and best returns in terms of economics of the crop with high net returns and Benefit cost ratio.

INTRODUCTION

The total world area under bulbous crops is around 50,000 ha out of which over 3500 ha is in India (Raj, 2005).Gladiolus stands 4th in the international cut flower trade after rose, carnation andchrysanthemum. The flowers are of various colours.Gladiolus can be cultivated on all types of soil having good structure and drainage.A soil pH between 6 and 7 is ideal; however, a soil with pH ranging from 5 to 7 can also be used for gladiolus cultivation. In soil by adding organic manures and microbial agents make easy uptake of nutrients when crop required comparing to chemical fertilizers (Vanilarasu and Balakrishnamurthy, 2014).The beneficial effect of Azotobacter is attributed to its N fixing capacity (15-30 kg/ha) and ability to produce growth promoting substances like auxins, gibberellins and cytokinins (Azcon and Barea, 1975).Phosphate Solubilizing Bacteria are useful for all the crops i.e., cereals, cash crops, leguminous crops, horticultural crops, vegetables, flowers etc. The effective strain of Phosphate Solubilized Bacteria increased the level of available P□O5 in the soil. With the increase in available P□O5 level, overall plant growth can be increased. In certain condition they also exhibit anti-fungal activities and thereby fungal diseases may be controlled indirectly.VarietyPusaSrijana is a gladiolus hybrid variety released from Indian Agricultural Research Institute, New Delhi and is a cross between Berlew × Heady Wine. PusaSrijana attains

Formatted: Justified

UNDER PEER REVIEW

a height of 103.22 cm having spike length of 85.25 cm, rachis length of 49.55 cm with 16.66 number of florets per spike.

MATERIAL AND METHODS

Experimental site

The present investigation "The effect of organic manures and biofertilizers on growth, flowering and yield of Gladiolus *(Gladiolus grandiflorus)* cv. PusaSrijana" was conducted during October,2021-March,2022 at Horticulture Research Farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom, University of Agriculture, Technology and Sciences, Prayagraj (Uttar Pradesh). All the facilities necessary for cultivation, including labour were made available in the department. Prayagraj is situated at an elevation of 78 meters above sea level at 25.87˚ North latitude and 81.15˚ E longitudes. This region has a sub-tropical climate prevailing in the South-East part of U.P. with both the extremes in temperature, i.e., the winter and the summer. In cold winters, the temperature sometimes is as low as 0˚C in December – January and very hot summer with temperature reaching up to 46˚C in the months of May and June. During winter, frosts and during summer, hot scorching winds are also not uncommon. The average rainfall is around 1013.4 (cm) with maximum concentration during July to September months with occasional showers in winters. The meteorological data for the experimental period collected from Meteorological Observatory at College of Forestry and Environment, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj.

Formatted: Justified

Chart 1 :Treatment details

Treatment	Treatment details
T_0	RDF only
T_1	Vermicompost + azotobacter @2kg/ha + PSB @2kg/ha
T_2	Vermicompost + azotobacter @2.5kg/ha + PSB @2.5kg/ha
T_3	Vermicompost + azotobacter @3kg/ha + PSB @3kg/ha
T_4	Vermicompost + azotobacter @3.5kg/ha + PSB @3.5kg/ha
T_5	Vermicompost + azotobacter @4kg/ha + PSB @4kg/ha
T_6	Vermicompost+ azotobacter @4.5kg/ha + PSB @4.5kg/ha
T_7	Vermicompost +azotobacter @5kg/ha + PSB @5kg/ha

* RDF(150kg : 120kg : 120kg per ha) is common in all the treatments with Vermicompost (8t/ha)

UNDER PEER REVIEW

Table 1 Growth parameters:

Treatments	Sprouting (no of days)	Sprouting %	Days to 7 leaf stage	Plant height 30 days (cm)	Plant height 45 days (cm)	Plant height 75 days (cm)
T0	14.22	85	63.53	33.56	52.53	76.06
T_1	14.17	88.39	58.16	38.63	59.96	80.36
T_2	13.87	88.72	57.33	43.63	66.33	83.36
T_3	13.07	90.13	56.65	41.7	67.16	85.86
T_4	11.59	91.84	56.26	45.13	67.53	85.86
T_5	11.26	93.98	51.14	46.13	70.16	94.83
T_6	9.2	96.14	48.4	46.63	72.2	96.33
T_7	7	97.14	44.76	52.86	78.3	103.23
SEM±	0.31	0.43	1.02	2.61	2.96	2.66
CD (5%)	0.92	1.26	3.00	7.65	8.65	7.77
CV	4.60	0.82	3.25	10.4	7.67	5.21

UNDER PEER REVIEW

Table2 Floral parameters:

Treatments	50% flowering	No. of florets/spike	Diameter of 1st flower(cm)	Diameter of 2nd flower(cm)	Spike length (cm)	Spike weight (gm)	Shelf life in days
T_0	121.62	7.58	8.07	7.43	64.4	60.16	5.66
T_1	121	10.1	8.17	7.56	66.76	62.4	6.44
T_2	120.66	13.95	8.37	8	75.76	63.13	7.28
T_3	116	10.89	9.43	8.53	77.86	69.23	9.32
T_4	114.66	11.69	10.2	8.66	76.96	64.36	10.14
T_5	113.36	12.52	10.53	9.13	81.46	75.8	12.83
T_6	112.33	13.79	10.77	9.4	84.26	83.73	14.29
T_7	108	15.39	11.13	9.7	84.96	84.16	14.79
SEM±	0.51	0.28	0.53	0.39	3.44	4.51	0.20
CD (5%)	1.48	0.82	1.56	1.13	10.06	13.20	0.59
CV	0.75	4.06	9.66	7.80	7.78	11.11	3.47

UNDER PEER REVIEW

Table 3 Yield attributing characters:

Treatments	No. corms/plant	Weight of corms/plant (gm)	No. of cormels/plant	Weight of cormels/plant (gm)
T0	1.33	30.4	19.69	12.65
T_1	1.77	31.71	21.08	12.71
T_2	2.2	33.48	21.85	15.00
T_3	2.23	44.05	22.77	16.05
T_4	2.42	46.00	24.12	17.25
T_5	2.46	46.95	25.19	18.75
T_6	2.79	49.32	26.21	19.80
T_7	3.00	52.00	26.45	21.32
SEM±	0.13	0.39	0.19	0.23
CD (5%)	0.38	1.15	0.57	0.66
CV	9.92	1.68	1.44	2.35

UNDER PEER REVIEW

Table 4 Yield parameters:

Treatment	No. of corms/ha	Yield of cormels (q/ha)	No. of spikes/ha
T_0	126112.5	11.94	94444.35
T_1	174601	12.48	98211.01
T_2	217138.1	14.79	98585.09
T_3	223784	16.08	100151.8
T_4	247689.4	17.61	102048
T_5	257816.7	19.59	104422.1
T_6	298072.8	21.15	106822.1
T_7	324293.2	23.02	107940.6
SEM±	13009.75	0.25	479.62
CD (5%)	38060.32	0.73	1403.14
CV	9.86	2.52	0.82

Results and discussion:

Growth parameters:

In terms of number of days to sprouting, themaximum days to sprouting (14.22) was recorded in the treatment T_0 and minimum days to sprouting (7.0) was found for the treatment T_7. In terms of sprouting % the Maximum sprouting% (97.14) was recorded in the treatment T_7 and minimum sprouting% (85) was found for the treatment T_0. In terms of 7 leaf stage the Maximum days taken to 7 leaf stage (63.53) was recorded in the treatment T_0 and minimum days taken to 7 leaf stage (44.76) was found for the treatment T_7. In terms of plant height recorded (30DAS) highest was recorded in T_7 (52.86cm) whereas lowest was recorded in treatment T_0 (33.56 cm). Plant height (45 DAS) was recorded highest in T_7 (78.3cm) whereas lowest in T_0 (52.53 cm). Plant height (75DAS) highest was recorded inT_7 (103.23cm) whereas lowest in T_0 (76.06cm) respectively.

Formatted: Justified

UNDER PEER REVIEW

Floral parameters:

In terms of days to 50 % flowering the maximum days to 50% flowering (121.62) was recorded in the treatment $T_0$and minimum days (108) was found for the treatment T_7. In terms of no of florets/spike the maximum no of florets/spike (15.39) was recorded in the treatment T_7 and minimum florets (7.58) was found for the treatment T_0. In terms of diameter of 1st& 2nd flower the maximum diameter (11.13cm & 9.7cm) was recorded in the treatment T_7 and minimum diameter of 1st& 2nd flower (8.07cm & 7.43 cm) was found for the treatment T_0. In terms of spike length the maximum (84.96cm) was recorded in the treatment T_7 and minimum length (64.4cm) was found for the treatment T_0. In terms of spike weight the maximum (84.16gm) was recorded in the treatment T_7 and minimum weight(60.16gm) was found for the treatment T_0respectively.

Shelf life:

In terms of shelf lifemaximum shelf life(14.79 days) was recorded in the treatment T_7 and minimum shelf life(5.66 days) was found for the treatment T_0 respectively.

Yield attributing characters:

In terms of no of corms/plant maximum (3) was recorded in the treatment T_7 and minimum no. of corms/plant(1.33) was found for the treatment T_0. In terms of weight of corms/plant maximum weight of corms (52.0 gm) was recorded in the treatment T_7 and minimum weight of corms/plant(30.4 gm) was found for the treatment T_0. In terms of no. of cormels/plant maximum (26.45) was recorded in the treatment T_7 and minimum no. of cormels/plant(19.69) was found for the treatment T_0. In terms of weight of cormels/plant maximum (21.32 gm) was recorded in the treatment T_7 and minimum weight of cormels/plant(12.65 gm) was found for the treatment T_0 respectively.

Yield parameters:

In terms of no. of corms/hectare maximum (324293.2) was recorded in the treatment T_7 and minimum no. of corms/hectare (126112.5) was found for the treatment T_0. In terms of yield of cormels (q/ha) maximum (23.02 q/ha) was recorded in the treatment T_7 and minimum yield of cormels (q/ha)(11.94 q/ha) was found for the treatment T_0. In terms of no. of spikes/hectare

Formatted: Justified

Formatted: Justified

Formatted: Justified

Formatted: Justified

UNDER PEER REVIEW

(107940.6) was recorded in the treatment T_7 and minimum no. of spikes/hectare (94444.35) was found for the treatment T_0 respectively.

Economics:

Economically best treatment was T_7, gave the highest net return up to 2026917.00 Rs/ha as well as B:C ratio 4.37, and T_0 lowest net return was at 864101.00 Rs/ha as well as B:C ratio 2.43.

Conclusion:

In this study, T_7 was found superior combination in days to sprouting,50% sprouting, days taken to 7 leaf stage, maximum plant height (30,45,75DAS), 50% flowering, florets/spike, flower diameter, length & weight of spike, shelf life, no of corms, weight of corms, no of cormels, weight of cormelswith high net return and B:C ratio. From this it can be concluded that the T_7, was the most promising treatment amongst all the other treatments.

Formatted: Justified

Formatted: Justified

UNDER PEER REVIEW

REFERENCES:

MukeshDawar, Anuj Kumar, M VidhyaSankar, Roshan Gallani (2019). Effect of biofertilizers on growth and flowering of tuberose (Polianthes tuberosa L.) under Malwa Plateau of Madhya Pradesh. *Journal of Ornamental Horticulture 22 (1and2), 21-27, 2019*

E. Sathyanarayana, SudhaPatil, M. Bahubali and S. L. Chawla (2016). Effect of INM on gladiolus (Gladiolus grandiflorus L.) cv. American Beauty under Navsari and Tansa Conditions, *International journal of pure and applied bioscienceISSN: 2320 – 7051*

P. B. Pansuriya, R. V. Chauhan, J. V. Varasani and S. R. Aghera (2016). Effect of INM on flowering, corm and cormel yield of gladiolus (gladiolus grandiflorus L.) cv. Psittacinus hybrid, *An international quarterly journal of life sciences.*

Ali, A., Mehmood, T., Hussain, R., Bashir, A., Raja, S., Najam-ud-Din and Ahmad, A.(2014). Investigation of biofertilizers influence on vegetative growth, flower quality, bulb yield and nutrient uptake in gladiolus (*Gladiolus grandiflorus L*). *International Journal of Plant, Animal and Environmental Sciences 4(1): 94-99.*

Baskaran, V., Misra, R.L. Singh, S.K. and Abirami(2014). Response of biofertilizers and commercial formulations on growth, yield and comm production of gladiolus. *Indian Journal of Horticulture. 71(2): 237-241.p*

Rahul Singh, Mukesh Kumar, Sameeksha Raj and Sanjay Kumar(2014). Flowering and corm production in gladiolus (gladiolus grandiflorus l.) cv.“White prosperity” as influenced by integrated nutrient management (INM). Annals of Horticulture 7(1) : 36-42 (2014).

Sharma Uday Beer, Khare R. K (2014). To study the effect of bio fertilizer and foliar spray of zinc under different NP levels on floral characteristics and economics of Gladiolus. *BhartiyaKrishiAnusandhanPatrikaYear : 2014, Volume : 29, Issue : 2 Print ISSN : 0303-3821. Online ISSN : 0976-4631.*

UNDER PEER REVIEW

Hadwani, M. K. ; Varu, D. K. ; NiketaPanjiar ; Babariya, V. J.(2013). Effect of integrated nutrient management on growth, yield and quality of ratoon tuberose (*Polianthes tuberose* L.) cv. Double.*Asian Journal of Horticulture Vol.8*

DhamiVandana, Rao V.K, Sachan Sanjay, Kumar Santosh (2013). To study the influence of biofertilizers, organic manures and chemical fertilizers on growth, flowering and yield of African marigold African marigold (Tageteserecta L.) cv. PusaNarangiGainda. *Journal of Ornamental Horticulture Volume: 16*

Chaudhuri, S.R., Patil, A.B. and Patel, N.K. (2013). Effect of organics, inorganics and biofertilizers on growth and yield of gladiolus (Gladiolus grandiflorus L.) cv. AmericanBeauty. *Bioinfolet, 10(4 B): 1214-1217.*

Dubey R.K., Misra R.L., Singh S.K., Manisha (2010). Efficacy of bio-and chemical fertilizers on certain floral qualities of gladiolus. *Indian Journal of Horticulture Year : 2010, Volume : 67*

Uday Sharma ; Chaudhary, S. V. S. ; Rajesh Thakur (2008). Response of gladiolus to integrated plant nutrient management. *Haryana Journal of Horticultural Sciences 2008 Vol.37*

R. Srivastava, M. Govil (2007). Influence of biofertilizers on growth and flowering in gladiolus cv. American beauty. *International Society for Horticultural Sciences 2007.*

Rakesh Kumar, Ramesh Kumar and Prabhat Kumar (2007). Effect of integrated use of chemical fertilizers, biofertilizers and biostimulants in gladiolus (*Gladiolus grandiflorus L.*) cv. Sancerre. *Progressive Horticulture Vol. 43(1).*

Dongardive, S. B. ; Golliwar, V. J. ; Bhongle, S. A. (2007). Effect of organic manure and biofertilizers on growth and flowering in *Gladiolus* cv. White prosperity. *Plant Archives 2007 Vol.7.*

Gotmare, P. T. ; Damke, M. M. ; Gonge, V. S. ; SnehalDeshmukh (2007). Influence of integrated nutrient management on vegetative growth parameters of marigold (*Tageteserecta* L.). *Asian Journal of Horticulture 2007 Vol.2*

Godse, S. B. ; Golliwar, V. J. ; Neha Chopde ; Bramhankar, K. S. ; Kore, M. S.(2006). Effect of organic manures and biofertilizers with reduced doses of inorganic fertilizers on growth, yield and quality of gladiolus. *Journal of Soils and Crops 2006 Vol.16.*

Mogal, S. A. ; Khiratkar, S. D. ; Chopde, N. K. ; Dalvi, A. M. ; Kuchanwar, O. D. ; Khobragade, Y. R., (2006). Effect of organic manures and biofertilizers with reduced doses of nitrogen on growth, yield and quality of China aster**.***Journal of Soils and Crops 2006 Vol.16*

Mukeshkumar, Sultan Singh, S.K. Sharma, D.S. Dahiya, L.S. Beniwal (2006). Effect of biofertilizers on growth and flowering of marigold cv. PusaNarangi. *Haryana Journal Horticulture science 36 (1&2).*

UNDER PEER R