

Effect Of Npk And Nano Fertilizers On Plant Growth, Yield And Quality Of Gladiolus (*Gladiolus Grandiflorus L.*)

ABSTRACT:

The current experiment was conducted in the Research Field of the Department of Horticulture, SHUATS, Prayagraj, from November 2021 to March 2022. The goal of the experiment, which used a Randomized Block Design (RBD) with eight treatments and three replications, was to determine which variety will grow best in Prayagraj agroclimatic conditions. It is determined from the current experimental results that the eight treatments in the American beauty variety under consideration demonstrated substantial diversity in all the parameters noted. to investigate "~~The Effect of NPK and Nano Fertilizer on Growth, Yield, and Quality of Gladiolus (Gladiolus grandiflorus L.)~~". The best growth (116.22), number of leaves (14.0), number of shoots 50% sprouting (19.73 cm), spike length (72.33 cm), and first spike emergence were all recorded (86.77 cm), days it took for the first flower to bloom (118.44 cm) 50% florets (106.88 cm), spike weight (12.77 cm), and first and second floret flower diameter (6.27, 6.01 cm) Corms per plant (2.33 cm), corms per plot (20.31 cm), vase life (11.5 cm), spikes (1.83 cm), and florets (number of) (12.77cm) compared to other treatments, T5 nano iron 0.25g/l offered the best results in terms of gladiolus growth, yield, and quality. The highest T5 nano iron 0.25g/l net income was (1563017) Rs./Ha, the highest gross return was (2187165) Rs./Ha, and the highest BC ratio was (1:2.5).

Key words: *Gladiolus, T5 nano iron, 0.25g/l, treatment effectiveness*

1. INTRODUCTION

The Latin word "Gladius," which means sword, is where the name "gladiolus" originates. It goes by the names "com flag" or "sword lily." Due to the presence of *Gladiolus illyricum* as a wild weed in com fields, **com flag** has another name throughout Europe. **com flag** came up with the term "gladiolus" to characterize the leaves' sword-like design. The gladiolus, or *Gladiolus grandiflorus*, is a well-known flowering plant from the Iridaceae family and subfamily Ixodidae. There are roughly 180 species in the genus *Gladiolus*, and there are more than 10,000 cultivars, of which about 20 are bred for cut flowers on a commercial scale. Many others are also employed in gardens, exhibitions, and other spaces **as** seasonal flowering plants.

It is found throughout Africa and **Mediterranean** region with its greatest concentration in South Africa. *Gladiolus* (*Gladiolus grandiflorus*) is said to be the "Queen of bulbous plant" because the gladiolus is top in the list of its beauty, glamour keeping quality various colour like white, crimson, pink, orange, salmon, red, purple cream and rose etc. shades and shapes.

Gladiolus is grown under varying climatic **condition**. It is cultivated during **rainy** and winter season for obtaining good quality cut flowers and corms of gladiolus. It grows well in good drained sandy loam soil, with sufficient organic matter. Its cultivation start from June-December. Generally gladiolus is planted from corms. The plant grows to a height of about 80 to 90 cm. The flowers are irregular and funnel shaped. They are boom on branched spikes. **Ilapogu, V. 2012.[1]**. The plant **start** flowering from 60 to 90 days after **planting** of conns. The spikes remain fresh for about 10 to 12 days depending upon the season. The gladiolus cut flowers stands well for transportation **to** long-distance. Hence, **its cultivation for** cut flowers can be taken up in far off areas.

Iron NPs due to their nano size as well as magnetic characteristics are considered as special nano fertilizers. The bio-compatibility **as well as interaction between plants and the Fe nanoparticles had lead to a great deal of attentions** According to the records, the Fe nanoparticles effects on plants are of two kinds like concentrations in lower doses of Fe NPs were found to have positive effects on the growth and physiology of crop plants, whereas high concentrations of Fe NPs application seems have resulted in toxic effects on plants (harmful). **Tayade et al. (2018).[2]**. Fe nanoparticles were reported with nutrient absorption promotion as well as photosynthetic efficiency enhancement whereas the information on anatomical as well as ultra-structural responses at the cellular level of plants under Fe NPs application is still lacking When Fe nanoparticles are absorbed by plants, their absorption, distribution, and accumulation have also attracted significantly higher interest when used as fertilizers in agricultural.

The higher rate of phosphorus and potassium tended to improve the flower quality, cormel growth and corm production. The application of potassium plays vital role in regulating the opening and closing stomata and water retention **Hembrom, R. and Singh, A.[3]**. Potassium has works catalyst of many biosynthetic reactions of the photosynthesis and there by assumed to affect the biomass production. **Barman D, Rajni K, Ram P.[4]**. The deficiency of potassium in gladiolus reduces the number of floret in there spike, shortens the spikes length and delay the flowering. Lack of potassium causes the less flowering of gladiolus

2. MATERIALS AND METHODS

The field experiment entitled “**Effect of NPK and Nano fertilizers on the growth, yield and quality of gladiolus (*Gladiolus grandiflorus* L.)” cv.-American Beauty**” was carried out during the year 2021-2022 at Sam Higginbottom University of Agriculture Technology And Sciences Allahabad.

2.1 Geographical location of the experimental site:

The experimental site is ~~being~~ located at a latitude of 25.41° North and **longitude** of 81.84° East, with an altitude of 98 meters above the mean sea level (MSL).

2.2 Climatic conditions in the experimental area:

The area of Prayagraj comes under **humid** sub-tropical climate, which experiences warm humid monsoon, hot dry summer and cold dry winter. The annual mean temperature is 26.1°C while monthly mean temperatures are 18-29°C. The daily average maximum temperature is about 22°C and the minimum temperature is 9°C. The average annual rainfall received is 1042.2 mm. At this location, the temperature reaches upto 46°C-48°C and the minimum temperature recorded was 4°C-5°C. The relative humidity ranges in this location **ranges** between 20-94%.. The meteorological data (October2021-March2022) with respect to total rainfall, maximum and minimum temperature, relative humidity.

2.3 The experimental design and parameters determined

21. Experimental layout

Eight treatments, each replicated three times, were used in the Randomized block design (RBD) that was used to lay out the trial. Acquisition of Varieties: Only American beauty varieties were employed for this experiment. bought from the Flora Nursery in Bareilly, Uttar Pradesh.

2.3.2 Growth Conditions

Plant height (in centimeters) and the number of leaves were measured at 30, 60, and 90 days following transplanting as growth indicators. Number of shoots per plant, days needed for 50% of them to sprout, Spike size (cm),

2.3.3 Flowering conditions.

Days needed for the first spike to appear (d), 50% of the florets, and the size of the first and second florets (cm)

2.4 Statistical investigation

Separate statistical analyses of the data and outcomes were performed using the analysis of variance method (ANOVA). The least significant difference test was used to compare the treatment mean differences at 5% probability levels.

RESULTS AND DISCUSSION

3.1 Growth Parameter

Significantly, the maximum plant height was recorded in the plant height in (Table.1). The Maximum Plant height at 90 days (116.22 cm) was recorded in the T₅ (Nano Iron 0.25g/l) followed by T₆ (Nano Iron 0.30g/l) with (112.55cm) and the minimum Plant Height at 90 days (94.55cm) was recorded in. T₂ (Nano Iron 0.10g/l). Similar results were found in plants **(Ghafari and Razmjoo,2015).[5]**.

Significantly, the maximum number of leaves was recorded in the (Table.1). The Maximum no.of leaves at 90 days (10.22cm) was recorded in the T₅ (Nano Iron 0.25g/l) followed by T₆ (Nano Iron 0.30g/l) with (8.66 cm) and the minimum no.of leaves at 90 days (6.11 cm) was recorded in T₂ (Nano Iron 0.10g/l). **(Pirzad and Shokrani, 2012).[6]**

Significantly, the maximum number of shoots was recorded in the (Table.1),. The Maximum no.of shoots at 90 days (2.44cm) was recorded in the T₅ (Nano Iron 0.25g/l) followed by T₆ (Nano Iron 0.30g/l) with (2.1cm) and the minimum no.of shoots at 90 days (1.00 cm) was recorded in T₂ (Nano Iron 0.10g/l). Similar results were found in **Memon, S. A., Bangulzai.[7]**. Effect of humic acid and iron sulphate on growth and yield of Zinnia (*Zinniaelegans*). *Journal of Agriculture Technology*, **10(6):1517-1529**.

Significantly, the maximum number of days for 50% sprouting was recorded in the , at 90 DAT. The Maximum no.of days for 50% sprouting at 90 days (19.73 cm) was recorded in the T₅ (Nano Iron 0.25g/l) followed by T₆ (Nano Iron 0.30g/l) with (20.55cm) and the minimum no.of days for 50% sprouting at 90 days (25.00 cm) was recorded in T₂ (Nano Iron 0.10g/l). Similar results were found in **in Askary, M., Amirjani, M. R., and Saberi, T. 2017[8]**.

Significantly, the maximum length of spike was recorded in the, at 90 DAT. The Maximum length of spike at 90 days (72.33 cm) was recorded in the T₅ (Nano Iron 0.25g/l) followed by T₆ (Nano Iron 0.30g/l) with (66.22 cm) and the minimum length of spike at 90 days (48.33 cm) was recorded in T₂ (Nano Iron 0.10g/l). in **Saberi, T. 2017 [9]**.

3.2 Flowering parameters

Significantly, the maximum first spike emergence was recorded in the , at 90 DAT. The Maximum first spike emergence at 90 days (86.77 cm) was recorded in the T₅ (Nano Iron 0.25g/l) followed by T₆ (Nano Iron 0.30g/l) with (84.66 cm) and the minimum first spike emergence at 90 days (76.8 cm) was recorded in T₂ (Nano Iron 0.10g/l) Similar results were found in. **Tayade et al. (2018)**.

Significantly, the maximum number of days for 50% florets was recorded in the , at 90 DAT. The Maximum no.of days for 50% florets at 90 days (106.88cm) was recorded in the T₅ (Nano Iron 0.25g/l) followed by T₆ (Nano Iron 0.30g/l) with (108.55cm) and the minimum no.of days for 50% florets at 90 days (115.00 cm) was recorded in T₂ (Nano Iron 0.10g/l). 7. Similar results were found in **Ilapogu, V. 2012..**

Significantly, the maximum flower diameter of **first** and second floret was recorded in the, at 90 DAT. The Maximum flower diameter of first and second floret at 90 days (6.27, 6.01 cm) was recorded in the T₅ (Nano Iron 0.25g/l) followed by T₆ (Nano Iron 0.30g/l) with (6.04, 6.01 cm) and the minimum flower diameter of first and second floret at 90 days (5.31, 5.15 cm) was recorded in T₂ (Nano Iron 0.10g/l). Similar results were found in **Shokrani (2012).[10]**.

Table 1. Growth performance parameters of different treatments, at different days after planting (DAT)

Treatment Details		Plant height (cm)			No. of leaves			No. of shoots per plant	Days to 50% sprouting	Spike length (cm)
		30 DAP	60 DAP	90 DAP	30 DAP	60 DAP	90 DAP	90DAP	90 DAP	90 DAP
T ₁	NPK 120:150:150	71.66	101.2	37.97	2.77	6.11	9.55	1.11	21.77	55.88
T ₂	Nano Iron 0.10g/l	62.88	94.55	33.22	2.33	6.44	7.62	1.00	25.00	48.33
T ₃	Nano Iron 0.15g/l	68.77	97.11	34.94	2.66	6.88	8	1.33	24.22	50.44
T ₄	Nano Iron 0.20g/l	70.11	103.4	35.66	3.11	7.22	9.22	1.33	22.77	52.66
T ₅	Nano Iron 0.25g/l	82.55	116.2	41.77	4.55	10.22	14	2.44	19.73	72.33
T ₆	Nano Iron 0.30g/l	77.77	112.5	40.66	4.22	8.66	12.55	1.88	20.55	66.22
T ₇	Nano Iron 0.35g/l	74.88	107.6	36.11	3.44	8.33	11.27	1.55	20.77	64.00
T ₈	Nano Iron 0.40g/l	73.55	100.33	35.77	3.44	7.44	10.61	1.44	21.00	59.00
	F-Test	S	S	S	S	S	S	S	S	S
	SED(≠)	1.65	3.88	3.43	0.16	0.43	0.758	0.166	0.75	7.180
	CD@5%	3.37	7.93	7.02	0.34	0.88	1.54	0.34	1.55	14.67
	CV	5.45	6.53	4.04	6.24	6.94	8.96	13.48	4.22	15.004

Table. 2 Flowering parameters performance of different treatments of Gladiolus

Treatment Details		Days taken for first spike emergency	50% Florets	Floret Diameter (First Floret)	Floret Diameter (Second Floret)
T ₁	NPK 120:150:150	81.44	111.66	5.66	5.243
T ₂	Nano Iron 0.10g/l	82.22	115.00	5.31	5.15
T ₃	Nano Iron 0.15g/l	86.77	112.8	5.47	5.611
T ₄	Nano Iron 0.20g/l	84.66	112.5	5.65	5.38
T ₅	Nano Iron 0.25g/l	76.8	106.88	6.27	6.01
T ₆	Nano Iron 0.30g/l	78.22	108.55	6.04	5.78
T ₇	Nano Iron 0.35g/l	79.33	109.66	5.86	5.41
T ₈	Nano Iron 0.40g/l	80.55	111.00	5.68	5.54
	F-Test	S	S	S	S
	SEd (±)	1.78	0.753	0.14	0.199
	CD@5%	3.64	1.53	0.3	0.40
	CV	2.68	0.831	3.12	4.43

4. Conclusion

From the research conducted, it is concluded that among the different treatments combination as planting material gave the best performance in terms of vegetative parameter, floral parameter, yield parameter and quality parameter gave the best result in **T nano iron 0.25g/l** as compared to other treatments in **Terms** of growth, yield and quality of gladiolus recorded highest BC ratio (1:2.5) in **T nano iron 0.25g/l**

REFERENCE

1. Ilapogu, V. 2012. Studies on the effect of boron and iron on vegetative growth, flowering, corm and cormel production in gladiolus (*Gladiolus grandiflorus* L.) cultivars.[1].
2. Tayade, M., Badge, S. and Nikam, B. 2018. Foliar application of zinc and iron as influenced on flowering and quality parameters of tuberose. *International Journal of Current Microbiology and Applied Sciences*, 7(1): 2239-2243.[2]
3. Hembrom, R. and Singh, A. K. 2015. Effect of iron and zinc on growth, flowering and bulb yield in lillium. *International Journal of Agriculture, Environment and Biotechnology*, 8(1):61-64.[3].
4. Barman D, Rajni K, Ram P, Upadhyaya RC. Corm multiplication of gladiolus as influenced by application of potassium and spike removal. *J. Ornament Hort.* 2005;8(2): 104 -107. 23.[4].
5. Ghafari H, Razmjoo J. Effect of foliar application of nano-iron oxidase, iron chelate and iron sulphate rates on yield and quality of wheat. *International J. of Agronomy and Plant Production*. 2013; 4(11):2997-[5].
6. Pirzad, A. and Shokrani, F. 2012. Effects of iron application on growth characters and flower yield of *Calendula officinalis* L. under water stress. *World Applied Sciences Journal*, 18 (9): 1203-[6].
7. Memon, S. A., Bangulzai Effect of humic acid and iron sulphate on growth and yield of *Zinnia (Zinnia elegans)*. *Journal of Agriculture Technology*, 10(6):1517-1529.[7].
8. Askary, M., Amirjani, M. R., and Saberi, T. 2017. Comparison of the effects of nano- iron fertilizer with iron-chelate on growth parameters and some biochemical properties of *Catharanthus roseus*. *Journal of Plant Nutrition*, 40(7): 974- 982.[8].
9. Saberi, T. 2017. Comparison of the effects of nano- iron fertilizer with iron-chelate on growth parameters and some biochemical properties of *Catharanthus roseus*. *Journal of Plant Nutrition*, 40(7): 974- 982.[9].
10. Shokrani, F. 2012. Effects of iron application on growth characters and flower yield of *Calendula officinalis* L. under water stress. *World Applied Sciences Journal*, 18 (9): 1203.[10].