

Original Research Article

Nine years of integrated nutrient management practices on soil microbial activities in a cereal-based cropping system

ABSTRACT

The present work is a follow-up study to identify the effects of nine years long-term integrated nutrient management practices on soil microbial activities in a cereal-based cropping system which was initiated in 2010. The microbial activities like microbial population and enzyme activities were estimated at three different soil depths; 0-15cm, 15-30cm and 30-45cm, in response to different treatment regimes of inorganic fertilizers, organic manures, biofertilizers, and lime. The results identified a decreasing trend in the microbial population and enzyme activity with increased soil depth. Of all the treatment regimes, the maximum enhancements of dehydrogenase and urease activities were observed when there was a combined application of lime with inorganics and organic manure. Additionally, this treatment also increased the bacterial population while decreasing the fungal and actinomycetes population in the soil. The increased soil microbial activities in this treatment can be attributed to the role of organic manures.

Keywords: Acidic Inceptisols, cereal based cropping system, integrated nutrient management, microbial population, enzyme activity

1. INTRODUCTION

One-third of the earth's surface area and about half of the world's potentially cropped land is acidic [1]. In India, soil acidity affects around one-third of the cultivated lands. Maximizing crop yields with good soil health and proper environmental and ecological balance is possible only with the balanced use of mineral fertilizers and organic sources of nutrients such as organic manures like farmyard manure (FYM), compost, green manures, and biofertilizers [2–4]. Application of organic manures like vermicompost having C:N ratio below 15 is suitable for crop production [5].

Sustainable crop production has become a great challenge now a days, particularly in problem soils like acid soil. Long-term experiments have shown that neither organic sources nor mineral fertilizers alone can achieve sustainability in crop production. The integrated use of organic and inorganic fertilizers is more effective in maintaining higher productivity and stability [2,3,6,7]. The use of imbalanced and inadequate fertilizers and restricted use of organic manures have made the soils not only deficient in nutrients but also deteriorate soil health [8] and decreased the yield of the crop.

To supply a recommended dose of nutrients, large quantities of organic ~~materials~~ materials are needed, and also a slow release of plant nutrients upon

decomposition from organic material deprives crop growth [9–11]. Microorganisms play a very crucial role in soil fertility. They also play an important role in the degradation of organic matter and also in the detoxification of toxic wastes and pollutants. Microbial activities particularly the population of bacteria, fungi, and actinomycetes, microbial biomass carbon, and activity of enzymes such as dehydrogenase, urease, and phosphatase are the most sensitive indicators of soil health [12]. Interest in soil enzyme activity has increased recently since their activities are believed to reflect the potential capacity of soil to perform nutrient transformations. By considering the above facts, this experiment was carried out to evaluate the change in biological properties in acid *Inseptisols* in different depths.

2. MATERIALS AND METHODS

The present long-term integrated nutrient management (INM) field experiment on cereal-vegetable-pulse cropping system was performed on the farmland of "AINP on Soil Biodiversity - Biofertilizers" in the College of Agriculture, Odisha University of Agriculture and Technology, Bhubaneswar situated at 20.26°N latitude, 85.81°E longitude and 25.9 m above mean sea level. The experimental area falls under a subhumid tropical climate. The mean annual rainfall was 1577 mm, and the mean maximum and minimum temperatures were 33.2 and 21.4°C, respectively.

The soils of the site belong to *Inceptisols* order with acidic soil reaction. The experiment was initiated in the year 2010, during 2018-19 the cropping cycle was sweetcorn, knolkhol, and blackgram. The experiment was laid out in a randomized block design (RBD) having three replications with treatments consisting of; T1 (control), T2 (NPK), T3 (NPK+FYM), T4 (NPK+VC), T5 (NPK+FYM+BF), T6 (NPK+VC+BF), T7 (NPK+FYM+Lime+BF), T8 (NPK+VC+Lime+BF), T9 (1/2 NPK+BF) and T10 (uncultivated Fallow). The NPK dose was given to the crop as per the soil test-based dose viz; 150:30:60 for sweetcorn, 150:40:60 for knolkhol, and 20:20:30 for blackgram in the form of N:P₂O₅:K₂O kg ha⁻¹. Lime was applied @ 0.1 LR to sweetcorn and @0.2 LR to knolkhol and blackgram crop.

Standard methods were adopted for the analysis of soil and organic inputs [13,14]. Organic sources applied were farm yard manure (FYM) @ 5t ha⁻¹ and vermicompost (VC) @ 2.5 t ha⁻¹ to each crop. Biofertilizers (BF) like *Rhizobium* to Blackgram and *Azotobacter*, *Azospirillum* and PSB (@1:1:1) to Knolkhol and Sweetcorn. The crop residues were incorporated into the soil after harvesting the economic yield portion of each crop. Representative soil samples were collected at the end of the ninth cropping cycle after harvesting blackgram in June 2019.

The collected fresh soil samples were stored immediately in a refrigerator at 4°C for microbial parameter analysis. Microbial populations in the soil such as bacteria, fungi, and actinomycetes were enumerated using nutrient agar, rose Bengal agar, and Kenknight's agar as a growth medium, respectively, following the dilution plating viable count method [15]. After the required incubation period, the colony forming units (cfu) were counted and expressed as cfu g⁻¹ of soil. Urease activity was determined according to [16] and reported as mg of NH₄⁺ -N released g⁻¹ soil h⁻¹. Dehydrogenase activity was determined by monitoring the rate of production of triphenyl formazan [17,18]. All the data were subjected to statistical analysis with software SPSS [19] for significant differences between treatments using analysis of variance (ANOVA) at a 5% significance level.

3. RESULTS AND DISCUSSION

3.1 INFLUENCE OF NINE-YEAR LONG-TERM INM PRACTICES ON MICROBIAL POPULATION

3.1.1 INFLUENCE ON BACTERIAL POPULATION

The data relating to bacterial populations in different depths of soil has been presented in Table 1. The heterotrophic bacteria population in 0-15 cm of soil was more than 15-30cm and 30-45 cm depth soil. The heterotrophic bacteria population on the surface soil (0-15cm) was highest (63) in the integrated package where lime was applied with a soil test dose of NPK, vermicompost and biofertilizers followed by a soil test dose of NPK, farm yard manure and biofertilizers (60), NPK + VC+BF (55), NPK + FYM+ BF (52), NPK +VC (37), ½ NPK + BF (37), NPK +FYM (35), Fallow (32), Control (29) and lowest was in completely inorganic package NPK (25). There was no significant difference (p=0.05) between vermicompost and

farm yard manure applied packages. In 15-30 cm depth, the bacterial population was more than 30-45 cm soil but less than 0-15 cm depth soil. The heterotrophic bacteria population in 15-30 cm depth varied between 14 and 50. The highest was in T8 (50) followed by T7 (46), T6 (42), T5 (38), T4 (29), T9 and T3 (25), T10 (21), T1(16), and the lowest was in T2 (14).

The bacterial population in the subsurface layer (15-30 cm) were significantly different among themselves ($p=0.05$) except for control and NPK packages where these two packages were statistically at par ($p=0.05$). The bacterial population in the deeper layer (30-45 cm) was ranged between $10-14 \times 10^5$ cfu g^{-1} soil and it was nonsignificant among the packages. In the vermicompost applied packages, the bacterial population were more than in farm yard manure applied packages due to more readily available nutrient for microbes. The addition of biofertilizer with organics enhances the bacterial population, and ameliorants like lime improved the bacterial population. The liming neutralizes the soil acidity, enhances the nutrient availability to the plant, and thus provides a congenial environment to the microbes [20]. The biofertilizers containing rhizobium secretes exopolysaccharide, phytohormone like indole acetic acid [21], and other metabolites which enhance the microbial population in the rhizosphere [22]. Integrated nutrient management with soil amelioration enhances the biological properties [12]. The application of only inorganic nutrients depletes the soil reaction (pH) [23] which resulted in the depletion of the bacterial population.

Table 1: Influence of nine years of INM practice on heterotrophic bacterial population ($\times 10^5$ cfu g^{-1} soil)

Treatments	Depth of Soil (cm)		
	0-15	15-30	30-45
T1: Control	29	16	11
T2: NPK	25	14	10
T3: NPK+FYM	35	25	11
T4: NPK+VC	37	29	12
T5: NPK+FYM+BF	52	38	12
T6: NPK+VC+BF	55	42	13
T7: NPK+FYM+Lime+BF	60	46	14
T8: NPK+VC+Lime+BF	63	50	14
T9: 1/2 NPK+BF	37	25	11
T10: Fallow	32	21	10
LSD (5%)	7.4	4.5	NS

3.1.2 INFLUENCE ON FUNGAL POPULATION

The data related to the influence of long-term manurial practice on soil fungal populations in different depths of soil has been presented in Table 2. The surface soil (0-15 cm) had a maximum fungal population than sub-surface (15-30 cm) and deeper soil (30-45 cm). The fungal population at 0-15 cm, 15-30 and 30-45 cm depth soil was varied between 22×10^4 cfu g^{-1} soil and 35×10^4 cfu g^{-1} soil; 12×10^4 cfu g^{-1} soil and 22×10^4 cfu g^{-1} soil; and 9×10^4 cfu g^{-1} soil and 12×10^4 cfu g^{-1} soil, respectively. In 0-15 cm layer of soil, the highest was observed in NPK+FYM+BF (35×10^4 cfu g^{-1} soil) followed by NPK+VC+BF (33×10^4 cfu g^{-1} soil), NPK+FYM (30×10^4 cfu g^{-1} soil), NPK + VC (30×10^4 cfu g^{-1} soil), NPK (27×10^4 cfu g^{-1} soil), NPK + FYM +Lime + BF (25×10^4 cfu g^{-1} soil), NPK + VC+ Lime +BF and control (22×10^4 cfu g^{-1} soil), 1/2 NPK + BF (20×10^4 cfu g^{-1} soil), and lowest was in fallow (18×10^4 cfu g^{-1} soil). The application of lime reduced the fungal population, it was due to the reduction of soil acidity as fungus can grow in a wide range of pH [24,25]. The fungal population was more in

FYM applied packages than VC applied packages. In 15- 30 cm soil the fungal population was highest ($22 \times 10^4 \text{cfu g}^{-1}$ soil) in T5 followed by T3 ($20 \times 10^4 \text{cfu g}^{-1}$ soil), T6 and T4 ($19 \times 10^4 \text{cfu g}^{-1}$ soil), T9 ($16 \times 10^4 \text{cfu g}^{-1}$ soil), T7 and T2 ($15 \times 10^4 \text{cfu g}^{-1}$ soil), T10 ($13 \times 10^4 \text{cfu g}^{-1}$ soil), T1 ($12 \times 10^4 \text{cfu g}^{-1}$ soil) and the lowest was in T8 ($11 \times 10^4 \text{cfu g}^{-1}$ soil). In 30-45 cm soil the fungal population was nonsignificant ($p=0.05$) with each package.

Table 2: Influence of nine years of INM practice on fungal population ($\times 10^4 \text{cfu g}^{-1}$ soil)

Treatments	Depth of Soil (cm)		
	0-15	15-30	30-45
T1: Control	22	12	9
T2: NPK	27	15	10
T3: NPK+FYM	32	20	12
T4: NPK+VC	30	19	10
T5: NPK+FYM+BF	35	22	12
T6: NPK+VC+BF	33	19	10
T7: NPK+FYM+Lime+BF	25	15	10
T8: NPK+VC+Lime+BF	22	11	9
T9: 1/2 NPK+BF	20	16	11
T10: Fallow	18	13	10
LSD (5%)	4.3	2.3	NS

3.1.3 INFLUENCE ON ACTINOMYCETES POPULATION

The data related to the actinomycetes population of soil in different soil depth has been presented in Table 3. The actinomycetes population in 0-15 cm soil layer ranged from $10 \times 10^4 \text{cfu g}^{-1}$ soil to $22 \times 10^4 \text{cfu g}^{-1}$ soil. The highest was in the package where soil test-based fertilizer was applied with farm yard manure and biofertilizers (T5) followed by package where soil test-based fertilizer was applied with vermicompost and biofertilizers (T6), NPK + FYM (T3), NPK + VC (T4) and $\frac{1}{2}$ NPK + BF (T9), fallow (T10), NPK+ FYM +Lime+ BF (T7) and control (T1), NPK+ VC+ Lime+ BF (T8) and lowest was in NPK (T2). In 15-30 cm depth of soil the actinomycetes population was varied between $6 \times 10^4 \text{cfu g}^{-1}$ soil and $15 \times 10^4 \text{cfu g}^{-1}$ soil. The actinomycetes population in 30-45 cm depth of soil was ranged from $4 \times 10^4 \text{cfu g}^{-1}$ soil to $7 \times 10^4 \text{cfu g}^{-1}$ soil. The actinomycetes population was nonsignificant ($p=0.05$) among the packages. The data reveals that the actinomycetes population was influenced positively by FYM application and negatively with liming of acid soil. Similar finding was reported by [26] in rhizosphere of red gram and [12] in rice-rice ecosystem.

Table 3: Influence of nine years of INM practice on actinomycetes ($\times 10^4 \text{cfu g}^{-1}$ soil) population

Treatments	Depth of Soil (cm)		
	0-15	15-30	30-45
T1: Control	12	8	5
T2: NPK	10	6	4
T3: NPK+FYM	18	12	5
T4: NPK+VC	15	10	5

T5: NPK+FYM+BF	22	15	7
T6: NPK+VC+BF	19	12	6
T7: NPK+FYM+Lime+BF	12	7	6
T8: NPK+VC+Lime+BF	10	6	5
T9: 1/2 NPK+BF	15	10	6
T10: Fallow	13	8	5
LSD (5%)	2.3	1.6	NS

3.2 INFLUENCE OF NINE YEARS OF INM PRACTICES ON SOIL ENZYME ACTIVITIES

3.2.1 INFLUENCE ON SOIL DEHYDROGENASE ACTIVITY

The dehydrogenase activity of the soil was presented in Fig 1. The activity of dehydrogenase enzyme in 0-15 cm soil depth was varied between 20 $\mu\text{g TPF g}^{-1}\text{dw}$ to 69 $\mu\text{g TPF g}^{-1}\text{dw}$. The highest was estimated in the integrated package where soil test dose of NPK along with vermicompost, lime and biofertilizer was applied (69 $\mu\text{g TPF g}^{-1}\text{dw}$) followed by the package where soil test dose of NPK along with farmyard manure, lime and biofertilizer was applied (60 $\mu\text{g TPF g}^{-1}\text{dw}$), soil test dose of NPK with vermicompost and biofertilizer (54 $\mu\text{g TPF g}^{-1}\text{dw}$), soil test dose of NPK with farmyard manure and biofertilizer (48 $\mu\text{g TPF g}^{-1}\text{dw}$), soil test dose of NPK along with vermicompost was applied (40 $\mu\text{g TPF g}^{-1}\text{dw}$), $\frac{1}{2}$ of soil test dose of NPK with biofertilizer (35 $\mu\text{g TPF g}^{-1}\text{dw}$), soil test dose of NPK with farmyard manure (33 $\mu\text{g TPF g}^{-1}\text{dw}$), uncultivated fallow (28 $\mu\text{g TPF g}^{-1}\text{dw}$), control (23 $\mu\text{g TPF g}^{-1}\text{dw}$) and lowest was in soil test dose of NPK (20 $\mu\text{g TPF g}^{-1}\text{dw}$).

Fig 1. Influence of nine years of INM practice on soil dehydrogenase activity ($\mu\text{g TPF g}^{-1}\text{dw}$)

In 15-30 cm depth the dehydrogenase activity was highest in T8 (55 $\mu\text{g TPF g}^{-1}\text{dw}$) followed by TT7 (52 $\mu\text{g TPF g}^{-1}\text{dw}$), T6 (46 $\mu\text{g TPF g}^{-1}\text{dw}$), T5 (42 $\mu\text{g TPF g}^{-1}\text{dw}$), T4 (33 $\mu\text{g TPF g}^{-1}\text{dw}$), T9 (30 $\mu\text{g TPF g}^{-1}\text{dw}$), T3 (27 $\mu\text{g TPF g}^{-1}\text{dw}$), T10 (22 $\mu\text{g TPF g}^{-1}\text{dw}$), T2 (19 $\mu\text{g TPF g}^{-1}\text{dw}$), and lowest was in control (17 $\mu\text{g TPF g}^{-1}\text{dw}$). The dehydrogenase activity of soil in 30-45 cm depth was varied between 12 $\mu\text{g TPF g}^{-1}\text{dw}$ and 17 $\mu\text{g TPF g}^{-1}\text{dw}$. The dehydrogenase activity was nonsignificant among the treatments. The dehydrogenase activity was decreased with increase in depth of soil. Liming of acid soil increased the dehydrogenase activity. The dehydrogenase activity was more in organic applied packages than inorganic packages. The application of organics increased the dehydrogenase activity

due to degradation of added materials which provides intra and extracellular enzymes and increase microbial activity in the soil [27,28]. The dehydrogenase activity of organic management soil was more than conventional management [29,30]. Liming of acid soil enhanced the dehydrogenase activity of soil by creating a congenial environment for microbial growth [31] The finding of this experiment was also agreed with the findings of [12,32,33].

3.2.2 INFLUENCE ON SOIL UREASE ACTIVITY

The urease activity of the soil in different depth has been presented in Fig 2. The urease activity in the 0-15 cm soil depth was varied between $5 \mu\text{g NH}_4^+ \text{g}^{-1} \text{h}^{-1}$ and $15 \mu\text{g NH}_4^+ \text{g}^{-1} \text{h}^{-1}$. The highest ($15 \mu\text{g NH}_4^+ \text{g}^{-1} \text{h}^{-1}$) was in lime applied treatments irrespective of the types of organic fertilizers followed by integrated management package without lime and without biofertilizer package. The urease activity of only inorganic package ($5 \mu\text{g NH}_4^+ \text{g}^{-1} \text{h}^{-1}$) was lower than control ($7 \mu\text{g NH}_4^+ \text{g}^{-1} \text{h}^{-1}$). The urease activity of fallow ($9 \mu\text{g NH}_4^+ \text{g}^{-1} \text{h}^{-1}$) was more than control and only inorganic package but these were statistically at par ($p=0.05$). In 15-30 cm depth the urease activity was ranged from $4 \mu\text{g NH}_4^+ \text{g}^{-1} \text{h}^{-1}$ to $13 \mu\text{g NH}_4^+ \text{g}^{-1} \text{h}^{-1}$. The urease activity in 30-45 cm depth was varied between $2.3 \mu\text{g NH}_4^+ \text{g}^{-1} \text{h}^{-1}$ and $5.1 \mu\text{g NH}_4^+ \text{g}^{-1} \text{h}^{-1}$ and all the packages were statistically at par ($p=0.05$). Similar findings have been reported by [12,34].

Fig 2. Influence of nine years of INM practice on soil urease activity ($\mu\text{g NH}_4^+ \text{g}^{-1} \text{h}^{-1}$)

The manuscript lacks a scientific discussion comparing published research; therefore, I suggest incorporating the following paragraphs that would improve the scientific quality of the manuscript. I continue to add a paragraph that summarizes the importance, usefulness and social relevance, contemporary of the study, specifically pointing out the Impact, Benefit and Social Projection, something like this (for example):

Our results establish that in this studied sustainable ecosystem it depends on the nutrient flows through the trophic levels, which are mainly mediated by microorganisms [35, 36, 37]. Due to environmental degradation and poor agricultural practices, the intrinsic biodiversity of soils has been subject to numerous variations, which has led to changes in the functioning of the native microbial communities of these soils [38, 39, 40]. As a consequence, the outcome of these modifications, in terms of the ability of ecosystems to maintain ecosystem functions and services, is of fundamental importance. In this context, it is a challenge to understand and predict the mechanisms that govern the actions of soil microbial diversity and the relationship between that biodiversity and other processes that occur in it [41, 42, 43].

The functions performed by microorganisms are essential for crop growth [44]. These functions include the decomposition of OM in all its fractions, the recycling of plant material, the mobilization and immobilization of minerals and pollutants, the improvement in soil aeration, the inhibition of pathogens, the increase in resistance in plants, the physical structuring of the soil and the increase in plant nutrition promoting its growth, among others [45, 46]. For this reason, the processes in which microorganisms participate are strongly influenced by the agricultural management under which they are found [47, 48].

4. CONCLUSION

From this experiment, it can be concluded that the microbial activities viz; microbial population, enzyme activities in soil decreases with increase in soil depth under INM package of practices over nine years of cereal-based cropping. The liming of acid soil under this cropping system enhanced the heterotrophic bacteria population in soil and enzyme activities like dehydrogenase and urease whereas the fungal and actinomycetes population in soil depleted. Only inorganic application depletes the microbial activity in terms of bacterial populations and both the enzyme activities. The addition of organics like, farm yard manure and vermicompost to soil increases over all microbial activity.

REFERENCES

I suggest the authors update the bibliography; therefore I suggest adding recent references which address the issue in question. Suggested citations are for genuine scientific reasons that emphasize the current topic of study in context.

1. Von Uexküll, H.R.; Mutert, E. Global Extent, Development and Economic Impact of Acid Soils. *Plant Soil* **1995**, *171*, 1–15.
2. Priyadarshini, J.; Panda, C.M.; Sethi, D. Effect of Integrated Nutrient Management Practices on Yield, Yield Attributes and Economics of Coriander (*Coriandrum Sativum* L.). *Int. J. Curr. Microbiol. Appl. Sci.* **2017**, *6*, 1306–1312.
3. KUSUMAVATHI, K.; PATTANAYAK, S.K.; MOHAPATRA, A.K.; SETHI, D. Effect of Nutrient Management on Soil Fertility in Rice (*Oryza Sativa*)-Greengram (*Vigna Radiata*) Cropping System. *Ann. Plant Soil Res.* **2018**, *20*, 330–337.
4. Sarkar, A.K.; Pattanayak, S.K.; Surendra, S.; Mahapatra, P.; Arvind, K.; Ghosh, G.K. Integrated Nutrient Management Strategies for Acidic Soils. *Indian J. Fertil.* **2020**, *16*, 476–491.

5. Pandit, L.; Sethi, D.; Pattanayak, S.K.; Nayak, Y. Bioconversion of Lignocellulosic Organic Wastes into Nutrient Rich Vermicompost by *Eudrilus Eugeniae*. *Bioresour. Technol. Reports***2020**, *12*, 100580.
6. Mallikarjun, M.; Maity, S.K. Energetic Evaluation of Integrated Nutrient Management for Nitrogen in Kharif Rice and Its Residual Effect on Yellow Sarson. *Res. J. Agric. Sci.***2017**, *8*, 1362–1365.
7. Prusty, M.; Swain, D.; Alim, M.A.; Ray, M.; Sethi, D. Effect of Integrated Nutrient Management on Yield, Economics and Post-Harvest Soil Properties of Sweet Corn Grown under Mid-Central Table Land Zone of Odisha. *Int. J. Plant Soil Sci.***2022**, *34*, 55–61.
8. Garnaik, S.; Samant, P.K.; Mandal, M.; Mohanty, T.R.; Dwibedi, S.K.; Patra, R.K.; Mohapatra, K.K.; Wanjari, R.H.; Sethi, D.; Sena, D.R. Untangling the Effect of Soil Quality on Rice Productivity under a 16-Years Long-Term Fertilizer Experiment Using Conditional Random Forest. *Comput. Electron. Agric.***2022**, *197*, 106965.
9. Goutami, N.; Rao, C.S.; Sireesha, A.; Rao, C.P.; Vijaya Gopal, A. Effect of Long-Term Use of Inorganic Fertilizers, Organic Manures and Their Combination on Soil Properties and Enzyme Activity in Rice-Rice Cropping System. *Int. J. Curr. Microbiol. App. Sci***2018**, *7*, 469–486.
10. Sethi, D.; Subudhi, S.; Rajput, V.D.; Kusumavathi, K.; Sahoo, T.R.; Dash, S.; Mangaraj, S.; Nayak, D.K.; Pattanayak, S.K.; Minkina, T. Exploring the Role of Mycorrhizal and Rhizobium Inoculation with Organic and Inorganic Fertilizers on the Nutrient Uptake and Growth of *Acacia Mangium* Saplings in Acidic Soil. *Forests***2021**, *12*, 1657.
11. SWAIN, P.; PANDA, N.; PATTANAYAK, S.K. Effect of Long Term Integrated Nutrient Management Practices on Yield and Nutrient Uptake by Finger Millet (*Eleusine Coracana* L.) in an Acidic Inceptisols. *Ann. Plant Soil Res.***2021**, *23*, 473–476.
12. Mandal, M.; Rout, K.K.; Purohit, D.; Majhi, P.; Singh, M. Evaluation of Rice-Rice System on Grain Yield, Chemical, and Biological Properties of an Acid Inceptisols. *J. Indian Soc. Soil Sci.***2018**, *66*, 208–214.
13. Page, A.I.; Miller, R.H.; Keeny, D.R. Methods of Soil Analysis. Part II. Chemical and Microbiological Methods. *Amer. Soc. Agron., Madison, Wisconsin, USA***1982**.
14. Panda, N. Soil, Plant, Water and Seed Testing. A Text Book 2019.
15. Alexander, M. Most Probable Number Method for Microbial Populations. *Methods Soil Anal. Part 2 Chem. Microbiol. Prop.***1983**, *9*, 815–820.
16. Tabatabai, M.A.; Bremner, J.M. Assay of Urease Activity in Soils. *Soil Biol. Biochem.***1972**, *4*, 479–487.
17. Klein, D.A.; Loh, T.C.; Goulding, R.L. A Rapid Procedure to Evaluate the Dehydrogenase Activity of Soils Low in Organic Matter. *Soil Biol. Biochem.***1971**, *3*, 385–387.
18. Casida Jr, L.E.; Klein, D.A.; Santoro, T. Soil Dehydrogenase Activity. *Soil Sci.***1964**, *98*, 371–376.
19. Kirkpatrick, L.A.; Feeney, B.C. *A Simple Guide to SPSS for Windows: For Version 12.0*; Wadsworth Publishing Company, 2004; ISBN 0534610064.

20. Sethi, D.; Mohanty, S.; Pradhan, M.; Dash, S.; Das, R. Effect of LD Slag Application on Yield Attributes, Yield and Protein Content of Groundnut Kernel in Acid Soil of Bhubaneswar, Odisha. *Int. J. Farm Sci.* **2017**, *7*, 79–82.
21. Subudhi, S.; Sethi, D.; Kumar Pattanayak, S. Characterization of Rhizobium Sp (SAR-5) Isolated from Root Nodule of Acacia Mangium L. *Indian J. Biochem. Biophys.* **2020**, *57*, 327–333.
22. Sethi, D.; Mohanty, S.; Pattanayak, S.K. Effect of Different Carbon, Nitrogen and Vitamine Sources on Exopolysaccharide Production of Rhizobium Species Isolated from Root Nodule of Redgram. **2019**.
23. Pattanayak, S.K.; Sarkar, A.K. Sustainable Management of Acid Soils: Technologies and Their Transfer. *Indian J. Fertil.* **2016**, *12*, 16–35.
24. Brady, N.C.; Weil, R.R. The Nature and Properties of Soils 13th Ed Prentice Hall. *New Jersey, USA* **2002**, 249.
25. Areyu, T. A Review on the Potential Effect of Lime on Soil Properties and Crop Productivity Improvements. *J. Environ. Earth Sci.* **2019**, *9*, 17–23.
26. Sethi, D.; Mohanty, S.; Pattanayak, S.K. Acid and Salt Tolerance Behavior of Rhizobium Isolates and Their Effect on Microbial Diversity in the Rhizosphere of Redgram (Cajanus Cajan L.). *Indian J. Biochem. Biophys.* **2019**, *56*, 245–252.
27. Bhattacharyya, P.; Chakrabarti, K.; Chakraborty, A. Microbial Biomass and Enzyme Activities in Submerged Rice Soil Amended with Municipal Solid Waste Compost and Decomposed Cow Manure. *Chemosphere* **2005**, *60*, 310–318, doi:10.1016/j.chemosphere.2004.11.097.
28. Zhong, W.; Gu, T.; Wang, W.; Zhang, B.; Lin, X.; Huang, Q.; Shen, W. The Effects of Mineral Fertilizer and Organic Manure on Soil Microbial Community and Diversity. *Plant Soil* **2010**, *326*, 511–522.
29. Bhat, N.A.; Riar, A.; Ramesh, A.; Iqbal, S.; Sharma, M.P.; Sharma, S.K.; Bhullar, G.S. Soil Biological Activity Contributing to Phosphorus Availability in Vertisols under Long-Term Organic and Conventional Agricultural Management. *Front. Plant Sci.* **2017**, *8*, 1523.
30. Purohit, D.; Mandal, M.; Dash, A.; Rout, K.K.; Panda, N.; Singh, M. Influence of Long-Term Fertilization on Soil Microbial Biomass and Dehydrogenase Activity in Relation to Crop Productivity in an Acid Inceptisols. *ORYZA-An Int. J. Rice* **2019**, *56*.
31. Sethi, D.; Mohanty, S.; Dash, S. Effect of LD Slag on Soil Microbial Population and Enzyme Activity in Rhizosphere of Groundnut in Acid Soil. *Crop Res.* **2017**, *52*.
32. Nath, D.J.; Ozah, B.; Baruah, R.; Barooah, R.C.; Borah, D.K.; Gupta, M. Soil Enzymes and Microbial Biomass Carbon under Rice-Toria Sequence as Influenced by Nutrient Management. *J. Indian Soc. Soil Sci.* **2012**, *60*, 20–24.
33. Patra, A.; Sharma, V.K.; Purakayastha, T.J.; Barman, M.; Kumar, S.; Chobhe, K.A.; Chakraborty, D.; Nath, D.J.; Anil, A.S. Effect of Long-Term Integrated Nutrient Management (INM) Practices on Soil Nutrients Availability and Enzymatic Activity under Acidic Inceptisol of North-Eastern Region of India. *Commun. Soil Sci. Plant Anal.* **2020**, *51*, 1137–1149.

34. Sigua, G.C.; Stone, K.C.; Bauer, P.J.; Szogi, A.A. Efficacy of Supplemental Irrigation and Nitrogen Management on Enhancing Nitrogen Availability and Urease Activity in Soils with Sorghum Production. *Sustainability* **2020**, *12*, 8358, doi:10.3390/su12208358.
35. Olivares, B. Description of soil management in agricultural production systems of sector hammock in Anzoátegui, Venezuela. *La Granja: Revista de Ciencias de la Vida*, 2016, 23(1): 14–24. <https://doi.org/10.17163/lgr.n23.2016.02>
36. Olivares, BO., Araya-Alman, M., Acevedo-Opazo, C. et al. Relationship Between Soil Properties and Banana Productivity in the Two Main Cultivation Areas in Venezuela. *J Soil Sci Plant Nutr.*2020, 20 (3): 2512-2524. <https://doi.org/10.1007/s42729-020-00317-8>
37. Olivares, BO., Calero, J., Rey, JC., Lobo, D., Landa, BB., Gómez, JA. Correlation of banana productivity levels and soil morphological properties using regularized optimal scaling regression. *Catena*, 2022, 208, 105718. <https://doi.org/10.1016/j.catena.2021.105718>
38. Olivares, B., Paredes, F., Rey, J., Lobo, D., Galvis-Causil, S. The relationship between the normalized difference vegetation index, rainfall, and potential evapotranspiration in a banana plantation of Venezuela. *SAINS TANAH - Journal of Soil Science and Agroclimatology*, 2021, 18(1), 58-64. <http://dx.doi.org/10.20961/stjssa.v18i1.50379>
39. Olivares, B., Rey, JC., Lobo, D., Navas-Cortés, JA., Gómez, JA., Landa, BB. Fusarium Wilt of Bananas: A Review of Agro-Environmental Factors in the Venezuelan Production System Affecting Its Development. *Agronomy*, 2021,11(5):986. <https://doi.org/10.3390/agronomy11050986>
40. Olivares, B., Hernández, R. Ecoterritorial sectorization for the sustainable agricultural production of potato (*Solanum tuberosum* L.) in Carabobo, Venezuela. *Agricultural Science and Technology*, 2019,20(2): 339-354. https://doi.org/10.21930/rcta.vol20_num2_art:1462
41. Olivares, B., Hernández, R. Application of multivariate techniques in the agricultural land's aptitude in Carabobo, Venezuela. *Tropical and Subtropical Agroecosystems*, 2020,23(2):1-12. <https://n9.cl/zeedh>
42. Olivares, B. Determination of the potential influence of soil in the differentiation of productivity and in the classification of susceptible areas to banana wilt in Venezuela. *UCOPress: Spain*, pp. 89-111, 2022. <https://helvia.uco.es/handle/10396/22355>
43. Olivares, B., Hernandez, R., Arias, A, Molina, JC., Pereira, Y. Eco-territorial adaptability of tomato crops for sustainable agricultural production in Carabobo, Venezuela. *Idesia*2020, 38(2):95-102. <http://dx.doi.org/10.4067/S0718-34292020000200095>
44. Olivares, B., Hernández, R; Arias, A; Molina, JC., Pereira, Y. Identificación de zonas agroclimáticas potenciales para producción de cebolla (*Allium cepa* L.) en Carabobo, Venezuela. *Journal of the Selva Andina Biosphere*,2018, 6 (2): 70-82. http://www.scielo.org.bo/pdf/jsab/v6n2/v6n2_a03.pdf
45. Olivares, B. Hernández, R; Arias, A; Molina, JC., Pereira, Y. Zonificación agroclimática del cultivo de maíz para la sostenibilidad de la producción agrícola en Carabobo, Venezuela. *Revista Universitaria de Geografía*, 2018,27 (2): 139-159. <https://n9.cl/l2m83>
46. Olivares, B., López-Beltrán, M., Lobo-Luján, D. Cambios de usos de suelo y vegetación

Formatted: Font: Italic, Spanish (Spain, International Sort)

Formatted: Spanish (Spain, International

Formatted: Spanish (Spain, International

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Font: Italic, Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Spanish (Spain, International

Formatted: Font: Italic, Spanish (Spain, International Sort)

Formatted: Spanish (Spain, International

Formatted: Font: Italic

Formatted: English (India)

Formatted: Spanish (Spain, International

Formatted: Font: Italic, Spanish (Spain, International Sort)

Formatted: Spanish (Spain, International

Formatted: Spanish (Spain, International

Formatted: Spanish (Spain, International Sort)

en la comunidad agraria Kashaama, Anzoátegui, Venezuela: 2001-2013. *Revista Geográfica De América Central*, 2019,2(63):269-291. <https://doi.org/10.15359/rgac.63-2.10>

47. Olivares, B., López, M. Normalized Difference Vegetation Index (NDVI) applied to the agricultural indigenous territory of Kashaama, Venezuela. *UNED Research Journal*.2019.11(2): 112-121. <https://doi.org/10.22458/urj.v11i2.2299>

48. Olivares, B., Lobo, D., Cortez, A., Rodríguez, M.F. y Rey, J.C. Socio-economic characteristics and methods of agricultural production of indigenous community Kashaama, Anzoategui, Venezuela. *Rev. Fac. Agron. (LUZ)*2017.34 (2): 187-215. <https://n9.cl/p2qc5>

Formatted: Font: Italic, Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Font: Italic

Formatted: Spanish (Spain, International Sort)

Formatted: Font: Italic, Spanish (Spain, International Sort)

Formatted: Font: Italic