

## Original Research Article

**“Studies on management of  
fruit and shoot borer [*Leucinodes orbonalis* (Guenee)] using chemicals and Neem oil in brinjal  
*Solanum melongena* L.”**

### **ABSTRACT**

The experiment was conducted at the research plot of the Department of Agricultural Entomology and Central Research Farm, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, during Rabi from November to February 2021-22 in Randomized Block Design (RBD) with three replications. Eight treatments were evaluated against *Leucinodes orbonalis* i.e., Chlorantraniliprole 18.5% EC, Spinosad 45% SC, Flubendiamide 20WG, Indoxacarb 14.5% SC, Emamectin benzoate 5 SG, Lambda cyhalothrin 4.9% CS, Neem oil 0.15% EC and control in RBD with three replication. We tested to compare the efficacy against *Leucinodes orbonalis* and their influences on yield of Brinjal. The best and most economical treatment was Chlorantraniliprole (1:10.06) which was par with Spinosad (1:8.51) followed by Flubendiamide (1:8.44), Indoxacarb (1:8.04), followed by Emamectin benzoate (1:7.12) and Lambda cyhalothrin (1:6.85), Neem oil (1:6.36) as compared to control (1:4.63). The data on the percent infestation of shoot and fruit borer on brinjal 3<sup>rd</sup>, 7<sup>th</sup> and 14<sup>th</sup> day after first spray reveal that all the chemical treatments were significantly superior over control. Ratio was Chlorantraniliprole 18.5% EC (10.85%, 7.29%), Spinosad 45% SC (11.95%, 9.12%), Flubendiamide 20WG (13.76%, 9.79%), Indoxacarb 14.5% SC (14.05%, 10.47%), Emamectin benzoate 5SG (15.59%, 13.46%), Lambda cyhalothrin 4.9% CS (17.09%, 13.46%), Neem oil 0.15% EC (19.97%, 17.43%). The yields among the treatment were significant. The highest yield was recorded in Chlorantraniliprole 18.5% EC (222.50q/ha), followed by Spinosad 45% SC (198.33q/ha).

**Keywords:** Cost Benefit Ratio, Chemical Insecticide, *Leucinodes orbonalis*, Neem oil.

## INTRODUCTION

Vegetable cultivation is one of the most profitable and dynamic branches of agriculture.

Vegetables are an important constituent of the human diet. Botanically brinjal is known as (*Solanum melongena* Linnaeus) (2n=24) popularly known as eggplant belongs to family Solanaceae and India is its center of origin and diversity. Brinjal is an important dietary vegetable crop. Brinjal (*Solanum melongena* Linnaeus) also known as eggplant is referred to as the "**King of vegetables**" and now grown as a vegetable throughout the tropical, sub-tropical and warm temperate areas of the world. It is the most important vegetable in the Indian Subcontinent that accounts for almost 50% of the world's area under its cultivation. Under sustainable farming, brinjal provides regular daily income to meet the day-to-day expenditure. (Kolhe, 2017)

Under sustainable farming, brinjal provides regular daily income to meet the day-to-day expenditure. (Murugesan, 2009) It has become an important economical source for farmers and field laborers. Brinjal, *Solanum melongena* L. is one of the popular vegetable grown as a poor man's crop in India; it is also grown extensively under diverse agro-climatic conditions throughout the year (Singh, 2018).

So far, *L. orbonalis* is considered as a major pest of brinjal as a shoot and fruit borer in the established crop in the main field. The main difficulty in developing a suitable control measure against this pest is that it belongs to one of the most serious categories of insect pest internal feeder. Once the larva bores into petiole and midrib of leaves and tender shoots, it causes dead hearts. In later stages, it also bores into flower buds causing high flower abortion. They also bore into fruits also leading to fruit drop, fruit rot or rendering them unfit for human consumption when fruits are able to reach maturity.

The Larvae of this pest cause 12-16% damage to shoots and 20-60% damage to fruits. The pest is very active during the rainy and summer season and often causes more than up to 95% in India. It is also reported that the infestation of fruit borer causes a reduction in Vitamin "C" content to an extent of 68 % in the infested fruits (Anwar et al., 2015).

The yield loss in brinjal due to the pest complex is to the extent of 70-80 per cent. Among the insect pests the most obnoxious detrimental and ubiquitous pest of brinjal is shoot and fruit borer (BSFB), *L. orbonalis* (Guenee), it alone causes damage as high as 85.90% and even up to 100% damage (Kaur et al., 2014). It is also recorded as major pest of brinjal since two decades.

**Comment [KFA 11-211]:** This is already stated above  
This is better 'Brinjal is an important dietary vegetable **King of vegetables**' is grown throughout the tropical, sub-tropical and warm temperate areas of the world

**Comment [KFA 11-212]:** To whom?  
Commercial, peasant farmers or

**Comment [KFA 11-213]:** This is already stated above but attributed to a different author

**Comment [KFA 11-214]:** This is first mention so give full name

**Comment [KFA 11-215]:** Give information on how the pest is generally managed in India and indicate whether they are effective or not before coming to this statement

**Comment [KFA 11-216]:** Add reference

**Comment [KFA 11-217]:** Do you mean there is high incidence of the pest during this period

**Comment [KFA 11-218]:** Use either more than or up to and not both

Different chemicals have been reported to be effective for the control of brinjal pest complex

**Comment [KFA 11-219]:**

**Comment [KFA 11-2110]:** This should be integrated in to the preceding paragraph

UNDER PEER REVIEW

## MATERIALS AND METHODS

The experiment was conducted at Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Prayagraj (U.P.) during the rabi season of 2021-2022 with a recommended package of practices excluding plant protection. Research field situated at 25°27' North latitude 80°05' East longitude and at an altitude of 98 meters above sea level. The maximum temperature reaches up to 42°C in summer and drops down to 4°C in winter.

The Brinjal seedlings of variety 'Dhruva' were transplanted after 30 days at 60 cm x 45 cm spacing. The experiment was laid down in randomized block design (RBD) with eight treatments replicated thrice with each plot size of 2m x 2m and proper irrigation was provided. The treatments comprising of, Chlorantraniliprole 18.5% SC, Spinosad 45% SC, Flubendiamide 20 WG, Indoxiacarb 14.5% SC, Emamectin benzoate 5SG, Lambda cyhalothrin 4.9% CS and Neem oil 0.15% EC were applied two times using knapsack sprayer in 15 days of interval.

From

each plot, five plants were selected randomly and labeled tagged for recording observations. As soon as the infestation of pest on shoot was initiated, the observations on total number of shoots and number of infested shoots and fruit infestation of five observational plants from each treatment replication wise were recorded at 3<sup>rd</sup>, 7<sup>th</sup> and 14<sup>th</sup> days after imposing treatments.

The

data recorded in the different treatments were subjected to statistical analysis after suitable transformation by following standard procedures of RBD experiment. After the last picking, total of all pickings of individual plots produce were calculated to work out the yield of the treatments. Yield of healthy fruits was converted into quintal per hectare.

The observations on the per cent infestation were recorded visually per plant from five randomly selected and tagged plants in each plot. The insecticides were sprayed at recommended doses when larval population reaches its ETL level. Larvae count was taken 24 hours before spraying at 5 tagged plants per treatment, which was further converted into per plant population and subsequent observation was recorded at 3<sup>rd</sup>, 7<sup>th</sup> and 14<sup>th</sup> days after spraying on same plants.

The healthy marketable yield obtained from different treatments was collected separately and weighed. The cost of insecticides used in this experiment was recorded during rabi season, when the experiment was conducted. The cost of botanicals used was obtained from nearby market. The total cost of plant protection consisted of cost of treatment, sprayer, rent and labour charges for the spray. There are two sprays throughout the research period and the overall plant

Comment [KFA 11-2111]: Give a citation of the recommended package otherwise give details

Comment [KFA 11-2112]: Look for the standard way of writing this

Comment [KFA 11-2113]: Give the average temperatures of the rabi season, as is was this time that the research was done

Comment [KFA 11-2114]: How was the seedlings raised

Comment [KFA 11-2115]: Is there improper irrigation

Comment [KFA 11-2116]: At what stage of the crop were the insecticides first applied

Comment [KFA 11-2117]: Not clear

Comment [KFA 11-2118]: ???

Comment [KFA 11-2119]: Be specific. State the type of transformation

Comment [KFA 11-2120]: Last picking of what

Comment [KFA 11-2121]: ????

Comment [KFA 11-2122]: This is already given

protection expenses was calculated. Total income was realized by multiplying the total yield per hectare by the prevailing market price, while the net benefit is obtained by subtracting the total cost of plant protection from the total income. Benefit over the control for each sprayed treatment was obtained by subtracting the income of the control treatment from that of each sprayed treatment.

Comment [KFA 11-2123]: Better use equations

## RESULTS AND DISCUSSION

Among insect pests infesting brinjal, shoot and fruit borer [*Leucinodes orbonalis*] is a major pest that damages tendershoots and fruits. Its larvae bore inside brinjal fruit and making fruit unfit for human consumption. *L. orbonalis* is the most destructive pest causing economic damage to the crop at all the growth stages. The deformity in shape and remain stunted in growth freshly hatched larvae bore into tender shoots and tunnel downwards. As a result formation of buds, flowers and fruits, the larvae bore inside and feed on inner tissues. However, various new molecules with different mode of action are available in the market which required to be tested for the control of *L. orbonalis* in brinjal. Hence, the experiment was planned to evaluate the efficacy of various synthetic insecticides against *L. orbonalis* in brinjal during rabi season.

Comment [KFA 11-2124]: revise

Comment [KFA 11-2125]: this should be part of the introduction. Besides white use the term new molecules instead of new insecticides with novel mode of action

Effect of different insecticides on the incidence of *Leucinodes orbonalis* showed that all the treatments were significantly superior in reducing the infestation of shoot and fruit borer resulting in increasing the yield, significantly as compared to control. The first spray was given applied after 30 days of transplanting. The percent infestation of shoot and fruit borer on brinjal after first spray (Shoot) revealed that all the chemical treatments were significantly superior over control. Among all the treatments lowest per cent shoot, infestation was recorded in Chlorantraniliprole 18% EC (10.85%), followed by Spinosad 45% SC (11.95%), Flubendamide 20 WG (13.76%), Indoxacarb 14.5% SC (14.05%), Emamectin benzoate 5 SG (15.59%) and Lambda cyhalothrin 4.9% CS (17.09%). The treatment Neem oil 0.15% EC (19.97%) was least effective among all the treatments and maximum shoot damage was recorded in control plot (24.49%). (Table 1)

Comment [KFA 11-2126]: move to materials and methods

The second spray was applied after 15 days of first spray and data on percent fruit damage was recorded. The pooled data for second spray shows minimum percent fruit damage in Chlorantraniliprole 18% EC (7.29%), Spinosad 45% SC (9.12%), Flubendamide 20 WG (9.79%), Indoxacarb 14.5% SC (10.47%), Emamectin benzoate 5 SG (13.46%), and Lambda cyhalothrin 4.9% CS (13.46%), The treatment Neem oil 0.15% EC (17.43%) was least effective

among all the treatments. The highest fruit damage was recorded in Control plot (24.94%). (Table 1)

All of the pesticides were discovered to be extremely effective and greatly *suppressed L. orbonalis as compared to the* control. The lowest documented level of brinjal shoot and fruit borer infection was found in Chlorantraniliprole 18% EC (222.50q/ha) **Tripura, A., et al (2017)**. Spinosad 45% SC (198.33 q/ha) this finding were associated with **Samota and Jat (2014)**. Flubendamide 20 WG (179.16 q/ha) and Indoxacarb 14.5% SC (163.33 q/ha). These findings were supported by **Mahata, S., et al. (2014)**, Emamectim benzoate 5 SG (143.33 q/ha) **Dey, S. (2019)**, and Lambda cyhalothrin 4.9% CS (136.66 q/ha) **Sen, et al. (2017)**. The treatment Neem oil 0.15% EC (127.50 q/ha), validated these finding **Sharma, J. H., et al. (2017)** and the control (90q/ha). (Table 1)

Formatted: Font: Italic

Comment [KFA 11-2127]: re

Comment [KFA 11-2128]: rewrite this

When the cost-benefit ratio was calculated, an intriguing result was obtained; the best and most cost-effective treatment, Chlorantraniliprole 18% EC (1:10.06), among those investigated, is identical to those found by **Pawar et al., (2019)**. The results of Spinosad 45% SC (1:8.51) and Flubendamide 20 WG (1:8.44) are comparable to those of **Jat, H.K., et al. (2020)** and **Shanmugam and Indhumathi (2015)** respectively. Emamectim benzoate 5 SG (1:7.12) and Indoxacarb 14.5% SC (1:8.04) are similar to findings from **Biswas, M., et al., (2020)**. lambda cyhalothrin 4.9% CS (1:6.85), Neem oil 0.15% EC (1:6.36), treatments that are similar to those found by **Sen, et al. (2017)**, **Sanjana and Tayde (2017)**, and control (1:4.63). (Table 1)

**Tableno.1Comparativeefficacyofselectedinsecticideswithneemoilagainstfruitandshootborer [*Leucinodesorbonalis*] on brinjal**


S.No.	Treatments	% Infestation of brinjal shoot and fruit borer						Yield (q/ha)	C:Bratio	
		First spray			Second spray					
		1DBS	3DAS	7DAS	14DAS	3DAS	7DAS			14DAS
T <sub>1</sub>	Chlorantraniliprole@18.5%EC	20.81	11.65 <sup>d</sup>	07.33 <sup>e</sup>	13.58 <sup>ab</sup>	07.33 <sup>e</sup>	05.18 <sup>e</sup>	09.38 <sup>B</sup>	222.50	1:10.06
T <sub>2</sub>	Spinosad@45% SC	19.77	11.92 <sup>d</sup>	09.38 <sup>de</sup>	14.56 <sup>d</sup>	08.69 <sup>de</sup>	08.30 <sup>de</sup>	10.39 <sup>e</sup>	198.33	1:8.51
T <sub>3</sub>	Flubendamide@20WG	19.76	13.97 <sup>cd</sup>	11.83 <sup>cd</sup>	15.48 <sup>d</sup>	10.04 <sup>de</sup>	07.08 <sup>d</sup>	12.27 <sup>e</sup>	179.16	1:8.44
T <sub>4</sub>	Indoxacarb@14.5 %SC	18.51	14.30 <sup>cd</sup>	12.39 <sup>c</sup>	15.48 <sup>cd</sup>	10.37 <sup>d</sup>	09.28 <sup>cd</sup>	11.76 <sup>de</sup>	163.33	1:8.04
T <sub>5</sub>	Emamectin benzoate @ 5SG	21.83	14.84 <sup>cd</sup>	12.50 <sup>c</sup>	19.43 <sup>cd</sup>	13.17 <sup>c</sup>	12.10 <sup>bc</sup>	15.13 <sup>de</sup>	143.33	1:7.12
T <sub>6</sub>	Lambdacyhalothrin @4.9 %CS	20.24	17.58 <sup>bc</sup>	13.49 <sup>c</sup>	20.22 <sup>bc</sup>	13.49 <sup>c</sup>	12.34 <sup>b</sup>	14.57 <sup>c</sup>	136.66	1:6.85
T <sub>7</sub>	Neemoil0.15%EC	23.08	20.23 <sup>ab</sup>	17.76 <sup>b</sup>	21.93 <sup>b</sup>	18.10 <sup>b</sup>	14.50 <sup>b</sup>	19.70 <sup>cd</sup>	127.50	1:6.36
T <sub>8</sub>	Control	20.43	22.97 <sup>a</sup>	24.56 <sup>a</sup>	25.95 <sup>a</sup>	23.82 <sup>a</sup>	24.77 <sup>a</sup>	26.24 <sup>a</sup>	90.00	1:4.63
	F-test	NS	S	S	S	S	S	S		

Comment [KFA 11-2129]: was the comparison of the other insecticides against neem oil? If so make is clear in you objective

Comment [KFA 11-2130]: give full meanings of all abbreviations as foot note under the table

S.Ed( $\pm$ )	01.84	02.28	01.29	02.15	01.29	01.31	01.55	
C.D.(P = 0.5)	-	04.89	02.78	04.61	02.78	02.83	03.33	

UNDER PEER REVIEW



From the present study, the results showed that chlorantraniliprole 18.5% EC is most effective treatment against brinjal fruit and shoot borer producing maximum yield and recorded high cost benefit ratio compared to other treatments. While Spinosad 45% SC, Flubendamide 20 WG, Indoxacarb 14.5% SC, Emamectin benzoate 5 SG has shown average results Lambda cyhalothrin 4.9% SC, Neem oil 0.15% EC, has least effectiveness and in botanicals Neem oil found to be least effective in managing *Leucinodes orbonalis*. Botanicals are the part of integrated pest management in order to avoid indiscriminate use of pesticides causing pollution in the environment and not much harmful to beneficial insects.

Comment [KFA 11-2131]: is this your conclusion? If so where is your heading

## References

- Anwar, S., Mari, J. M., Khanzada, M. A and Ullah, F. (2015). Efficacy of insecticides against infestation of brinjal fruit borer, *Leucinodes orbonalis* Guenee (Pyralidae: Lepidoptera) under field conditions. *Journal of Entomology and Zoology Studies*, 3(3), 292-295.
- Biswas, M., Mishra, I., and Mishra, B. K. (2020). Efficacy of newer molecules of insecticides against brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee (Lepidoptera: Crambidae) and fruit yield, *Journal of Entomology and Zoology Studies*, 8(2): 1562-1566.
- Dey, S. (2019). Evaluation of some novel insecticides on brinjal fruit and shoot borer (BFSB) in West Bengal, India. *The Pharma Innovation Journal*, 8 (12): 337-341.
- Jat, H.K., Shrivastava, V.K. and Dubey, R. (2020) Management of the brinjal shoot and fruit borer *Leucinodes orbonalis* (Guen.) through newer insecticides in brinjal. *Journal of Entomology and Zoology Studies*, 8(5): 1259-1261.
- Kaur, J., Kang, B. K. and Singh, B. (2014) Base line data for insecticide resistance monitoring in brinjal shoot and fruit borer, *Leucinodes orbonalis*, Guenee. *International Quarterly Journal of Life Sciences*, 9(4): 1395-1398.
- Kolhe, P. S., Kumar, A., and Tayde, A. R. (2017). Field Efficacy of Certain Chemicals and Neem Products against Shoot and Fruit Borer (*Leucinodes orbonalis* Guenee) on Brinjal (*Solanum melongena* L.) in Trans Yamuna Region of Allahabad. *International Journal of current Microbiology and Applied Sciences*, 6(9), 1320-1327.
- Mahata, S., Das, B. C., Patra, S., Biswas, A. K., Chatterjee, M. L., and Samanta, A. (2014). New Diamide insecticides against fruit and shoot borer (*Leucinodes orbonalis* Guen.) in Brinjal. *Pesticide Research Journal*, 26(2), 197-201.

- Murugesan, N. and Murugesan, T. (2009)** Bioefficacy of Some Plant Products Against Brinjal Fruit Borer, *Leucinodes orbonalis* (Guenee) (Lepidoptera: Pyralidae). *Journal of Biopesticides*, **2**(1):60-63.
- Pawar, S. A., Bhalekar, M. N., and Datkhile R. V. (2019)**. "Evaluation of different insecticide use strategies for the management of shoot and fruit borer in Brinjal." *Journal of Entomology and Zoology Studies*, **7**(6): 1044-1047.
- Sanjana, V.V.S.P. and Tayde, A.R. (2019)**. Field Efficacy of Some Insecticides, Neem oil and Spinosad against Shoot and Fruit Borer, *Leucinodes orbonalis* (Guenee) on brinjal. *Journal of Entomology and Zoology Studies*, **7**(5): 563-566.
- Samota, R. G. and Jat, B. L. (2014)**. Bio-efficacy of newer insecticides and bio-pesticides against *Leucinodes orbonalis* on brinjal. *Annals of Agri Bio Research*, **19**(2):285-289.
- Shanmugam, P. S., Indhumathi, K., Vennila, M. A. and Tamilselvan, N. (2015)**. Evaluation of bio-intensive pest management modules against brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee (Lepidoptera: Crambidae) under precision farming system. *Pest Management in Horticultural Ecosystems*, **21**(2):154-158.
- Sharma, J. H. and Tayde, A. R. (2017)**. Evaluation of bio-rational pesticides, against brinjal fruit and shoot borer, *Leucinodes orbonalis* Guen. On brinjal at Allahabad Agroclimatic region. *International Journal of Current Microbiology and Applied Sciences*, **6**(6), 2049-2054.
- Singh, J.P., Singh, R. and Singh, S. (2018)**. Efficacy of Newer Insecticides and Biopesticides Against Shoot and Fruit Borer, *Leucinodes orbonalis* (Guenee) brinjal (*Solanum melongena* Linnaeus). *Journal of Pharmacognosy and Phytochemistry*, **6**(4):339-347.
- Sen, K., Samanta, A., Alam, S.K.F., Dhar, P.P. and Samanta, A. (2017)**. Field Evaluation of a New Ready Mix Formulation Ampligo 150ZC (Chlorantraniliprole 9.3% + Lambda Cyhalothrin 4.6% ZC) against Shoot and Fruit Borer (*Leucinodes orbonalis* Guen.) infestation in Brinjal. *Journal of Pharmacognosy and Phytochemistry*, **6**(5):1674-1678.
- Tripura, A., Chatterjee, M. L., Pande, R., and Patra, S. (2017)**. Biorational management of brinjal shoot and fruit borer (*Leucinodes orbonalis* guenee) in mid hills of meghalaya. *Journal of Entomology and Zoology Studies*, **5**(4), 41-45.

UNDER PEER REVIEW

