

## Original Research Article

### **Comparative efficacy and economics of chemicals insecticides and bioagents against diamondback moth (*Plutella xylostella*) on cabbage(*Brassica oleracea* var. *capitata* L.)**

#### **Abstract**

Field trial was conducted during *rabi* season 2021-2022 at Central Research Farm (CRF), SHUATS. The experiment was laid out in Randomised Block Design with eight treatments each replicated thrice using a variety Green Soccer (546). The treatments were Chlorantraniliprole 18.5 %SC -T<sub>1</sub>, Indoxacarb 14.5%SC-T<sub>2</sub>, Emamectin benzoate 5%SG-T<sub>3</sub>, Spinosad 45% SC-T<sub>4</sub>, *Beauveria bassiana* (1x10<sup>8</sup>CFU/ml)-T<sub>5</sub>, *Metarhizium anisopilae* (1x10<sup>8</sup>CFU/ml)-T<sub>6</sub>, *Bacillus thuringiensis* (1x10<sup>8</sup>CFU/ml)-T<sub>7</sub> and untreated control -T<sub>8</sub>. Mean reduction in the larval population per plant revealed that all the treatments were significantly superior over the control (6.51). Chlorantraniliprole 18.5% SC recorded lowest mean larval population of *P. xylostella* (1.52), followed by Spinosad 45% SC (1.75), Indoxacarb 14.5% SC (2.00), Emamectin Benzoate 5%SG (2.28), *B. thuringiensis* (2.68), *B. bassiana* (2.87), *M. anisopilae* (3.02). Highest yield (280 q/ha) as well as B:C ratio (1:6.37) was obtained from the treatment Chlorantraniliprole 18.5% SC followed by Indoxacarb 14.5% SC (1:6.33), Emamectin Benzoate 5%SG (1:6.25), *B. thuringiensis* (1:5.97), Spinosad 45% SC (1:5.75), *B. bassiana* (1:5.72), *M. anisopilae* (1:5.50) as compared to control (1:3.34).

**Keywords:** Chlorantraniliprole, *Bacillus thuringiensis*, *Beauveria bassiana*, Cost benefit ratio, *Plutella xylostella*.

#### **Introduction**

Cabbage (*Brassica oleracea* var. *capitata*) is one of the most popular Cole vegetables grown in India. It is originated in Europe and in the Mediterranean region after cauliflower.

Cabbage is also used in herbal medicine. Cabbage juice can reduce constipation and has also been used as a laxative, as an antidote to mushroom poisoning, or a treatment for hangovers and headaches. In fact, cabbage has historically been used to stop sunstroke, or to relieve fevers. The leaves were also used to soothe swollen feet and to treat childhood croup. Brassica vegetables have also anti-inflammatory activity and have been used to different irritations of the human body. (Alexandra *et al.*,2020).

Regular consumption of dark green leafy vegetables is highly recommended because of their potential in reducing chronic diseases. (Miller-Cebert *et al.*,2009) and glucosinolates in cabbage reduced risk of cancer induction and development. (Kang *et al.*,2006).It is known to possess medicinal properties and its enlarged terminal buds is a rich source of Ca, P, Na, K, S, Vitamin A, Vitamin C and dietary fibre. It is said to be good for person suffering from diabetes. It may be used to prepare soup, stew, as stuffing for cake (Norton *et al.*,1997).

In India, West Bengal accounts highest production of cabbage in the world which is 2288.50 tonnes, which has the share of 25.32 percent followed by Orissa 1058.78, tonnes, Madhya Pradesh 686.91 tonnes, Bihar 673.44 tonnes, and Uttar Pradesh 302.97. (NHB,2017-2018).

The brassica crop has a multiple insect pest complex. A total of 37 insect pests have been reported to feed on cabbage in India (Lal *et al.*,1975). The important insect pest species are Diamondback moth (*Plutella xylostella* L), Cabbage caterpillar (*Pieris brassicae* Linnaeus), Cabbage semi-looper (*Thysanoplusia orichalcea* Fabricius) and (*Autographa nigrisigna* Walker), Tobacco caterpillar (*Spodopteralitura* Fabricius), Cabbage leaf Webber (*Crocodolomia binotalis* Zeller), Cabbage borer (*Hellula undalis* Fabricius) and Cabbage aphid (*Brevicoryne brassicae* W).Of these Diamondback moth, *Plutella xylostella* (L.) is the most destructive pest (Kumar *et al.*,2007) and is the limiting factor for the successful cultivation of cruciferous crops. *Plutella xylostella* was first recorded in 1746 and probably from European origin. About 128 countries or regions reported infestation by this insect pest in 1972.The level of infestation varies from place to place for example the infestation is serious in south and southeast Asian countries and moderate in other Asian countries than the

Mediterranean region. *Plutella xylostella*. (L.) is a common pest (**Harcourt, 1963**).

UNDER PEER REVIEW

## Materials and methods

The experiment was conducted at Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Prayagraj Uttar Pradesh (U.P) during the *rabi* season of 2021-2022 with a recommended package of practices excluding plant protection. Cabbage seedlings (var 'Green Soccer-546') transplanted after 40 days at 60 cm x 45 cm spacing. The experiment was laid down in Randomized Block Design (RBD) with eight treatments replicated thrice with each plot size of 2m X 2m and proper irrigation was provided. The treatments comprising of Chlorantraniliprole 18.5%SC, Spinosad 45% SC, Indoxacarb 14.5% SC, Emamectin benzoate 5% SG, *Bacillus thuringiensis* ( $1 \times 10^8$ CFU/ml), *Beauveria bassiana* ( $1 \times 10^8$ CFU/ml), *Metarhizium anisopliae* ( $1 \times 10^8$ CFU/ml), and were applied in two sprayings at 15 days interval with recommended doses when larval population reaches its ETL level.

Observations on total number of larvae on cabbage of five observational plants from each treatment replication wise were recorded at 1 Day before spraying, 3<sup>rd</sup>, 7<sup>th</sup> and 14<sup>th</sup> days after imposing treatments. The data recorded in the different treatments were subjected to statistical analysis after suitable transformation by following standard procedures of RBD experiment. After harvesting of cabbage from each individual plots produce were calculated to work out the yield of the treatments. Yield of healthy heads was converted into quintal per hectare.

The cost of Insecticides and biopesticides used in the experiment was obtained from the local market. The total cost of plant protection consisted of cost of treatment, sprayer, rent and labour charges for the spray. There are two sprays throughout the research period and the overall plant protection expenses was calculated. Total income was realized by multiplying the total yield per hectare by the prevailing market price, while the net benefit is obtained by subtracting the total cost of plant protection from the total income. Benefit over the control for each sprayed treatment was obtained by subtracting the income of the control treatment from that of each sprayed treatment.

## Results and discussion

Effect of different insecticides and biopesticides on the incidence of *Plutella xylostella* revealed that all the treatments were significantly superior in reducing the infestation of Diamondback moth resulting in increasing the yield, significantly as compared to control. The first spray was given after 30 days of transplanting. The larval population of Diamondback moth on cabbage after first spray revealed that all the chemical treatments were significantly superior over control. Among all the treatments lowest larval population, was recorded in Chlorantraniliprole 18.5% SC (2.17) followed by Spinosad 45% SC (2.37), Indoxacarb 14.5% SC (2.65), Emamectin benzoate 5SG (2.95), *Bacillus thuringiensis* ( $1 \times 10^8$ CFU/ml) (3.44) and *Beauveria bassiana* ( $1 \times 10^8$ CFU/ml) (3.55). The treatment *Metarhizium anisopliae* ( $1 \times 10^8$ CFU/ml) (3.66) was least effective among all the treatments but maximum damage was recorded in control plot (5.77) (Table 1).

The second spray was after 15 days of first spray. The data for second spray shows minimum larval population in Chlorantraniliprole 18.5% SC (0.86) followed by Spinosad 45%SC (1.13), Indoxacarb 14.5% SC (1.35), Emamectin benzoate 5% SG (1.62), *Bacillus thuringiensis* ( $1 \times 10^8$ CFU/ml) (1.93) and *Beauveria bassiana* ( $1 \times 10^8$ CFU/ml) (2.2) The treatment *Metarhizium anisopliae* ( $1 \times 10^8$ CFU/ml) (2.37) was least effective among all the treatments. The highest mean larval population was recorded in Control plot (7.24) (Table 1).

All the insecticides were found very effective and significantly over control. The data for overall mean larval population was recorded of which least larval population was recorded in Chlorantraniliprole 18.5% SC (1.52), Spinosad 45%SC (1.75), Indoxacarb 14.5% SC (2.00), Emamectin benzoate 5%SG (2.28), *Bacillus thuringiensis* ( $1 \times 10^8$ CFU/ml) (2.68) and *Beauveria bassiana* ( $1 \times 10^8$ CFU/ml) (2.87) The treatment *Metarhizium anisopliae* ( $1 \times 10^8$ CFU/ml) (3.02) was least effective among all the treatments but control treatment had higher mean larval population of 6.51 (Table 1).

Highest yield and benefit cost ratio was recorded in Chlorantraniliprole 18.5% SC (325 q/ha) (1:6.37) is similar to the findings of **Sharma et al.,(2016)**, Spinosad 45% SC (310 q/ha) (1:5.75) is similar to the findings of **Gill et al.,(2008) and Gaddam et al.,(2021)**, Indoxacarb 14.5% SC (300 q/ha) (1:6.33) is similar to the findings of **Harika et al.,(2019)**

and Gaddam *et al.*,(2021), Enamectin benzoate 5% SG (295 q/ha) (1:6.25) is similar to the findings of Akbar *et al.*, (2014) , *Bacillus thuringiensis* ( $1 \times 10^8$ CFU/ml) (280 q/ha ) (1:5.97) is similar to the findings of Choyon *et al.*,(2022) and *Beauvaria bassiana* ( $1 \times 10^8$ CFU/ml) (270q/ha) (1:5.72) is similar to the findings of Debbarma *et al.*,(2017) The treatments *Metarhizium anisopliae* ( $1 \times 10^8$ CFU/ml) (260 q/ha) (1:5.50) is similar to the findings of Singh *et al.*, (2015), and the lowest yield was recorded in control (150 q/ha) (1:3.34) (Table 1).

UNDER PEER REVIEW

**Table 1: Effect of certain insecticides and bioagents on larval population of diamondback moth (*Plutella xylostella*).**

S.No.	Treatments	Larval population of diamondback moth								Overall mean population	Yield (q/ha)	C:B ratio	
		1 <sup>st</sup> spray				2 <sup>nd</sup> spray							
		1DBS	3DAS	7DAS	14DAS	MEAN	3DAS	7DAS	14DAS				MEAN
T1	Chlorantraniliprole 18.5 % SC (0.3ml/l)	04.06 (11.60)*	02.6 <sup>d</sup> (9.27)*	01.8 <sup>e</sup> (7.70)*	02.13 <sup>e</sup> (8.39)*	2.17 (8.46)*	01.13 <sup>e</sup> (6.10)*	00.93 <sup>f</sup> (5.53)*	00.53 <sup>f</sup> (4.17)*	0.86 (5.27)*	01.52	325	1:6.37
T2	Indoxacarb 14.5% SC (1ml/l)	04.26 (11.91)*	03.13 <sup>c</sup> (10.18)*	02.33 <sup>de</sup> (8.76)*	02.5 <sup>cd</sup> (9.09)*	2.65 (9.35)*	01.6 <sup>cd</sup> (7.25)*	01.26 <sup>ef</sup> (6.45)*	01.2 <sup>de</sup> (6.27)*	1.35 (6.66)*	02.00	300	1:6.33
T3	Emamectin benzoate 5 %SG(0.6 gm/l)	04.6 (12.32)*	03.26 <sup>c</sup> (10.39)*	02.66 <sup>cd</sup> (9.37)*	02.93 <sup>b</sup> (9.85)*	2.95 (9.82)*	01.73 <sup>c</sup> (7.56)*	01.53 <sup>de</sup> (7.1)*	01.6 <sup>cd</sup> (7.24)*	1.62 (7.31)*	02.28	295	1:6.25
T4	Spinosad 45% SC (2ml/l)	04.4 (12.09)*	02.73 <sup>d</sup> (9.50)*	02.06 <sup>e</sup> (8.23)*	02.33 <sup>de</sup> (8.78)*	2.37 (8.84)*	01.26 <sup>de</sup> (6.44)*	01.2 <sup>ef</sup> (6.27)*	00.93 <sup>ef</sup> (5.51)*	1.13 (6.08)*	01.75	310	1:5.75
T5	<i>Beauveria bassiana</i> (1x10 <sup>8</sup> CFU/ml) (2ml/l)	04.46 (12.19)*	04.73 <sup>b</sup> (12.58)*	03.2 <sup>bc</sup> (10.30)*	02.73 <sup>bc</sup> (9.51)*	3.55 (10.79)*	02.46 <sup>b</sup> (9.02)*	02.2 <sup>bc</sup> (8.51)*	01.93 <sup>bc</sup> (7.98)*	2.2 (8.51)*	02.87	270	1:5.72
T6	<i>Metarhizium anisopliae</i> (1x10 <sup>8</sup> CFU/ml) (2ml/l)	04.8 (12.63)*	04.86 <sup>ab</sup> (12.72)*	03.26 <sup>b</sup> (10.39)*	02.86 <sup>b</sup> (9.47)*	3.66 (10.95)*	02.66 <sup>b</sup> (9.39)*	02.4 <sup>b</sup> (8.90)*	02.06 <sup>b</sup> (8.26)*	2.37 (8.85)*	03.02	260	1:5.50
T7	<i>Bacillus thuringiensis</i> (1x10 <sup>8</sup> CFU/ml) (2ml/l)	04.46 (12.19)*	04.6 <sup>b</sup> (12.37)*	03.06 <sup>bc</sup> (10.08)*	02.66 <sup>bc</sup> (9.38)*	3.44 (10.61)*	02.33 <sup>b</sup> (8.77)*	01.86 <sup>cd</sup> (7.85)*	01.6 <sup>cd</sup> (7.26)*	1.93 (7.92)*	02.68	280	1:5.97
T0	Control	04.33 (11.97)*	05.2 <sup>a</sup> (13.16)*	05.73 <sup>a</sup> (13.84)*	06.4 <sup>a</sup> (14.65)*	5.77 (13.89)*	06.73 <sup>a</sup> (15.03)*	07.2 <sup>a</sup> (15.56)*	07.8 <sup>a</sup> (16.21)*	7.24 (15.60)*	06.51	150	1:3.34
	F-test	NS	S	S	S	S	S	S	S	S	-	-	-
	C.D. at 0.5%	---	00.35	00.55	00.30	0.99	00.41	00.41	00.40	0.51	-	-	-
	S.EdA (±)	00.29	00.16	00.82	00.14	0.46	00.18	00.19	00.18	0.24	-	-	-

DBS\*= Days before spraying, \*Figures in parenthesis are Arc sin transformed values.

## CONCLUSION

From the present study, the results it showed that T<sub>1</sub> Chlorantraniliprole 18.5 % SC most effective treatment against diamondback moth of Mean larval population and producing maximum yield and recorded highest Cost-Benefit ratio compared to other treatments. While T<sub>2</sub> Indoxacarb 14.5% SC, T<sub>3</sub> Emamectin benzoate 5 %SG, T<sub>4</sub> Spinosad 45% SC , has shown average results has proved to be least effective chemicals. *Beauveria bassiana*, *Metarhizium annisopliae* and *Bacillus thuringiensis*, found to be least effective in managing *Plutella xylostella*. Botanicals are the part of integrated pest management in order to avoid indiscriminate use of pesticides causing pollution in the environment and not much harmful to beneficial insects.

**Acknowledgements:** The authors are grateful to Prof. (Dr.) Rajendra B. Lal Hon'ble Vice Chancellor SHUATS, Prof. Dr. Shailesh Marker, Director of research, Dr. Deepak Lal, Dean of PG studies, Dr. Gautam Gosh, Dean, Naini Agricultural Institute, Sam Higginbottom University of Agriculture Technology and Sciences, for taking their keen interest and encouragement to carry out this research work.

## References

1. **Akbar, Ilyas., Sattar, Shahid., Khurshid, Isma. and Akbar, Aneela. (2014).** Comparative efficacy of synthetic insecticides and botanical extracts against diamondback moth (*Plutella xylostella*)(Lepidoptera: Plutellidae) in cauliflower. *European. Academic. Research*, 2: 6051-6066.
2. **Alexandra, S. I. M. and Andrea Daniela, O. N. A. (2020)** Cabbage (*Brassica oleracea* L.). Overview of the health benefits and therapeutical uses. *Hop and Medicinal Plants*, 1(2):150-169 .
3. **Choyon, M. M., Akhter, N., Rahman, M. M. and Hossain, M. E. (2022).** Effectiveness of some bio-pesticides in managing major Lepidopteran insect Pests of Cabbage (*Brassica oleracea* var. capitata L.). *Asian Research Journal of Agriculture*, 15(3): 44-51.
4. **Debbarma, A., Singh, K. I., Gupta, M. K. and Sobitadevi, P. (2017).** Bio-rational management of major lepidopterous pests and their influence on yield of cabbage crop under Manipur valley. *Journal of Entomology. and Zoology. Studies*, 5(5): 1546-1551.
5. **Gaddam, N. R., Srivastava, V. K., Tayde, A. R. and Tripathi, A. (2021).** Comparative efficacy of microbials and botanicals against diamondback moth (*Plutella xylostella* Linn) on cabbage. *Journal of Entomology and Zoology Studies*, 9(1): 497-499.
6. **Gill, C. K., Kaur, S. and Joia, B. S. (2008).** Efficacy of new insecticides for the management of diamondback moth, *Plutella xylostella* (Linnaeus) on cauliflower and cabbage. *Journal of Insect Science*, 21(2):171-177.
7. **Harcourt, D.G. (1963)** Major mortality factors in the population dynamics of the Diamondback moth, *Plutella maculipennis* (Curt.) (Lepidoptera: Plutellidae) *Memoirs of the Entomological Society of Canada*, 95(32): 55-66.
8. **Harika, G., Dhurua, S., Suresh, M. and Sreesandhya, N. (2019).** Evaluation of Certain Insecticides against Diamondback moth (DBM) *Plutella xylostella* on Cauliflower. *International Journal of Bio-resource and stress Management*, 10(1): 070-076.

9. **Kang, J. Y., Ibrahim, K. E., Juvik, J. A., Kim, D. H. and Kang, W. J. (2006).** Genetic and environmental variation of glucosinolate content in Chinese cabbage. *Horticulture Science*, 41(6): 1382-1385.
10. **Kumar, P., Prasad, C. S. and Tiwari, G. N. (2007).** Population intensity of insect pests of cabbage in relation to weather parameters. *Annals of Plant Protection Sciences*, 15(1): 245-246.
11. **Lal, O.P. (1975).** A Compendium of insect pest of vegetables in India. *Bulletin of Entomological Research*, 16: 31-56.
12. **Miller-Cebert, R. L., Sistani, N. A. and Cebert, E. (2009).** Comparative protein and foliate content among canola cultivars and other cruciferous leafy vegetables. *Journal of Food, Agriculture And Environment*, 7(2):42-49.
13. **Norton, G. A. (1997)** The economic and social context of pest, diseases and weed problems. In origin of pest, parasites, disease and weed problems. Ed. By J. M. Cherret and G. R. Sagar. *Symposium of the british Ecological Society*, 205-226.
14. **Sharma, S., Singh, R. and Gill, C. K. (2016).** Efficacy of anthranilic insecticide E2Y45 20 SC (Chlorantraniliprole) against *Plutella xylostella* L. in cabbage, *Brassica oleracea* var. capitata. *Journal of Applied and Natural Science*, 8(3): 1584-1588.
15. **Singh, K. I., Debbarma, A. and Singh, H. R. (2015).** Field efficacy of certain microbial insecticides against *Plutella xylostella* Linnaeus and *Pieris brassicae* Linnaeus under cabbage-crop-ecosystem of Manipur. *Journal of Biological Control*, 29(4): 194-202.