

Original Research Article

“Effects of different growing media and biocapsules on seed germination, seedling growth and survival percentage of papaya (*Carica papaya*) cv. Pusa Nanha”

ABSTRACT

The study entitled, **Effects of different growing media and biocapsules on seed germination, seedling growth and survival percentage of papaya (*Carica papaya*) cv. Pusa Nanha** was carried out during March – June 2022 at Department of Horticulture, Naini Agricultural Institute, SHUATS, Prayagraj. The experiment was carried out to see the performance of the most suitable soil media for growth and establishment of Papaya seedlings. There were nine treatments including control viz, M0 Control (Garden soil), M1(Soil + FYM), M2 (Soil+FYM+biocapsules), M3 (Soil+FYM+Neemcake), M4 (Soil + FYM + Neemcake + biocapsules), M5 (Soil +FYM + Vermicompost), M6 (Soil + Sand + Vermicompost + biocapsules), M7 (Soil + FYM + Sand) M8 (Soil+FYM+Sand+Biocapsules) replicated three times with 10 seeds per replication. Growth medium significantly affected the growth, biomass, and root morphological indexes of papaya seedlings. The results reveal that the treatment M6 (Soil+Sand+Vermicompost+biocapsules) was the most suitable treatment over all the other treatments in relation to germination and growth parameters.

Keywords: Papaya, growing media, FYM, Vermicompost, Biocapsule, Neemcake

INTRODUCTION:

Papaya (*Carica papaya L.*) belongs to caricaceae family and its native place is Tropical America. It is an important fruit crop of tropical and sub-tropical countries of the world, covering 32 °N and 32 °S on the Globe mainly grown in America, Brazil, Columbia, Costa Rica, Hawaii, India, Mexico, Nigeria, Phillipines and Bangladesh. Papaya is called papaw or pawpaw, an ideal fruit for growing in kitchen gardens, backyards of homes as well as field, especially near the cities or big towns. It is also grown extensively as a filler plant in orchards. It is a quick growing typically single-stemmed, short lived, large, perennial herb, and starts bearing within 8-10 months of transplanting. It is one the most popular commercial fruit in India.

In India papaya is extensively grown throughout the country viz. Andhra Pradesh, Tamil Nadu, Assam, Bihar, Maharashtra, Gujarat, Haryana, M.P. etc. occupying an area of 132.18 thousand hectares with

annual production of 5381.73 thousand tones. Whereas, it is grown on 0.40 thousand hectares with annual production of 13.2 thousand tones in Rajasthan (**Anonymous,2014**). Among the various varieties of papaya Pusa Dwarf, Taiwan, Pusa Delicious, CO-1, CO-2, CO-3, CO-5, CO-6, Coorg Honey Dew, Pant C-1, Sunrise Solo, Red Lady have attained the commercial status throughout the country.

Papaya is a very wholesome, refreshing and delicious fruit and rich source of vitamins. Immature papaya fruit is a rich source of papain, whereas mature fruit is rich source of carbohydrates (7.2%), minerals, Vitamin A (2020 IU/100g) and Vitamin C (85mg/100g) (**Gupta and Naik.,2007**). It has high medicinal value. Papain prepared from dried latex of its immature fruits is used in meat tenderizing, manufacture of chewing gum, cosmetics, degumming of natural silk and to give shrink resistance to wool. Papaya has an in vitro antiproliferative effect in liver cancer cells, possibly due to lycopene (**Asmah et al., 2002**). Its fruits are consumed fresh as well as processed commercially into jam, jelly, nectar, squash, cheese, puree, etc. papaya is highly remunerative due to high yield potential and good filler fruit crop that provide extra income to the orchardist. By adopting proper management practices and selection of suitable varieties a gross income of Rs 1.90 lac can be obtained within a year of planting from papaya crop. (**Shukla et al.,2008**)

Papaya is a quick growing, continuous fruiting, evergreen plant and requires a good fertile soil or growing media for both growth, development and quality of fruits. The use of suitable growing media or substrates for sowing of seeds directly affects the germination, development and functional rooting system. A good growing medium provides sufficient anchorage or support to the plant, serves as reservoir for nutrients and water, allow oxygen diffusion to the roots and permit gaseous exchange between the root and atmosphere outside the root substratum . The quality of seedlings is very much influenced by growing media under nursery. The quality of seedlings obtained from a nursery influences re-establishment in the field and the eventual productivity of an orchard (**Baiyeri, 2006**).

Since papaya is commonly propagated by seeds, the various factors like type of potting media used, water, oxygen, temperature, light, plant species etc. are the most important for successful production of the crop. (**Hartman et al., 2001**). The germination of papaya seeds was reported slow, erratic and incomplete (**Chako and Singh,1966**).

Germination of papaya seed faces certain problems of seedling mortality due to damping off diseases in nursery stage. Initial mortality and improper germination is one of the major causes of reduced survival per cent of papaya plants. The development of root system is suppressed and plants are more susceptible to soil borne diseases in heavy soils without drainage (**Beattie and White, 1992**). The papaya seeds are endorsed within a gelatinous sarcotesta (aril or outer seed coat which is formed from the outer integument). Growing media are an integral part of most horticultural production systems. Generally, media for fruit crop seedlings are composed of soil and different organic matters. The soil is generally used as a basic medium because it is cheapest and easy to procure supplementing of the soil which is aimed to make media more porous while the organic matter (vermicompost, FYM, Neemcake) are added so as to enrich adequate nutrients for the seedlings. A growing medium is a substance through which roots grow to extract water and nutrients. The growing medium also plays an important role in seed germination not only it does act as a support, but also a source of key nutrients for plant growth. The composition of the medium influences the quality of the seedlings (**Wilson et al.,2001**). The important media used to raise nursery plants along with their characters are Soil, vermicopost, FYM, neemcake, sand, biocapsules are included in the present study. Soil having good texture should be preferred for using a propagation media.

Sand is used as a rooting media for adding the coarse texture needed to induce proper drainage and aeration. Sand is the heaviest of all the rooting media. Therefore it should be a combination with some other organic material.

Farmyard Manure (FYM) is primarily made from cow dung, cow urine, waste grass, and other dairy waste. Properties of FYM are, FYM is nutrient-rich. A small portion of N is available directly to the plants, while a greater portion is made available when FYM decomposes. Farmyard manure has been the main source of organic matter for the supply of essential minerals needed by the plants. Heavy applications of organic matter are required for successful production of fruits and vegetables (Chaudhary, 1996).

Neem Cake is the de-oiled residue that can be used after Neem kernels are crushed for their oil. In this seed kernel are nutrients like NPK (nitrogen-phosphorus-potassium). These nutrients are nematicidal in nature so the seed cake ends up with these properties. Neem Cake is used in agriculture, horticulture, floriculture and the turf industry as an organic fertilizer as well as a natural nematicide. Neem Cake is used as an organic fertilizer because of the various micro and macro nutrients which it is composed of. It will control, at the same time, soil based pathogens as well as nematodes. It will also inhibit nitrification of the soil plus enhance the efficiency of nitrogen providing fertilizer.

Vermicompost refers to a mixture of worm casting, organic material, humus, living earthworms their cocoons and other organisms. Earthworm reduces C:N ratio, increases humic acid content, cation exchange capacity and water soluble carbohydrates (Talashilkar *et al.*, 1999). Vermicompost is a rich source of nutrients with good water holding capacity.

Biocapsules is a revolutionary technology that ensures the successful delivery of biologically competent beneficial microbes. Biocapsule, a biofertiliser technology developed by - Indian Institute of Spices Research, has started gaining acceptance among the farming community. Recently, ICAR (Indian Council of Agricultural Research) scientists have developed the technology to pack bio-fertilizers in tiny capsules. This eliminates the need for farmers to carry the sacks of biofertilizers.

- It is being called the **one-gram capsule, which have many advantages:**
 - One-gram capsules are very efficient as it contains the microbial population equivalent to what is present in a one-kg pack of powder-based biofertilizer or a one-litre bottle.
 - Also, as these microbial strains are retained in the dormant stage, there is no issue of their viable loss in room temperatures as is the case with many liquid-based bio formulations.

Unique Benefits of biocapsule:

- Each capsule contain 10^9 propagule.
- Enhance seed germination & suppress diseases.
- Low cost, ecologically safe micro for all agriculture and horticulture crop.
- Promote crop growth & increase in yield.
- Long lasting viability up to 2 yrs.
- Easy to use & handling (Drench, seed treatment & use in compost)

Materials and methods:

The experiment entitled was carried out at Department Nursery and Horticulture Research Farm , Naini Agriculture Institute, SHUATS, Prayagraj, Uttar Pradesh during the month of March-June 2022. There were nine treatment combination including control viz, M0 Control (Garden soil), M1(Soil + FYM), M2 (Soil + FYM + biocapsules), M3 (Soil + FYM + Neemcake), M4 (Soil + FYM + Neemcake + biocapsules), M5 (Soil + FYM + Vermicompost), M6 (Soil + Sand + Vermicompost + biocapsules), M7 (Soil + FYM + Sand) M8 (Soil + FYM + Sand + Biocapsules) replicated three times with 10 seeds per replication. Seeds were sown in the month of march. The polybags were irrigated after sowing and repeated everyday until emergence and in alternate days after germination.

Experimental design and measured parameters:

The experiment was carried out in a complete random block design with 30 polybags per treatment. All the germination and growth parameters were recorded at the time of germination till transplanting (60 days after sowing).

The days taken for first seed germination were calculated from the date of sowing up to germination of the first seedling. Germination per cent was calculated as the number of seeds sown and the number of seeds germinated, expressed in percentage.

$$\text{Germination percent} = \frac{\text{No of seeds germinated}}{\text{Total no. of seeds sown}} \times 100$$

Height of the seedling was recorded by measuring the height of five random seedlings by metric scale from the base to the tip of the shoot of the seedling. The numbers of leaves of five seedlings were counted and average value was calculated. The stem girth of seedlings were measured with the help of vernier caliper at height 1cm above ground level. The fresh weight of seedlings was weighed on digital weighing balance expressed in gram. The primary root of seedlings were measured by metric scale from the base of the shoot to tip of root at the time of transplanting. The Root: Shoot ratio was calculated by dividing dry weight of roots by dry weight of shoots of seedlings at the time of transplanting. The chlorophyll content of the leave of the plant was measured using a SPAD (Soil Plant Analysis Development) meter. Survival percentage of papaya seedlings was observed by transplanting them in the field and was calculated using the formula:

$$\text{Survival percentage} = \frac{\text{Total number of surviving seedlings}}{\text{Total number of seedlings transplanted}} \times 100$$

Results and discussion:

The results shows that growing media and biocapsule had beneficial effects on germination, seedling growth and survival percentage of papaya.

Seed germination parameters:

Effects of growing media and biocapsules on seed germination parameters are given in table 1. The treatment T6 was found best followed by treatment T2 regarding germination parameters as it had the best water holding capacity, porosity and nutrients which would have enhanced the availability of water, aeration and nutrition during germination of seed. Germination started at 11.3 days after sowing on Soil +FYM + Vermicompost (1:1:1) + Biocapsule (T6). Similar findings were found in **Venkatesh et al., (2009)** in Pitsal (*Pterocarpus marsipium*, **Abirami (2010)** in nutmeg, **Bhardwaj (2013)** in papaya, **Bhagat et al. (2013)** in rough lemon, **Kumar et al., (2015)** in papaya, **Kaur (2017)** in mango stones and **Gawankar et al., (2021)** in jackfruit. Maximum germination percent 96.67 was also found in T6. This might be because of the media containing organic manures possess organic acid within them. Therefore, more available moisture and some acids may have helped in better germination percentage. Biocapsules also provide microbes that are beneficial to plants. Among the microbes, rhizobacteria, which functions as plant growth promoting rhizobacteria (PGPR) such as *Pseudomonas* spp. and *Bacillus* sp., can serve as fertilizer. These organisms have proven to accelerate germination and improve overall growth of the plant. Similar findings were found in **Abirami (2010)** in nutmeg, **Dayeswari et al., (2017)** in papaya, **Chiranjeevi et al., (2018)** in aonla, **Kumararvind et al., (2020)** in papaya, **Irsa et al., (2021)** in marigold.

Table 1: Effects of different growing media and biocapsule on germination parameters of papaya plant.

Treatments	Initiation period	Germination percent %
T0	18.8	56.67
T1	18.4	63.33
T2	11.6	93.33
T3	15.6	73.33
T4	13.8	83.33
T5	12.7	86.67
T6	11.3	96.67
T7	16.23	66.67
T8	15.4	80.00
C.D.(P=0.05)	0.68	17.90
C.V	2.64	13.30

Seedling growth parameters:

Growth increase in papaya seedling significantly affected by growing media and biocapsule is presented in the data in table 2 and 3. Highest seedling height (23.41 cm) was recorded in T6. Maximum number of leaves (10.8) was recorded in T6 which was at par with T2 (10.2). Soil, vermicompost, FYM provides adequate nutrients and enhances both the physical and biological properties and the water holding capacity of soil while iocapsule provide PGPR (plant growth-promoting rhizobacteria) which results in acceleration of plant growth and protection against plant pathogens. Similar findings

were found in **R.M. Atiyeh et al.,(2004)** in tomato, **Annapurna et al., (2005)** in *Santalum album* L, **Asciutto et al., (2006)** in *Impatiens wallerana*, **Samir et al., (2016)** in Khirni, **Dayeswari et al., (2017)** in papaya, **Kumar et al. , (2018)** in *Albizia lebbeck*

Maximum stem girth (7.09mm), longest root length (12.83cm), highest seedling vigour (3503.33), highest chlorophyll content (38.37 $\mu\text{mol per m}^2$) and highest fresh weight of seedling (14.74 g) were recorded in treatment T6. This might be due to vermicopost which contains plant growth regulations like auxis, gibberllins and cytokinins which are responsible for increase in plant growth and biocapsule which provide PGPR (plant growth-promoting rhizobacteria) which results in acceleration of plant growth. Similar findings were found in **Prabha et al., (2007)** in banana, **Bachman and Metzger (2008)** in tomato, **Parasana et al., (2012)** in mango, **Chiranjeevi et al., (2018)** in aonla.

Similarly highest fresh weight of shoot (10.56 g) and root (4.29 g), highest dry weight of shoot (4.02) and root (0.73), highest total biomass (4.73 g) was recorded in treatment T6. The highest root:shoot ratio (0.18) was recorded in treatment T6 and T2. This maybe attributed to general improvement in the physical and chemical properties of the rooting media which improved the fresh and dry weight of the seedling. Similar findings were found in **Annapurna et al., (2005)** in *Santalum album*, **Asciutto et al., (2006)** in *Impatiens wallerana*.

The maximum survival percentage (82.67) was recorded with treatment M6, followed by M2 (81.33) while minimum was observed under M0 (control) (58.67). Better survival percentage might be due to the reason that the media combination was helpful in reducing damping off diseases in seedling due to preparation in root zone. Good physical and biological conditions in FYM and vermicopost along with biocapsules had positive effect on root and shoot growth which helps in better survival. Similar findings were found in **Bhardwaj (2013)** in papaya, **Ritesh et al., (2016)** in papaya, **Chiranjeevi et al., (2018)** in aonla, **Irsa et al., (2021)** in marigold.

Table 2: Effects of different growing media and biocapsule on growth of papaya seedlings.

Treatments	Seedling height (cm)	No. of leaves	Stem girth (cm)	Root length(cm)	Seedling vigour index	Chlorophyll content ($\mu\text{mol per m}^2$)	Fresh weight of seedling (g)
T0	11.71	5.7	5.21	6.35	1023.8	29.24	4.24

T1	11.93	6.2	5.40	7.88	1255.33	31.41	4.71
T2	22.05	10.2	6.15	12.73	3338.6	37.88	14.18
T3	19.23	8.8	5.72	10.06	2149.57	30.34	9.72
T4	21.09	9.4	6.21	12.38	2789.67	33.05	13.58
T5	22.19	10.03	6.44	12.52	3007.60	34.16	13.77
T6	23.41	10.8	7.09	12.83	3503.33	38.37	14.74
T7	12.93	7.8	5.57	8.871	1454.43	30.23	8.95
T8	19.81	9.2	6.06	10.75	2445.20	33.77	13.75
C.D.(P=0.05)	0.30	0.18	0.31	1.36	669.35	8.77	0.33
C.V	0.94	1.21	3.04	13.30	12.05	5.04	1.80

Table 3: Effects of different growing media and biocapsule on biomass production and survival per cent of papaya plant.

Treatments	Fresh weight of shoot (g)	Fresh weight of root(g)	Dry weight of shoot (g)	Dry weight of root(g)	Root:Shoot ratio	Total biomass (g)	Survival percentage%
T0	3.1	1.32	0.8	0.21	0.24	1.03	58.67
T1	3.42	1.29	1.6	0.26	0.17	1.86	61.33
T2	10.04	4.09	3.88	0.72	0.18	4.6	81.33
T3	7.69	2.48	2.4	0.32	0.12	2.8	69.33
T4	9.62	3.61	3.26	0.46	0.14	3.72	74.7
T5	10.03	3.78	3.60	0.52	0.14	4.11	77.33
T6	10.56	4.29	4.02	0.73	0.18	4.73	82.67
T7	6.59	2.24	2.11	0.26	0.13	2.36	65.33
T8	7.72	3.13	2.84	0.36	0.12	3.2	69.33
C.D.(P=0.05)	0.51	0.25	0.23	0.06	0.04	0.21	10.29
C.V	3.87	4.96	4.77	8.34	15.61	3.79	8.36

Conclusion:

On the basis of present investigation, it is concluded that the treatment M6 (Soil+FYM+Vermicompost+Biocapsule) was found to be the most suitable over all the other treatments in relation to germination and seedling growth parameters (seedling height, no. of leaves, stem girth, fresh and dry weight of seedling) of Papaya.

The treatment M6 (Soil+FYM+Vermicompost+Biocapsule) also recorded the maximum survival percentage after transplanting.

References:

- Abirami K., Rema J., Mathew, P. A. Srinivasan, V. and Hamza, S. 2010.** Effect of different propagation media on seed germination, seedling growth and vigour of nutmeg (*Myristica fragrans* Houtt.). *Journal of medicinal plants research*,4(19) : 2054- 2058.
- Annapurna, D., Rathore, T.S. and Joshi, G. 2005.** Refinement of potting medium ingredients for production of high quality seedlings of Sandalwood (*Santalum album* L.). *Australian Forestry*, 68(1) : 44-49.
- Anonymous, 2014.** Indian Horticulture Database, National Horticulture Board, Ministry of agriculture, Government of India.
- Aracon, N.Q., Lee, S., Edward, C.A. and Atiyen, R. 2004.** Effect of humic acid derived from cattle, food and paper waste vermicompost on growth of green house plants. *Pedobiologia*, 47:741-744
- Asciutto, K., Rivera, M.C., Wright, E.R., Morisigue, D., and López, M.V. 2006.** Effect of vermicompost on the growth and health of *Impatiens wallerana*. *International Journal of experimental botany*,75: 115-123.
- Asmah, R., Rozita, R., Wan, N., L'zzah W.M., Zain, S.E. and Huzzaimah, A.S. 2002.** Antiproliferative activity of pure lycopene compared to both extracted lycopene and juices from watermelon (*Citrullus vulgaris*) and papaya (*Carica papaya*) on human breast and liver cancer cell. *Linus Journal Medicinal Science*,2(2):55-58.
- Bachman, G.R. and Metzger, J.D. 2007.** Growth of bedding plants in commercial potting substrate amended with vermicompost. *Bioresource Technology*, 99: 3155-3161.
- Baiyeri, K.P. and Mbah, B.N. 2006.** Effects of soilless and soil-based nursery media on seedling emergence, growth and response to water stress of African Breadfruit (*Treculia africana* Decne). *African Journal of Biotechnology*, 5(15):1405-1410
- Beattie, D.J. and White, J.W. 1992.** Liliaceae. Hybrids and species In: De Hertogh A.A. and Le Naed M (eds). The physiology of flower buds. *Elsevier, Amsterdam*, 423-454
- Bhagat, S., Thakur, A. and Dhaliwal, H.S. 2013.** Organic amendments influence growth, buddability and budding success in rough lemon (*Citrus jambhiri* Lush.). *Biological Agriculture & Horticulture: An International Journal for Sustainable Production Systems*, 29 (1) : 46-57.
- Bhardwaj, R.L. 2013.** Effect of nine different propagation media on seed germination and the initial performance of papaya (*Carica papaya* L.) seedlings. *Journal of Horticultural Science & Biotechnology*,88(5) : 531-536
- Chaudhary, M.I. 1996.** Soil and Fertilizer. In: *Horticulture (Ed.): M.N. Malik. pp. 259-260., National Book Foundation, Islamabad, Pakistan.*
- Chako, E.K. and Singh, R.N. 1966.** Studies of the longevity of papaya, Phalsa, Guava and Mango seeds. *International Seed Testing Association* 36:147-158.
- Donaka Dayeswari , Surendranath Rayaprolu and Auxilia Jone 2017** Effect of Potting Media on Seed Germination, Seedling Growth and Vigour in TNAU Papaya Co.8 (*Carica papaya* L.) ISSN: 2320 – 7051 *Int. J. Pure App. Biosci.* 5 (3): 505-512 (2017)
- Donaka Dayeswari , Surendranath Rayaprolu and Auxilia Jone 2017** Effect of Potting Media on Seed Germination, Seedling Growth and Vigour in TNAU Papaya Co.8 (*Carica papaya* L.) ISSN: 2320 – 7051 *Int. J. Pure App. Biosci.* 5 (3): 505-512 (2017)
- Gupta, S.N. and Naik, K.B. 2007.** In: *Instant Horticulture, Jain Brothers, New Delhi.*
- Hartman, H.T., Kester, D.E., Davies, F.T. and Geneve, R.L. 2001.** Plant propagation: Principles and practices, 7th Edition, *Prentice Hall publishers, New Jersey.*
- Irsa Shafique, Saiqa Andleeb, Malik Saim, Aftab Farrukh, Naeem Shaukat, Ali Summaya, Yahya Fayaz, Ahmed Tauseef, Tabasum Tariq Sultan, Beenish Shahid, Abdul Hameed Khan,**

Ghafoor ul Islam, Wajid Arshad Abbasi 2021. Efficiency of cow dung based vermi-compost on seed germination and plant growth parameters of *Tagetes erectus* (Marigold). *Heliyon Volume 7, Issue 1*.

Kumararvind, Ritesh; Patel, K. M.; Upadhyay, N. V 2020. Effect of Different Growing Media and Containers on Germination and Establishment of Seedlings of Papaya (*Carica papaya L.*) Cv. Madhubindu. *Society for Advancement of Science and Rural Development*.

Malay Samir, Ratna Rai and Birendra Prasad 2016. Effect of organic manures on seed germination and seedling growth of khirni. *Indian Forester, 142 (7 : 666-669, 2016*

M.R. Chiranjeevi, Shivanand Hongal, G.M. Vinay, B.M. Muralidhara and M.K. Sneha 2018. Influence of Media and Biofertilizers on Seed Germination and Seedling Vigour of Aonla. *International Journal of Current Microbiology and Applied Sciences ISSN: 2319-7706 Volume 7 Number 01 0*

M.S.Gawankar, P.M.Haldankar, P.C.Haldavanekar, B.R.Salvi, M.M. Burondkar, C.D. Pawar and B. M. Jamadagni 2021. Effect of Potting Media on Seed Germination and Seedling Vigour of Jackfruit (*Artocarpus heterophyllus Lam.*) under the Konkan Region of Coastal Maharashtra. *J. Indian Soc. Coastal Agric. Res. 39(1): 61-65*.

Naresh Kumar, K. HandaI, Inder Dev, Asha Ram, A. R. Uthappa, A. Shukla Lal Chand 2018. Effect of pre-sowing treatments and growing media on seed germination and seedling growth of *Albizia lebbek(L.) Benth.* *Journal of Applied and Natural Science 10(3): 860 -86ISSN : 0974-9411 (Print), 2231-5209*

Parasana, J. S. ; Ray, N. R. ; Satodiya, B. N. ; Patel, K. A. ; Panchal, G. P. 2012. Effect of mixture of growing media on germination and seedlings growth of different mango (*Mangifera indica L.*) cultivars under net house conditions. *Asian Journal of Horticulture Vol.7 No.2 pp.409-411 ref.8*

Prabha, K.P., Loretta, Y.L. and Usha, R.K. 2007. An experimental study of vermin-biowaste composting for agricultural soil improvement. *Bioresource Technology, 99: 1672-1681*.

Ritesh .K; Patel K: M; Upadhyay N.V. 2016. Effect of different growing media and containers on germination and establishment of seedlings of papaya (*Carica papaya L.*) cv. Madhubindu. *Trends in Biosciences, 8(1):227-23*

R.M. Atiyeh, S. Subler, C.A. Edwards, G.Bachman, J.D.Metzger, W.Shuster 2004. Effects of vermicomposts and composts on plant growth in horticultural container media and soil. *Pedobiologia Volume 44, Issue 5*

Sanjay Kumar Nagar, N. J. Vihol, Ritesh Kumar Arvind and Shashi 2016. Effect of growing media on germination and establishment of seedlings of papaya (*Carica papaya L.*) under net house conditions. *The bioscan 11(3): 1465-1468, 2016*

Shukla, A.K., Kaushik, R.A., Mahavar, L.N., Pareek, S., Pandey, and Sarolia, D.K. 2008. Fruits area production and importance. *In: Adhunikpaloyadan: Sanchar Kendra, MPUAT (Udaipur) P:09*.

Sukhjit Kaur 2017 Effect of Growing Media Mixtures on Seed Germination and Seedling Growth of Different Mango (*Mangifera indica L.*) Cultivars under Submountaneous Conditions of Punjab. *Chemical Science Review and Letters ISSN 2278-6783*

Wilson, S.B., Stofella, P.J. and Graetz, D.A. 2001. Use of compost as a media in amendment for containerized production of two subtropical perennials. *Journal of environmental Horticulture, 19(1):37-42*.

Talashilkar, S.C., Bhangarath, P.P. and Metha, V.B. 1999. Changes in chemical properties during composting of organic residues as influenced by earthwormactivity. *Journal of the Indian society of soil science, 47:50-53*

Venkatesh, L. ; Lakshmipathaiiah, O. R. ; Nagarajaih, C 2009. Effect of potting media on germination and seedling growth of *Pterocarpus marsupium*. *Environment and Ecology Vol.27 No.2 pp.563-564 ref.3*

Yahira, M. 1979. Effects of seed pretreatment on the promotion of germination in Papaya (*Carica papaya.L.*). *Memories of the faculty of agriculture, Kago Shima University, 15:4*

UNDER PEER REVIEW