

Original Research Article

Effect of organic manures and spacing on growth and yield of sweetcorn (*Zea mays saccharata* L.)

Comment [I1]: No italic

ABSTRACT

Comment [I2]: Check spelling and correct it

A field experiment was conducted during *Rabi* 2021-22 at Crop Research Farm, Department of Agronomy, Nani Agricultural Institute, Sam Higginbottom University of Agriculture Technology And Sciences, Prayagraj, Uttar Pradesh, India. The experiment was laid out in Randomized Block Design with nine treatments each replicated thrice on the basis of one year experimentation. The treatments consisted of FYM 18t/ha, Poultry manure 3t/ha, Vermicompost 3t/ha and three levels of spacing 40cm×25cm, 50cm×25cm, 60cm×25cm. The results showed that treatment-6 (Poultry manure 3t/ha + 60 cm x 25 cm) was recorded significantly higher plant height (176.59 cm), maximum number of leaves/plant (14.37) and higher plant dry weight (144.03 g/plant). Whereas the maximum number of cobs/plant (2.60), maximum cob length (16.57 cm), maximum girth of cob/plant (15.37), maximum cob weight with husk (323.90 g), maximum cob weight without husk (252.10 g), higher cob yield with husk (13.15 t/ha), higher cob yield without husk (8.51 t/ha).

Comment [I3]: and

Comment [I4]: tonha⁻¹

Comment [I5]:

Comment [I6]:

Comment [I7]: i.e.:

Comment [I8]: include observation variables

Comment [I9]:

Comment [I10]:

Comment [I11]:

Key words: Sweet corn, FYM, Poultry manure, Vermicompost, Spacing, growth and yield.

Comment [I12]: Write in full

INTRODUCTION

Sweet corn (*Zea mays Saccharata L.*) is a versatile crop, which finds a place in the human diet, animal feed, fodder and industrial raw material. Specialty corn such as baby corn and sweet corn have emerged as alternative food sources, especially for the rich society. Sweet corn is used as a human food in the soft dough stage with succulent grain. The higher content of water-soluble polysaccharide in the kernel adds texture and quality in addition to sweetness (Venkatesh et al., 2003). Maize (*Zea mays L.*) is the third most important cereal crop after wheat and rice, in the world. Maize is cultivated in tropical, sub-tropical and temperate countries of the world. Among various specialty corns, sweet corn is a mutant type with one or more recessive alleles in homozygous condition, which enables the endosperm to accumulate twice the sugar content as that of seed corn and controls the conversion of sugar into starch inside the endosperm of kernel Pande (2014).

In the world Maize is produced in an area of 193.7 million hectare with 12% of world food crop production of 1.1 billion tonnes and the productivity is 5.75 tonnes/hectare. In India, maize is produced in an area of 8.15 million hectares with 27 billion tonnes production and an average productivity of 3.8 tonnes/hectare. In the state of Uttar Pradesh maize is produced in an area of 3.05 lakh hectares with the production of 18.62 lakh tonnes and the productivity is 6 tonnes/hectare (FAOSTAT, 2022).

In order to achieve higher cob yields, maintenance of plant spacing is the most important factor. A spatial arrangement of plant governs the shape and size of the leaf area per plant, which in turn influences efficient interception of radiant energy, proliferation, growth of roots and their activity. Maximum yield can be expected only when plant population allows individual plant to achieve their maximum inherent potential. Thus, there is need to work out an optimum population density by adjusting inter and intra row spacing in relation to other agronomic factors (Naik et al., 2019).

Yields in organic farming are lower than chemical farming during initial years of practice and it takes few years to stability the yields. However, in the longrun, if properly followed, yields with organic farming would be far greater than those obtained with chemical farming. Huge quantities of organic materials such as farm yard manure, poultry manure, vermicompost, green manures and crop residues can substitute the inorganic fertilizers to a large extent to maintain the productivity and environmental quality Pande (2014).

Comment [I13]: delete

Comment [I14]: delete

Comment [I15]: Pande et al., 2014

Comment [I16]: see reference

Comment [I17]: tonha⁻¹

Comment [I18]:

Comment [I19]: or FAO, 2021?

Comment [I20]: not listed in the reference

Comment [I21]: Include the spacing that is commonly used or variations in the spacing of previous studies on corn as the reason for choosing the spacing in the research method.

Comment [I22]: use the latest similar reference

Comment [I23]: (Pande et al., 2014.)

Comment [I24]: see reference

Farmyard manure (FYM) is an important organic resource for agricultural production in livestock-based farming systems in the semi-arid tropics of India. However, Farm yard manure application to the crops is being practiced for long period. Well decomposed FYM in addition to supplying plant nutrients acts as binding material and improves the soil physical properties (Jayant Kumar Paikra 2018).

Comment [I25]: use one syllable, example: Jayant et al., 2018

Comment [I26]: check the reference, not a single author

Poultry manure, sometimes called chicken manure, is an excellent soil amendment that provides nutrients for growing crops and also improves soil quality when applied wisely, because it has high organic matter content combined with available nutrients for plant growth. Poultry waste consists of droppings, wasted feed, broken eggs, feathers, and sometimes sawdust from poultry floor. It also includes the dead birds and hatchery waste, all of which are high in protein and contain substantial amount of calcium and phosphorus due to high level of mineral supplement in their diet. Poultry manure has been reported to contain more plant nutrients than all other organic manures (Ali, 2005). Hence the present investigation was carried out to find out the "Effect of organic manures and spacing on growth and yield of sweet corn (*Zea mays Saccharata L.*)"

Comment [I27]: which part is Ali's statement? sentences and statements are too long and difficult for readers to understand

Comment [I28]: include the results of the latest research related to the use of chicken manure to support the determination of the treatment dose in the research method

MATERIALS AND METHODS

A field experiment was conducted during *Rabi* season of 2021-22, at Crop research farm of Department of Agronomy at Sam Higginbottom University of Agriculture, Technology, and Sciences, Prayagraj (U.P) India, which is located at 25° 24' 42" N latitude, 81° 50' 56" E longitude and 98 m altitude above the mean sea level (MSL). The crop was sown in 14 december 2021. The experiment was laid out in Randomized Block Design comprising of 9 treatments which are replicated thrice. Each treatment net plot size is 9m². Organic manures were applied and spacing was maintained as per the treatment details in combinations as follows, T₁: FYM 18t/ha + 40 cm x 25 cm, T₂: FYM 18 t/ha + 50 cm x 25 cm, T₃: FYM 18 t/ha + 60 cm x 25 cm, T₄: Poultry manure 3 t/ha + 40 cm x 25 cm, T₅: Poultry manure 3 t/ha + 50 cm x 25 cm, T₆: Poultry manure 3 t/ha + 60 cm x 25 cm, T₇: Vermicompost 3 t/ha + 40 cm x 25 cm, T₈: Vermicompost 3 t/ha + 50 cm x 25 cm and T₉: Vermicompost 3 t/ha + 60 cm x 25 cm. The growth parameters and yield attributes were recorded at harvest from randomly selected plants in each plot. The data was computed and analysed by following statistical method of Gomez and Gomez (1984).

Comment [I29]: include the stages or research procedures in full so that readers can understand

Comment [I30]: Example: land preparation, research plotsect

Comment [I31]: include observation variables, observation time, data analysis, use of further tests if the data is significant etc.

Comment [I32]: ton ha⁻¹

Comment [I33]: fix everything

Comment [I34]: delete

Comment [I35]: (Gomez and Gomez, 1984).

RESULTS AND DISCUSSION

Effect of Organic manures and Spacing on growth parameters of sweetcorn.

Plant height (cm)

It is evident from Table-1 that significantly highest plant height (176.59 cm) was recorded in the treatment6(Poultry manure 3 t/ha + 60 cm x 25 cm). However, the treatment5(Poultry manure 3 t/ha + 50 cm x 25 cm) and treatment9(Vermicompost 3 t/ha + 60 cm x 25 cm) were found to be statistically at par with treatment 6(Poultry manure 3 t/ha + 60 cm x 25 cm). Significant increase in plant height was with the application of poultry manure 3t/ha might be due to chicken manure produced longer plants. This increase in crop height can be attributed to the continued supply of nutrients in poultry manure. In addition, the development of shoot apical meristem may be a reason to gain better height. It appeared that the use of chicken manure enhanced the functions of the apical meristem, which led to an increase in height. The optimal height of the plants improves the acquisition of solar energy, which will help maintain photosynthesis Asfaw (2022).

Comment [I36]: adjust it with the way of writing in the previous suggestion

Comment [I37]: fix everything for each page and check it

Number of Leaves/plant

Significantly maximum number of leaves/plant (14.37) was recorded in the treatment 6 (Poultry manure 3 t/ha + 60 cm x 25 cm). However, treatment 5(Poultry manure 3 t/ha + 50 cm x 25 cm) and treatment 9(Vermicompost 3 t/ha + 60 cm x 25 cm) were found to be statistically at par with treatment 6(Poultry manure 3 t/ha + 60 cm x 25 cm). Significant and maximum number of leaves/ plant increased in poultry manure (3t/ha) might be due to the manure application will have a positive effect on plant growth and energy. This is because the leaves are the main components of photosynthesis in plants. The increase in the number of leaves was a precursor to a large number of assimilates and this allows for further transfer of grain. In addition, the high number of leaves in poultry manure treated plants contributes to better canopy production and weed compression. The poultry manure has high nutrition which promotes plant growth and makes nutrients available to the plant and poultry manure was found to improve leaf quality by providing an adequate amount of nutrients that accelerate leaf growth. Similar findings were reported by Asfaw (2022).

Comment [I38]: distinguish between the results of the study and the statement put forward by Asfaw

Comment [I39]: Asfaw's statement is used to support research results

Plant dry weight(g)

Significantly maximum dry weight (144.03 g/plant) was recorded in treatment 6 Poultry manure 3 t/ha + 60 cm x 25 cm. However, treatment 5 (Poultry manure 3 t/ha + 50

Comment [I40]: distinguish between the results of the study and the statement put forward by Asfaw

Comment [I41]: Asfaw's statement is used to support research results

Comment [I42]: add another reference or use more than one supporting reference, but not contradictory

cm x 25 cm) and treatment 9 (Vermicompost 3 t/ha + 60 cm x 25 cm) were found to be statistically at par with treatment 6 (Poultry manure 3 t/ha + 60 cm x 25 cm). Significant increase in plant dry weight was with the application of poultry manure 3t/ha might be due to the nutrients triggered the vigorous growth of plants thereby achieving higher leaf area, LAI and this further boosted the dry matter production and hastened the flowering and maturity period. Superior growth attributes obtained with high rates of Poultry manure **Uwah et al. (2014)**.

Comment [I43]: distinguish between the results of the study and the statement put forward by Uwah et al

Effect of organic manures and spacing on Yield and Yield Attributes of sweetcorn:

Number of Cobs/plant

Significantly maximum number of cobs/plant (2.60) was recorded with the treatment 6 (Poultry manure 3 t/ha + 60 cm x 25 cm). However, treatment 5 (Poultry manure 3 t/ha + 50 cm x 25 cm) and treatment 9 (Vermicompost 3 t/ha + 60 cm x 25 cm) were found to be statistically at par with treatment 6 (Poultry manure 3 t/ha + 60 cm x 25 cm). Significant and maximum number of cobs/plant is increased with the spacing of 60 cm x 25 cm might be due to greater availability of photosynthates, metabolites and nutrients to develop reproductive structures seems to have resulted in increased number of cobs/plant (**Naik et al. 2019**).

Length of Cob/plant (cm)

Significantly maximum length of cob/plant (16.57 cm) was recorded with the treatment 6 (Poultry manure 3 t/ha + 60 cm x 25 cm). However, treatment 5 (Poultry manure 3 t/ha + 50 cm x 25 cm) and treatment 9 (Vermicompost 3 t/ha + 60 cm x 25 cm) were found to be statistically at par with treatment 6 (Poultry manure 3 t/ha + 60 cm x 25 cm). Significant and maximum length of cob/plant is increased with the spacing of 60 cm x 25 cm might be due to increased number of leaves, leading to higher photosynthetic rate and accumulation of more assimilates which in turn increased the sink size. Higher nutrient uptake by sweet corn was also evident in the present investigation. The present findings are in line with the results obtained by **Kar et al. (2006)**.

Comment [I44]: include the results of the research or the statement so that it supports the results of this study

Girth of Cob/plant (cm)

Significantly maximum girth of cob/plant (15.37 cm) was recorded with the treatment 6 (Poultry manure 3 t/ha + 60 cm x 25 cm). However, treatment 5 (Poultry manure 3 t/ha + 50 cm x 25 cm) and treatment 9 (Vermicompost 3 t/ha + 60 cm x 25 cm) were found to be statistically at par with treatment 6 (Poultry manure 3 t/ha + 60 cm x 25 cm). Significant and

maximum girth of cob/plant is increased with the spacing of 60 cm×25 cm might be due to wider spacing provided uniform spread of plants because of less crowding which resulted into healthy cobs and thereby increases the weight, length of cob, and girth of corn **Neelam and Dutta (2018)**.

Comment [I45]: (Neelam and Dutta, 2018).

Cob weight (g)

a) With husk

Significantly maximum cob weight with husk (323.90 g) was recorded with the treatment 6 (Poultry manure 3 t/ha + 60 cm x 25 cm). However, treatment 5 (Poultry manure 3 t/ha + 50 cm x 25 cm) and treatment 9 (Vermicompost 3 t/ha + 60 cm x 25 cm) which were found to be statistically at par with treatment 6 (Poultry manure 3 t/ha + 60 cm x 25 cm).

b) Without husk

Significantly maximum cob weight without husk (252.10 g) was recorded with the treatment 6 (Poultry manure 3 t/ha + 60 cm x 25 cm). However, treatments 5 (Poultry manure 3 t/ha + 50 cm x 25 cm) and treatment 9 (Vermicompost 3 t/ha + 60 cm x 25 cm) were found to be statistically at par with treatment 6 (Poultry manure 3 t/ha + 60 cm x 25 cm). Significant and maximum cob weight is increased with the spacing of 60 cm × 25 cm might be due to the lesser competition for radiation and nutrients that allowed the plants to accumulate more biomass with higher capability to convert more photosynthates into sink resulting in higher values for the aforementioned attributes **Uwah (2014)**.

Cob yield (t/ha)

a) With husk

Significantly highest Cob yield (13.15 t/ha) was recorded with the treatment 6 (Poultry manure 3 t/ha + 60 cm x 25 cm). However, treatment 5 (Poultry manure 3 t/ha + 50 cm x 25 cm) and treatment 9 (Vermicompost 3 t/ha + 60 cm x 25 cm) were found to be statistically at par with treatment 6 (Poultry manure 3 t/ha + 60 cm x 25 cm).

Comment [I46]: add supporting references

b) Without husk

Significantly highest Cob yield (8.51 t/ha) was recorded with the treatment 6 (Poultry manure 3 t/ha + 60 cm x 25 cm). However, treatments 5 (Poultry manure 3 t/ha + 50 cm x 25 cm) and treatment 9 (Vermicompost 3 t/ha + 60 cm x 25 cm) were found to be statistically at

par with treatment 6 (Poultry manure 3 t/ha + 60 cm x 25 cm). Significant and highest cob yield is increased with the application of poultry manure 3t/ha might be due to increased availability of nutrients to the sweet maize crop at higher rate of poultry manure resulted in the production of longer ears, accompanied by increased grain filling that produced more grains/ear. superior ear and grain development was due to increase availability of adequate supply of nutrients from the poultry manure and greater production of photosynthates and their efficient translocation for development of reproductive parts The influence of Poultry manure on sweet maize yield parameters could be ascribed to the ability of this organic manure to supply nutrients and organic matter to the soil and in improving soil physical condition **Uwah (2014)**.

Comment [147]: distinguish between the results of the study and the statement put forward byUwah et al

UNDER PEER REVIEW

Table 1. Effect of Organic manures and Spacing on growth parameters of Sweet corn.

S.No	Treatments	Plant height (cm)	Leaves/plant	Dry matter (g plant ⁻¹)
1.	FYM 18 t/ha + 40 cm x 25 cm	173.08	13.44	141.51
2.	FYM 18 t/ha+ 50 cm x 25 cm	173.40	13.49	141.83
3.	FYM 18 t/ha+ 60 cm x 25 cm	174..89	13.97	142.82
4.	Poultry manure 3 t/ha+ 40 cm x 25 cm	174.39	13.87	142.54
5.	Poultry manure 3 t/ha + 50 cm x 25 cm	175.93	14.18	143.34
6.	Poultry manure 3 t/ha + 60 cm x 25 cm	176.59	14.37	144.03
7.	Vermicompost 3 t/ha + 40 cm x 25 cm	173.86	13.71	142.24
8.	Vermicompost 3 t/ha + 50 cm x 25 cm	175.43	14.06	143.13
9.	Vermicompost 3 t/ha + 60 cm x 25 cm	176.26	14.27	143.76
	SEm (±)	0.24	0.08	0.24
	CD (P 0.05)	0.71	0.23	0.73

Comment [148]: ton ha⁻¹

Comment [149]: check and repaired for all

Table 2. Effect of Organic manures and Spacing on Yield attributes and Yield of Sweet corn.

Treatments	No. of Cobs/plant	Length of Cob/plant (cm)	Girth of Cob/plant (cm)	Cob weight (g)		Cob yield (t/ha)	
				With Husk	Without Husk	With Husk	Without Husk
1. FYM 18 t/ha+ 40 cm x 25 cm	1.83	15.42	14.10	305.20	236.66	11.75	6.72
2. FYM 18 t/ha+ 50 cm x 25 cm	1.90	15.59	14.23	307.44	238.83	11.85	6.92
3. FYM 18 t/ha+ 60 cm x 25 cm	2.13	16.01	14.74	318.59	243.90	12.51	7.72
4. Poultry manure 3 t/ha+ 40 cm x 25 cm	2.00	15.90	14.54	315.18	242.60	12.24	7.43
5. Poultry manure 3 t/ha + 50 cm x 25 cm	2.30	16.35	15.11	321.44	248.61	12.89	8.08
6. Poultry manure 3 t/ha + 60 cm x 25 cm	2.60	16.57	15.37	323.90	252.10	13.15	8.51
7. Vermicompost 3 t/ha + 40 cm x 25 cm	1.97	15.68	14.44	313.10	240.66	12.04	7.08
8. Vermicompost 3 t/ha + 50 cm x 25 cm	2.23	16.15	15.01	311.99	245.78	12.76	7.92
9. Vermicompost 3 t/ha + 60 cm x 25 cm	2.43	16.47	15.25	322.17	250.47	13.04	8.33
F test	S	S	S	S	S	S	S
S. EM (\pm)	0.11	0.04	0.10	0.82	1.20	0.09	0.16
CD (P = 0.05)	0.34	0.13	0.29	2.46	3.59	0.27	0.49

Comment [150]: see please the suggest above

CONCLUSION

Based on the findings of the investigation it may be concluded that treatment with Poultry manure 3t/ha and spacing 60 cm x 25 cm performed exceptionally in all growth and yield parameters and in obtaining maximum cob yield of sweet corn. Hence, Poultry manure 3t/ha with spacing 60 cm x 25 cm may be more preferable and can be recommended to the farmers.

Comment [I51]: ton ha⁻¹

Comment [I52]:

REFERENCES

Ali, G.A. (2005). Uses of manure and fertilizer as soil management Technique for sustainable crop production. Paper presented at workshop organized by Taraba State Local Government Service Commission on 8 and 9, December 2005.

Asfaw, M.D. (2022). Effects of animal manures on growth and yield of maize (*Zea mays* L.) *Journal of Plant Science and Phytopathology*. **6**:033-039.

Comment [I53]: italic

FAO. 2021. World Food and Agriculture - Statistical Yearbook 2021. Rome.

Gomez, K.A. and Gomez, A.A., (1984). Statistical procedures for agricultural research (2 ed.). John Wiley and Sons, New York, 680

Jayant Kumar Paikra, Sanjay, K., Dwivedi and Harishankar. (2018). Effect of Integrated Nutrient Management on Productivity and Profitability of Sweet Corn (*Zea mays* L. saccharata) *ISSN: 2319-7692* Special Issue-6 pp. 2762-2769.

Comment [I54]: italic

Comment [I55]: delete

Comment [I56]: Enter the name of the journal

Kar, P. P., Barik, K. C., Mahapatra, R. K., Garnayak, L. M, Rath B. S, Bastia, D. K and Khanda, C. M. (2006). Effect of planting geometry and nitrogen on yield, economics and nitrogen uptake of sweet corn. *Indian Journal of Agronomy*, **51**(1): 43-45.

Neelam and Dutta, R. (2018). Production of Baby Corn as Influenced by Spacing and Nutrient Management. *International Journal of Current Microbiology and Applied Sciences*, **7**(12): 1332-1339.

Naik, A.A., Reddy, M.F., Babu, P.V.R., Kavitha,P. (2019) Effect of plant density and nitrogen management on growth, yield and economics of sweet corn (*Zea mays* var. Saccharata) *The Pharma Innovation Journal* **8**(6): 839-842

Comment [157]: italic

Pande, S., Naik, M.R., Naidu, S.M.M. (2014). Effect of Different Sources of Organic Manures on Sweet Corn (*Zea mays* Saccharata) *Environment & Ecology* **33** (2A): 810—813

Comment [158]: italic

Uwah, D.F., Ogar, C.C. (2014). Effects of Poultry Manure and Plant Population on Soil Properties and Agronomic Performance of Sweet Maize (*Zea mays* L. saccharata Strut) *International Journal of Applied Science and Technology***4**(4): 190-201.

Comment [159]: italic

Venkatesh S, Sanjay R, Shekhar JC. (2003). Sweet corn. Speciality corn Technical Series 1, Directorate of Maize Research New Delhi, 1-3.

UNDER PEER REVIEW