

## Original Research Article

### Effect of Drip Fertigation Levels on Soil Fertility and Productivity of Cauliflower cv.

~~PSBK-1~~ Pusa Snow Ball K-1

Comment [PA1]: Write it in long form

#### ABSTRACT

The field experiment were conducted during 2013-14 and 2014-15 to study the effect of drip fertigation on nutrient availability, nutrient uptake, plant parameters and yield of cauliflower. The field experiments were laid out in randomized block design with four replications and six fertigation levels (100, 90, 80, 70, 60 and 110 per cent of recommended dose). Fertigation with recommended dose of fertilizer resulted in highest available N (365.3 kg ha<sup>-1</sup>), P (95.8 kg ha<sup>-1</sup>) and K (477.8 kg ha<sup>-1</sup>). Drip fertigation with recommended dose also increased the available Zn, Cu, Fe and Mn content by 8, 29, 21 and 40 per cent, respectively over 60% RD treatment. The productivity of cauliflower was maximum with the application of 100 percent nutrients through drip irrigation and recorded an increase of 17.9 percent curd yield over 60 percent RD treatment. The study suggests that application of recommended dose of fertilizer through drip irrigation in 10 splits leads to higher and sustainable cauliflower production.

Comment [PA2]: No results for this

Comment [PA3]: Start with low to high %

Comment [PA4]: What is recommended dose in kg/ha

Comment [PA5]: What is RD? if is recommended dose should be introduced before in the previous sentences

**Key words:** Fertigation, drip system, water soluble fertilizer, nutrient uptake, cauliflower yield.

#### INTRODUCTION

The continuous improvement in productivity with optimum utilization of water, fertilizer and natural resources is essential for sustainability of any production system. Apart from the economic considerations, it is well established that the adverse effect of injudicious use of water and fertilizers can also have adverse implications on the environment. Hence, there is a need for technological interventions that will help in minimizing the use of precious resources (nutrient and water) and maximizing crop production without any detrimental effects on the environment. Among the various techniques of water and nutrient application, micro irrigation practices are very efficient and water conserving in nature. The drip or trickle irrigation is gaining much importance now a days due to the many unique advantages it provides like conservation of soil moisture, optimum utilization of water resources, lesser wastage of nutrient, proper and sustained water and nutrient availability to the crop. It delivers water and nutrients directly to the root zone of the crop and water is applied in precise amount which synchronizes with the requirement of the crop. The major advantages of fertigation are in saving of labour, appropriate timing of application of water and nutrients and their uniform distribution. Apart from other advantages like minimum leaching and volatilization losses, higher fertilizer use efficiency besides higher crop yields (Raina *et al.*, 2011).

Comment [PA6]: citation

Comment [PA7]: citation?

Comment [PA8]: citation

Comment [PA9]: citation

Comment [PA10]: use recent reference atleast of 2019-2022

Cauliflower is one of the important winter vegetable crop of mid hill region of Himachal Pradesh and is grown on an area of 79.5 thousand hectare with production of 1521.1 thousand metric tonnes and productivity of 19.1 MT/ha (NHB, 2013). The major constraints in cauliflower production include the traditional way of fertilizer application and lack of irrigation water. Drip irrigation is the most suitable method for uniform head size and high quality produce. Emphasis on the fertilizer application rate-source, timing and nutrient balance is critical for achieving good productivity of this crop. Most of the work on fertigation has been carried out using conventional NPK fertilizer. The information, on use of water soluble fertilizers, which has recently been introduced in the country especially for fertigation, is scanty. In view of the aforesaid facts, the present investigation was undertaken to study the availability of nutrient elements applied through drip irrigation system, leaf nutrient content, nutrient uptake and yield of cauliflower.

**Comment [PA11]:** Give studies conducted in conventional

**Comment [PA12]:** No results for this

**Comment [PA13]:** You need to review more literature on the other studies, on the effect of drip fertigation doses on nutrients uptake, availability and yield

## MATERIALS AND METHODS

### Study area

The experiment was conducted during two crop year (2013-2015) at the experimental farm of Department of Soil Science and Water Management, Dr. YS Parmar University of Horticulture and Forestry, Nauni, Solan (HP). The soil (Eutrochrept) was gravelly sandy loam in texture and neutral in reaction.

Include the map of study area and coordinates such longitude, latitude, altitude (masl), also climatic data

Soil sampling protocol/producers should be added

### Experimental design and treatments

The experiment was laid out with six treatments combinations replicated four times in RBD factorial design. The treatments comprised of six levels of nutrients through drip irrigation or fertigation viz. T<sub>1</sub> (100% of Recommended dose), T<sub>2</sub> (90% of Recommended dose), T<sub>3</sub> (80% of Recommended dose), T<sub>4</sub> (70% of Recommended dose), T<sub>5</sub> (60% of Recommended dose), T<sub>6</sub>

(110% of Recommended dose). Cauliflower seeds of *PusaSnow Ball K-1* variety were sown in well prepared raised nursery beds and one month old seedling were transplanted in the field at the spacing of 60 cm x 45 cm. Drip system comprised of five emitters in each row, placed at an angle of 180° to each other and at a distance of 23cm. The trials were conducted during the month of December to March in 2013-14 and 2014-2015 under temperate and dry condition. Twenty four raised plots of dimension 5 m x 1.8 m were made. The whole dose of FYM was applied before start of experiment. Each plot had four rows accommodating 33 plants and total 792 seedlings were transplanted in 24 plots. Recommended doses of N, P and K are 125, 108 and 76kg/ha, respectively for cauliflower in Himachal Pradesh. Fertigation was done through venturi in 10 equal split applications at 10 days interval. The water soluble fertilizer **NPK (WSF-19:19:19)** was used for fertigation. To meet the NPK requirement of different treatments, WSF was supplemented with urea and muriate of potash (MOP). After each fertigation, drip system was thoroughly flushed for 5 minutes. Different doses of fertilizers were applied by regulating the supply with the help of closing and opening knobs put at appropriate places. All the cultural operations were done as per package and practices and nutrients were applied as per the treatment details. Soil samples from 15 cm depth from each treatment/ plot were collected. The entire volume of soil was mixed thoroughly, air dried, sieved through 2mm sieve and stored in polythene bags for further analysis.

**Comment [PA14]:** Give the bases of selecting the 5 of RD? Why not 0, 20, 40, 60, 80, 100, 120% Sometime lowering of nutrients doses especially NPK level to 40, 60 could improve nutrients use efficiency

**Comment [PA15]:** Why you didn't include the control (0 NPK)? May be we could find that 0 application could have improved yield... may be the soil could supply the required amount

**Comment [PA16]:** Why this variety not others?

**Comment [PA17]:** Give the amount of FYM applied and the nutrition characteristics/contents

**Comment [PA18]:** To be seen in abstract

**Comment [PA19]:** Each RD give the amount of NPK for example 80% of 125 kg N ha =

**Comment [PA20]:** NPK ratio?

**Comment [PA21]:** At what stage of crop growth was the soil sample taken for analysis? How did you collect the samples, and how many samples per plot?

**Comment [PA22]:** Did you conduct soil analysis before experiment establishment?

Table Q

Recommended dose %	Amount of NPK Kg/ha
100	
90	
80	
70	
60	
110	

Formatted Table

**Comment [PA23]:** Put amount of NPK in terms of kg /ha

**Comment [PA24]:** Why no control in your experiment?

### Soil analysis

The soil samples were analysed for pH, EC, Organic carbon (OC) available N, P, K, Ca and Mg following standard methods. Micronutrient cation, i.e. Cu, Fe, Mn and Zn were determined by the method adopted by Lindsay and Norvell (1978), respectively. The physico-chemical properties of initial soil used in the present study are presented in (Table-1).

**Comment [PA25]:** Include these methods here (extract them from table 1)

[Put table 1 here](#)

### Collection of growth and yield data

The observations were recorded on different quantitative characters of cauliflower, viz. stalk length, number of leaves, gross weight, curd weight and curd yield. Curd yield ( $\text{qha}^{-1}$ ) was calculated from the weight of curds obtained per plot (kg). ~~The data recorded for various parameters under field experiments were subjected to statistical analysis using randomized block design (RBD) and treatment means were separated by critical difference at 5% level of significance (Gomez and Gomez 1984). The pooled data of chemical properties, nutrient status and plant parameters for two crop years have been presented.~~

**Comment [PA26]:** How did you collect these data? Tools used? Method of measurements??

### Data analysis

~~The data recorded for various parameters under field experiments were subjected to statistical analysis using randomized block design (RBD) and treatment means were separated by critical difference at 5% level of significance (Gomez and Gomez 1984). The pooled data of chemical properties, nutrient status and plant parameters for two crop years have been presented.~~

## RESULTS AND DISCUSSION

### *Chemical properties and nutrient status of soil*

Irrespective of treatments, the mean pH, EC and organic carbon was not significantly influenced by various treatments in both the years presented in (Table 2). However, different fertigation treatments significantly influenced the availability of nutrients. In general, the decreasing level of nutrients significantly decreased the content of available nutrients. The highest build up of available nitrogen was recorded under 100% RD (365.3 kg ha<sup>-1</sup>) which was statistically at par with T<sub>6</sub> (354.4 kg ha<sup>-1</sup>) and T<sub>2</sub> (352.0 kg ha<sup>-1</sup>). However, these treatments registered significantly higher available nitrogen content over the application of nutrients to the tune of 80, 70 and 60 percent of recommended dose. Treatment comprising fertigation with 100% RD increased the soil available nitrogen status by 20 per cent over the initial value recorded before execution of trial. Such an increase under fertigation with 90 and 110 per cent of RD were recorded to be 15.7 and 4.3 respectively. Quantity of N being added in balance application have helped higher available nitrogen status with 100% RD level. These results are in consonance with those of Fanish and Muthukrishnan 2013, Janagarathinam 2007 and Behera *et al.* 2013. Similarly, available P contents were relatively greater (95.84 kg ha<sup>-1</sup>) under 100 percent fertigation treatment. However, differences in available P among 100, 110 and 90 percent of RD were not significant. Significantly higher available P under 100% RD treatment may be due to the complete solubility of phosphorus source and frequent and small application rates through drip system. The findings corroborates with the results reported by Rauschkolbet *et al.* 1976 and Harjinder *et al.* 2004. It is also observed that effect of different levels of fertigation treatments on available K, closely followed available N and P in soil. The high availability of potassium may be due to majority of applied K was held in the surface soil and the downward movement was slow resulting in very small movement to deeper layer. The results are in agreement with the findings of Fanish and Muthukrishnan 2013, Zeng *et al.* 2000, Mmolawa and Or 2000, Mishra *et al.* 2006 and Suganya 2007. The data presented in (Table 3) indicated that imposition of different treatment did not influenced the availability of calcium, magnesium and micronutrient content of soil. However, higher content of available Ca (5.99 [cmol (p<sup>+</sup>)kg<sup>-1</sup>]), Mg (2.01 [cmol (p<sup>+</sup>)kg<sup>-1</sup>]), Zn (2.95 mg kg<sup>-1</sup>),

**Comment [PA27]:** Show level of significant p > 0.05 vice versa

**Comment [PA28]:** In methodology part put a table to show these amount check table named Q above

**Comment [PA29]:** Very long literature

**Comment [PA30]:** Arrange accordingly

Cu ( $3.99\text{mg kg}^{-1}$ ), Fe( $26.69\text{ mg kg}^{-1}$ ) and Mn ( $10.52\text{kg ha}^{-1}$ ) were recorded under 100% RD ( $T_1$ ) fertigation treatment.

*Put table 2 & 3 here*

### ***Plant parameter and yield***

The plant parameters of cauliflower i.e. number of leaves, gross weight, curd weight and yield was significantly influenced by different levels of fertigation (Table 4). The number of leaves among nutrient levels was significantly higher under  $T_1$ (19) which was at par with  $T_2$ ,  $T_3$  and  $T_6$ . The higher number of leaves with 100 percent RD treatment may be attributed to balanced nutrition, better water and nutrients utilization which results into more vegetative growth and higher photosynthetic rate. Similar findings on cauliflower crop have also been reported by Yangle and Tumbare (2014). These results are also in consonance with those of Olsen and Grevsen (2000) and Kageet *al.* (2001) and Aladakattiet *al.* (2012). The increase in gross weight was noted to be 18.5 per cent higher under 100% RD compared to 60% RD. This could be due to high yield and vegetative growth because of balanced fertilization in fractionated supplies have met the nutrient requirement of cauliflower at different growth stages. The curd weight is an important parameter that determines the commercial viability and acceptability of a variety and is one of the most important traits attaining highest consideration in research programs. Application of recommended level of nutrients ( $T_1$ ) produced maximum ( $873.8\text{ g plant}^{-1}$ ) curd weight which was at par with  $T_6$ ( $844.4\text{ g plant}^{-1}$ ). A comparison of data among different fertigation level reveals that  $T_1$ ,  $T_2$ ,  $T_3$ ,  $T_4$ ,  $T_5$  and  $T_6$  registered curd yield in the tune of 291.0, 274.5, 266.5, 254.3, 246.9 and  $281.2\text{g ha}^{-1}$ , respectively. The increase in curd yield was noted to be 18 per cent higher under 100% RD treatment over to 60% RD treatment. These results suggested that increasing level of fertigation significantly improved the curd yield. Significantly higher yield under 100% RD treatment may be attributed to much better water and nutrient utilization as nutrients were applied in ten equal splits. These fractionated supplies in

Comment [PA31]: Include some pictures/plates

Comment [PA32]: What about other %?

optimum nutrient concentration might have met the nutrient requirement of cauliflower at various growth stages thus leading to higher curd weight and yield. Kapoor *et al.* (2014), Sathya *et al.* (2008) and Brahma *et al.* (2010) observed significantly higher yield of tomato on fertigation. Similar observations were also made by Patel and Rajput (2003) in broccoli. Yanglem and Tumbare (2014) have also made the similar findings and attributed to such effect at higher fertigation level, crop meet out its nutritional requirement at respective growth stage which lead to luxurious growth and there by enhancement of yield in cauliflower.

**Table 1. Some important physico- chemical properties of the experimental soil**

Characteristics	Value	Methods/ References
pH ( soil: water 1:2)	6.56	1:2 Soil : water suspension, measured with digital pH meter (Jackson, 2005)
EC (soil : water 1: 2 )dSm <sup>-1</sup>	0.18	1:2 Soil : water suspension, measured with digital EC meter (Jackson, 2005)
Textural class	Gravelly sandy loam	Methods????
Organic carbon (%)	1.00	Walkley and Black wet digestion method (Walkley and Black, 1934)
Available N (kg ha <sup>-1</sup> )	264.5	Alkaline potassium permanganate method (Subbiah and Asija, 1956)
Available P (kg ha <sup>-1</sup> )	42.5	Olsen's method ( Olsen <i>et al.</i> , 1954)
Available K (kg ha <sup>-1</sup> )	377.3	Ammonium acetate method (Merwin and Peech, 1951)
Available Ca (cmol (p <sup>+</sup> ) kg <sup>-1</sup> )	4.31	Ammonium acetate method (Merwin and Peech, 1951)
Available Mg (cmol (p <sup>+</sup> ) kg <sup>-1</sup> )	1.41	Ammonium acetate method (Merwin and Peech, 1951)
DTPA Zn (mg kg <sup>-1</sup> )	2.58	DTPA extractant(Lindsay and Norvell,1978)
DTPA Cu (mg kg <sup>-1</sup> )	1.60	DTPA extractant(Lindsay and Norvell,1978)

**Comment [PA33]:** Shift this table to section of soil analysis



**Table 4. Effect of drip fertigation on plant growth and yield of cauliflower.**

Treatments	Stalk length (cm)	No. of leaves	Gross weight (g)	Curd weight (g plant <sup>-1</sup> )	Yield (q ha <sup>-1</sup> )
T <sub>1</sub>	5.0	19	1281.0	873.8	291.0
T <sub>2</sub>	4.7	18	1149.6	825.5	274.5
T <sub>3</sub>	4.6	18	1123.1	800.3	266.5
T <sub>4</sub>	4.3	17	1114.5	763.8	254.3
T <sub>5</sub>	4.2	16	1080.8	741.4	246.9
T <sub>6</sub>	4.8	18	1209.1	844.4	281.2
CD(0.05)	NS	1.08	65.16	41.64	13.92

## CONCLUSION

The results revealed that application of 100% RD of nutrients in ten splits through drip irrigation was found better in soil fertility status, nutrient uptake, plant parameter and curd yield over rest of the fertigation treatments. Compared with other treatments, use of 100% RD through fertigation resulted in significantly greater availability of nutrients, as also nutrient uptake that contributed to higher yield of cauliflower. Therefore it may be concluded that better soil health, higher and sustainable cauliflower production could be realized by applying 100% of recommended NPK in splits through drip irrigation.

Comment [PA37]: No results of this

## REFERENCES

- Aladakatti, Y. R., Palled, Y. B., Chetti, M. B., Halikatti, S. I., Alagundagi, S. C., Patil, P. L., Patil, V. C., Janawade, A. D. 2012. Effect of nitrogen, phosphorus and potassium levels on growth and yield of stevia (*Stevia rebaudiana* Bertoni). Karnataka Journal of Agricultural Sciences. 25(1), 25–29.
- Behera, M. S., Verma, O. P., Mahapatra, P. K., Singandhupe, R. B., Kumar, A., Kannan, K., Brahma, P. S. 2013. Effect of fertigation on stevia (*Stevia rebaudiana*) under drip irrigation. Indian Journal of Agronomy. 58(2), 243-250.
- Brahma, S., Borbora, D. P., Kachari, M., Hazarika, T. K., Das, K. 2010. Growth, yield and economics of broccoli under different levels of nitrogen fertigation. Indian Journal of Horticulture. 67, 279-282.
- Gomez, Gomez., 1984. Statistical procedures for agricultural research. John Wiley and sons. 627.
- Fanish, A. S., Muthukrishnan, P. 2013. Nutrient distribution under drip fertigation systems. World Journal of Agricultural Sciences. 9(3), 277-283.

- Herjinder, S., Narda, N. K., Chawla, J. K. 2004. Effect of irrigation frequency on subsurface drip irrigated vegetables. *Horticulture Technology*.13, 115-120.
- Jackson, M. L. 2005. *Soil Chemical Analysis*. Prentice Hall of India Pvt. Ltd., New Delhi.
- Janagarathinam, T. 2007. Optimization of water and nutrient requirement by drip fertigation for enhanced productivity of white onion (*Allium cepa*. var. *cepa* L.). Ph.D. Thesis. Tamil Nadu Agriculture University, Coimbatore.
- Kage, H., Alt, C., Stutzel, H. 2001. Predicting dry matter production of cauliflower (*Brassica oleracea* L. *botrytis*) under unstressed conditions: I) photosynthetic parameters of cauliflower leaves and their implication for calculation of dry matter production. *Scientia Horticulturae*. 87(3), 155-170.
- Kapoor, R., Sandal, S. K., Sharma, S. K., Kumar, A., Saroch, K. 2014. Effect of varying drip irrigation levels and NPK fertigation on soil water dynamics, productivity and water use efficiency of cauliflower (*Brassica oleracea* var. *botrytis*) in wet temperate zone of Himachal Pradesh. *Indian Journal of Soil Conservation*. 42(3), 249-254.
- Lindsay, W. H., Norvell, W. A. 1978. Development of DTPA soil test for Zn, Fe, Mn and Cu. *Soil Science Society of American Journal*. 42, 420-428.
- Merwin, H. D., Peech, M. 1951. Exchangeability of soils potassium in the silt and clay fractions as influenced by the nature of the complimentary exchangeable cations. *Soil Science Society of American Proceedings*.15, 125-128.
- Mishra, P., Singh, D. K., Ravi, Babu, R. 2006. Impact of method of irrigation and fertilizer application on distribution of potassium in soil under radish. *Institution of Engineers (India) Journal*.87, 20-23.
- Mmolawa, K., Or, D. 2000. Root zone solute dynamics under drip irrigation: A Review. *Plant and Soil*. 222, 163-190.
- NHB. 2013. Annual production of vegetable. National Horticulture Board.
- Olesen, J. E., Grevsen, K. 2000. A simulation model of climate effects on plant productivity and variability in cauliflower (*Brassica oleracea botrytis* L.). *Scientia Horticulturae*. 83(2): 83-107.
- Olsen, S. R., Cole, C. V., Watanable, F. S., Dean, L. A. 1954. Estimation of available phosphorous in soil by extraction with sodium bicarbonate. *USDA Circular* 939, 1-19.
- Patel, N., Rajput, T. B. S. 2003. Yield response of some vegetable crops to different levels of fertigation. *Annals of Agricultural Research*.24, 542-45.
- Raina, J. N, Sharma, Tarika, Shashi, Suman. 2011. Effect of drip fertigation with different fertilizers on nutrient distribution in soil, leaf nutrient content and yield of apricot (*Prunus armeniaca* L.). *Journal of the Indian Society of Soil Science*.59, 268-277.
- Rauschkolb, R. S, Rolsten, D. E., Miller, R. J., Carlton, A. B., Burau, R. G. 1976. Phosphorus fertilization with drip irrigation. *Soil Science Society of America Journal*.40, 68-72.

- Sathya, S., Pitchai, J. G., Indirani, R., Kannathasan, M. 2008. Effect of fertigation on availability of nutrients (N, P & K) in soil-A Review *Agricultural Reviews*. 29(3), 214-219.
- Subbiah, B. V., Asija, G. L. 1956. Rapid method for estimation of available nitrogen in soils. *Current Science*. 25, 259-260.
- Suganya, S., Anitha, A., Appavu, K. 2007. Moisture and nutrient distribution system under drip fertigation systems. In: Third International Ground Water Conference on "Water, Environment and Agriculture-Present problems and future challenges". pp: 7-10.
- Walkley, A., Black, I. A. 1934. Examination of the method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Science*. 37, 29-38.
- Yangle, S. D., Tumbare, A. D. 2014. Influence of irrigation regimes and fertigation levels on yield and physiological parameters in cauliflower. *An International Quarterly Journal of Life Sciences*. 9(2), 589-594.
- Zeng, D. Q., Brown, P. H., Holtz, B. A. 2000. Potassium fertigation improves soil K distribution, Builds Pistachio yield and quality. *Fluid Journal*. 1-2.

UNDER PEER REVIEW