

Original Research Article

Evaluation of *Beauveria bassiana*, Neem oil and Selected Insecticides on Population of Armyworm *Spodoptera frugiperda* (J.E.Smith) on Maize (*Zea mays* L.)

Abstract

A field trial was conducted at Crop Research Farm (CRF), SHUATS Naini, Prayagraj, during Kharif 2021 in Randomized block design with seven treatments viz. Imidacloprid 17.8%SL, Thiodicarb 75% WP, Spinosad 45% SC, Emamectin benzoate 5% SG, Neem oil, *Beauveria bassiana* 1×10^8 CFU/ml, Thiamethoxam 12.6% + Lambda cyhalothrin 9.5 %ZC and untreated control in three replication. Data were taken on fall armyworm population. The larval population of fall armyworm *Spodoptera frugiperda* (J.E.Smith) on three, seven and fourteen days after spraying revealed that the treatment Emamectin benzoate 5% SG (5.15) proved to be the most effective treatment followed by Thiodicarb 75% WP (6.44), Thiamethoxam 12.6% + lambda cyhalothrin 9.5 %ZC (7.41), Spinosad 45% SC (8.23), Imidacloprid 17.8%SL (8.79). whereas, Neem oil 3% (10.24) and *B. bassiana* 1×10^8 CFU/ml (10.93) were of least effective against *S. frugiperda*. plots treated with Emamectin benzoate 5% SG showed highest yield (35.31q/ha), followed by Thiodicarb 75% WP (34.21q/ha), Thiamethoxam 12.6% + Lambda cyhalothrin 9.5 %ZC (32.58q/ha), Spinosad 45% SC (32.08q/ha), Imidacloprid 17.8%SL (31.21qt/ha), Neem oil 3% (29.63q/ha) and *B. bassiana* 1×10^8 CFU/ml (28.02), as compared to control (22.44q/ha).

Keywords: Benefit-Cost Ratio, Efficacy, Fall of ,armyworm, Infestation, Maize.

INTRODUCTION

Maize (*Zea mays* L.) is the second most cereal crop belongs to Poaceae family. It is one of the most flexible growing crops with greater adaptability to different agro-climatic conditions. Because of higher genetic yield potential among the cereals, this crop is globally known as the Queen of cereals. Maize kernel is an edible and nutritive part of the crop. Maize kernel contains vitamin C, vitamin E, vitamin K. Potassium is a major nutritional content present, which has a good significance because an average human diet is deficient in it (Kumar and Jhariya, 2013).

Recently, the occurrence of a new invasive pest *Spodoptera frugiperda* (J.E. Smith), a lepidopteron insect has been suspected on maize crop in Karnataka (Shylesha et al., 2018). The insect is native to the tropical and sub-tropical regions of North, and South America, where it has been considered a key pest in maize and several other crops for decades (Smith et al., 1797) (Its very very old, write recent reference). Fall armyworm was detected for the first time on the African continent during January 2016 in Nigeria, and by 2019 had been reported in almost all of sub-Saharan Africa, as well as Southeast Asia, causing substantial yield (Diva et al., 2021). The caterpillar feeds on all plant stages by consuming the foliage and mostly prefers premature corn (Moraes et al., 2015). In the event of food depletion and crowding, larvae march out of crop in search of food, which gives them name Fall armyworm. The densities of caterpillar reduced due to their cannibalistic behaviour (Capinera et al., 2008, Sisodiya et al., 2018). *S. frugiperda*, is a polyphagous migratory insect that is able to cause considerable economic losses in over 80 different crops (Bohnenblust et al., 2014). FAW was observed to cause up to 72 % yield loss in maize.

Maize is most vulnerable to fall armyworm, *S. frugiperda*, which causes severe losses to it. Though, application of effective chemicals and botanicals with different mode of action at proper crop stage is significant for its management. The applications of various insecticides with different mode of action strengthen insecticides resistance management strategy. Therefore, keeping in view the aforementioned facts, the present investigation was carried out with the aim to develop a new management strategy for control of *S. frugiperda* at farmer's field economically.

MATERIALS AND METHODS

The experiment was conducted at Central Research Farm (CRF), Department of Entomology, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (UP) during 2021 during Kharif season. The experiment was laid out using GA-85 maize cultivars. The experiment was sown in Randomized Block Design with three replications consisting of 7 treatments having one absolute control. Five insecticides, Neem oil and one entomopathogenic fungus *Beauveria bassiana* were used. Treatments comprising of Imidacloprid 17.8% SL (T1), Thiodicarb 75% WP (T2), Spinosad 45% SC (T3), Emamectin benzoate 5% SG (T4), Neem oil 3% (T5), *B. bassiana* 1×10^8 CFU/ml (T6), Thiamethoxam 12.6% + Lambda cyhalothrin 9.5% ZC (T7). The seed rate of 20 kg / ha (very very low amount) was utilized to cultivate the crop. Plots (2×2m) was made. Sowing was done with 30 cm × 10 cm ha (very very low space) spacing and applied dose of farm yard manure was 20t/ha and N, P, K were 50, 25 and 25kg/ha, (what is the kind of these fertilizers , concentration of each element and these amounts are very very low) respectively. The population of *Spodoptera frugiperda* (J.E. Smith) was recorded one day before spraying and on 3rd, 7th and 14th day after insecticidal application. The obtained data were subjected to statistical analysis. The populations of *S. frugiperda* was recorded on 5 randomly selected and tagged plants from each plot for investigating larval population and cost benefit ratio.

The cost of insecticides and biocides used in these experiments was obtained from nearby market. The total cost of plant protection, consisted of plant protection, consisted of cost of treatments sprayer, rent and labour charges for the spray. There are two sprays throughout the research period and the overall plant protection expenses were calculated. Total income was realised by multiplying the total yield per hectare by prevailing market price, while the net benefit is obtained by subtracting the total cost of plant protection from the total income. Benefit over the control for each sprayed treatment was obtained by subtracting the income of the control treatment from that of each sprayed treatment.

RESULT AND DISCUSSION

Effect of different insecticides and biocides on the incidence of *Spodoptera frugiperda* (J.E. Smith) showed that all the treatments were significantly superior in reducing the infestation of fall of armyworm resulting in increasing the yield significantly as compared to control. The first spray was given after 40 days of planting. The larval population of fall armyworm on maize after first spray revealed that all the chemical treatments were significantly superior over

control. Among all the treatments lowest larval population, was recorded in Emamectin benzoate 5%SG(6.62) followed by Thiodicarb 75%WP (7.93), Thiamethoxam 12.6% + Lambda cyhalothrin 9.5 %ZC (8.99), Spinosad 45% SC (9.77), Imidacloprid 17.8%SL (10.223). Neem oil 3% (11.15). The treatment with *Beveria bassiana* 1×10^8 CFU/ml (11.92) was found to be least effective but comparatively superior over the control.

The second spray was applied 15 days after first spray and larval population was recorded. The data for second spray showed minimum larval population in Emamectin benzoate 5%SG (3.68) followed by Thiodicarb 75%WP (4.95), Thiamethoxam 12.6% + lambda cyhalothrin 9.5 %ZC (5.84), Spinosad 45% SC (6.706), Imidacloprid 17.8%SL (7.376) and Neem oil 3% (8.95). The treatment with *B. bassiana* 1×10^8 CFU/ml (9.95) was found to be least effective among all the treatments but comparatively superior over the control.

All the insecticides were found very effective and significantly over control. The data for overall mean larval population were recorded of which least larval population was recorded in Emamectin benzoate 5%SG (5.15), Thiodicarb 75%WP (6.4), Thiamethoxam 12.6% + Lambda cyhalothrin 9.5 %ZC (7.41), Spinosad 45% SC(8.23), Imidacloprid 17.8%SC (8.7), Neem oil 3% (10.24). The treatment with *B. bassiana* 1×10^8 CFU/ml (10.93) was least effective among all the treatments and control(22.34).

The highest yield and benefit cost ratio due to controlling army worm was recorded by Emamectin benzoate 5%SG (35.31q/ha) (1:1.80). Thiodicarb 75%WP (34.21 q/ha) (1:1.67), Thiamethoxam 12.6% + lambda cyhalothrin 9.5 %ZC (32.58q/ha) (1:1.57), Spinosad 45% SC (32.08q/ha) (1:1.56), Imidacloprid 17.8%SC (31.21q/ha) (1:1.43), Neem oil 3% (29.63 q/ha) (1:1.32). These results are in accordance with the obtained data by Kunkel *et al.*(1999) ; Mallapur *et al.*(2019); Deshmukh *et al.* (2020); Sangleet *et al.* (2020) and Thumaret *et al.*(2020). On the other hand, the treatment with *B. bassiana* 1×10^8 CFU/ml (28.02 q/ha) (1:1.21) was the lowest efficient one in this regard followed by Neem oil 3% (29.63 q/ha) (1:1.32). These results are supported by the obtained results by Wale *et al.* (2022) and Tulashie *et al.* (2021). Control treatment yielded poor yield (22.47q/ha) and (1:0.85).

Table 1. Efficacy of *Beauveria bassiana*, Neem oil and selected insecticides on population of fall armyworm *Spodoptera frugiperda* on maize during Kharif season of 2021

No.	Treatments	Population of <i>S. frugiperda</i>							Overall mean	Yield (q/ha)	B:C Ratio ***
		First spray				Second spray					
		1DBS *	3DAS **	7DAS	14DAS	3DAS	7DAS	14DAS			
T ₁	Imidacloprid 17.8% SL	16.80	11.40 ^{bc} _d	9.20 ^{cd}	10.40 ^c _d	9.20 ^d	5.60 ^c	7.33 ^{bc}	8.79	31.21	1: 1.43
T ₂	Thiodicarb 75% WP	17.10	9.33 ^E	6.47 ^F	8.00 ^F	6.73 ^F	3.60 ^{Ef}	4.53 ^{Ef}	6.44	34.21	1: 1.67
T ₃	Spinosad 45% SC	16.10	10.80 ^{cd}	8.60 ^{de}	9.60 ^{de}	8.26 ^{dc}	5.46 ^{cd}	6.40 ^{cd}	8.23	32.08	1: 1.56
T ₄	Emamectin benzoate 5% SG	16.60	7.73 ^f	5.53 ^f	6.60 ^g	4.93 ^g	2.73 ^f	3.40 ^f	5.15	35.31	1: 1.80
T ₅	Neem oil 3%	17.30	11.90 ^{bc}	10.2 ^{bc}	11.33 ^{bc}	10.50 ^c	8.80 ^b	7.53 ^{bc}	10.24	29.63	1: 1.32
T ₆	<i>B. bassiana</i> 1x10 ⁸ CFU/ml	13.80	12.46 ^b	10.9 ^b	12.46 ^b	11.60 ^b	9.80 ^b	8.40 ^b	10.93	28.02	1: 1.21
T ₇	Thiamethoxam 12.6%+lambda cyhalothrin 9.5% ZC	19.70	10.30 ^{de}	7.80 ^e	8.86 ^{ef}	7.46 ^{ef}	4.46 ^{de}	5.60 ^{de}	7.40	32.58	1: 1.57
T ₀	Control	21.30	21.60 ^a	21.8 ^a	22.13 ^a	22.53 ^a	22.70 ^a	23.38 ^a	22.34	22.47	1: 0.85
	F-test	NS	S	S	S	S	S	S	S	-----	-----
	S. Ed (±)	N/A	0.66	0.52	0.59	0.59	0.44	0.53	0.44	-----	-----
	C.D.(P =0.5)	-	1.412	1.114	1.256	1.256	0.95	1.13	0.94	-----	-----

*DBS=Day before spray, ** DAS= Day after spray , *** =Mention what is to mean.

UNDER PEER REVIEW

CONCLUSION

It may be stated that the synthetic insecticides and the botanical Emamectin benzoate 5%SG could be suggested for the management of *Spodoptera frugiperda* on maize plants, which proved to be most effective and economical. Similarly, the use of Thiodicarb 75% WP, Thiamethoxam 12.6% + lambda cyhalothrin 9.5 %ZC, Spinosad 45% SC, Imidacloprid 17.8%SL, can also be thought for the management of fall armyworm. However, the application of Neem oil 3% and *B.bassiana* 1×10^8 CFU/ml could not exert much encouraging role for fall armyworm management. These products help in reducing pollution in the environment, hence they can be suitably incorporated as treatment from an IPM perspective.

UNDER PEER REVIEW

REFERENCES

- Bohnenblust, E. W., Breining, J. A., Shaffer, J. A., Fleischer, S. J., Roth, G. W. and Tooker, J. F. (2014).** Current European corn borer, *Ostrinia nubilalis*, injury levels in the northeastern United States and the value of Bt field corn. *Pest management science*, **70**(11):1711-1719.
- Capinera, J. (2008).** Fall armyworm, *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae). *Encyclopedia of Entomology*, 1409-1412.
- Deshmukh, S., Pavithra, H. B., Kalleshwaraswamy, C. M., Shivanna, B. K., Maruthi, M. S. and Mota-Sanchez, D. (2020).** Field efficacy of insecticides for management of invasive fall armyworm, *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) on maize in India. *Florida Entomologist*, **103**(2):221-227.
- Divya, J., Kalleshwaraswamy, C. M., Deshmukh, S., Ambarish, S. and Sunil, C. (2021).** Evaluation of whorl application of insecticides mixed with sand against fall army worm *Spodoptera frugiperda* in maize. *Indian Journal of Entomology*, 1-5.
- Kumar, D. and Jhariya, A. N. (2013).** Nutritional, medicinal and economical importance of corn: A mini review. *Research Journal of Pharmaceutical Sciences*, **2**(7): 1-6.
- Mallapur, C. P., Naik, A. K., Hagari, S., Praveen, T. and Naik, M. (2019).** Laboratory and field evaluation of new insecticide molecules against fall armyworm, *Spodoptera frugiperda* (J.E. Smith) on maize. *Journal of Entomology and Zoology Studies*, **7**(4): 869-875.
- Moraes, A. R. Lourencao, A. L. and Paterniani, Meag. Z. (2015).** Resistance of conventional and isogenic transgenic maize hybrids to *Spodoptera frugiperda* (Lepidoptera: Noctuidae). *Bragantia*, **74**:50-57.
- Sangle, S. V., Jayewar, N. E. and Kadam, D. R. (2020).** Efficacy of insecticides on larval population of fall armyworm, *Spodoptera frugiperda* on maize. *Journal of Entomology and Zoology Studies*, **8**(6), 1831-1834.

- Shylesha, A. N., Jalali, S. K., Gupta, Ankita, Varshney, Richa., Venkatesan, T., Shetty, P. and Raghavendra, A. (2018).** Studies on new invasive pest *Spodopterafrugiperda* (J.E. Smith)(Lepidoptera: Noctuidae) and its natural enemies. *Journal of Biological Control*, **32**(3), 1-7.
- Sisodiya, D. B., Raghunandan, B. L., Bhatt, N. A., Verma, H. S., Shewale, C. P., Timbadiya, B. G. and Borad, P. K. (2018).** The fall armyworm, *Spodopterafrugiperda* (J.E. Smith)(Lepidoptera: Noctuidae); first report of new invasive pest in maize fields of Gujarat, India. *Journal of Entomology and Zoology Studies*, **6**(5), 2089-2091.
- Smith, J. E., (1797).** First record of the fall armyworm, *Spodopterafrugiperda* (Lepidoptera: Noctuidae). *African Entomology*, **26**(1): 244-246.
- Thumar, R. K., Zala, M. B., Varma, H. S., Dhobi, C. B., Patel, B. N., Patel, M. B. and Borad, P. K. (2020).** Evaluation of insecticides against fall armyworm, *Spodopterafrugiperda*(J.E. Smith) infesting maize. *International Journal of Chemical Studies*, **8**(4): 100-104.
- Tulashie, S. K., Adjei, F., Abraham, J. and Addo, E. (2021).** Potential of neem extracts as natural insecticide against fall armyworm (*Spodopterafrugiperda*(J.E. Smith)(Lepidoptera: Noctuidae)). *Case Studies in Chemical and Environmental Engineering*, **4**:1-7.
- Wale, S. D., Mahadik, S. S. and Hole, U. B. (2022).** Bio-intensive management of fall armyworm, *Spodopterafrugiperda* infesting maize. *Annals of Plant Protection Sciences*, **30**(1), 73-77.