

Original Research Article

Cell wall changes of papaya fruit treated with 1-Methylcyclopropene: postharvest analysis through SEM and FT-IR spectroscopy

ABSTRACT

1-Methylcyclopropene (1-MCP), is a widely researched and commercially used molecule to delay fruit ripening and improve the shelf life, both climacteric and non-climacteric fruits. Increasing the shelf life and keeping quality, especially of important climacteric fruits like papaya using 1-MCP, which undergoes a rapid softening during the ripening process required a detailed investigation of the structural changes of fruit cell wall. These changes include modifications in the cell wall polysaccharide structure due to increased activity of hydrolyzing enzymes, pectin, hemicellulose and cellulose degradation that results in fruit softening. Pectin degradation in papaya determines the extent of softening during ripening. Hence, the present investigation describes the role of 1-MCP in retaining the firmness, postharvest. The ultrastructural changes of papaya peel and pulp were studied using Scanning Electron Microscope (SEM). The pectin degradation in untreated and 1-MCP treated fruits were analyzed through a rapid and reliable technique, FTIR spectroscopy (Fourier transform-infrared (FT-IR)). The study revealed that postharvest application of 1-MCP at 900 ppb to papaya fruits, stored under cold storage (14°C) was effective in retaining higher fruit firmness by suppressing the rapid cell wall modifications and activity of pectin degradation, which are triggered by ethylene.

Keywords: *1-Methylcyclopropene (1-MCP), softening, ethylene, postharvest, Scanning Electron Microscope (SEM), Fourier transform-infrared (FT-IR).*

1. INTRODUCTION

Papaya (*Carica papaya* L.) is an important tropical, but a highly perishable climacteric fruit. Climacteric fruits undergo rapid changes during ripening process that include dramatic rise in respiration rate, faster peel color changes, rapid loss of firmness leading to softening and quality loss. Postharvest softening basically occurs due to the enhanced activity of cell wall degrading enzymes, triggered by ethylene (major hormone that regulates ripening associated changes), especially in climacteric fruits. A characteristic feature of the softening process, is degradation of the ordered arrangement of cell wall and middle lamella polysaccharides, due to an increase in the activity of the cell wall degrading/hydrolyzing enzymes (Dick and Labavitch, 1989; Zhao and Paull, 1996). Hence, in other terms cell wall modification

during ripening is accompanied by pectin solubilization and hemicellulose depolymerization (Paull et al., 1999; Ali et al., 2004). Pectin hydrolyzation is attributed to significantly affect fruit firmness, among the alterations in cell wall (Lazan et al., 1995). Retardation of faster ripening, associated rapid morpho-physiological and biochemical changes have been achieved by application of 1-Methylcyclopropene (1-MCP) in several climacteric and non-climacteric fruits. 1-MCP is a non-toxic molecule (approved by USEPA) that inhibit the ethylene action and improve shelf life of several fruits and vegetables. 1-MCP delays fruit ripening and softening by suppressing the activities of cell wall hydrolytic enzymes (Jeong et al., 2002). Hence, the present investigation is aimed at (1) studying the ultra-structural changes in the cell wall of papaya fruits as affected by 1-MCP, revealed through Scanning Electron Microscope technique (SEM) (2) pectin solubilization and degree of methyl esterification (DME) of pectin using Fourier transform-infrared (FT-IR) spectroscopy. FT-IR is a reliable, rapid and highly sensitive method to analyze the DME of pectin (Barros et al., 2002; Manrique and Lajolo, 2002).

2. MATERIALS AND METHODS

2.1 Application of 1-Methylcyclopropene

Papaya (TNAU var. CO.8) fruit harvested at color break stage were selected for uniformity, devoid of injury or diseases manually and sorted into two groups for application of 1-MCP treatments. 1-MCP application procedure is as previously described (Srividhya and Jeykumar, 2017). Fruits were then loaded into plastic crates and cooled down to attain a pulp temperature of 15-20°C (measured using stick thermometer). One batch was treated with 1-MCP at 900 ppb, while the other batch was left untreated. The fruit were pre-treated with 100 ppm ethylene to trigger the ethylene receptor in order to avoid rubbery texture as reported in previous studies (Manenoi et al., 2007). 1-MCP (obtained from Agrofresh Inc., Mumbai) at 900 ppb concentration was applied to papaya fruit by vaporization in a closed container for a duration of 14 hours @ 14°C, RH 90-95 per cent. After 1-MCP vaporization, half of the fruit from each treatment box were loaded into crates and kept at ambient temperature of 27±2°C and rest of the fruit were kept at cold storage (14°C). Thus, the treated and untreated fruits were allowed to undergo normal ripening at the respective storage conditions. The samples were then subjected to SEM and FTIR analysis. To determine the effects 1-MCP on papaya fruit morphology stored under ambient and cold storage, the peel samples were collected from all set of treatments on day seven (since ambient stored untreated fruit had achieved the last stage of consumer acceptance).

2.2 Ultrastructural peel changes using SEM

Ultrastructural changes revealed through SEM in papaya fruit peel can be complemented to cell wall changes as observed using FT-IR spectroscopy. The papaya fruit peel sample was collected from the middle portion of the fruit. At the time of analysis, sample was placed to a carbon tape and affixed to a carbon stub which was then vacuum desiccated for 2 min to remove excess moisture from the stub. The peel sample was then viewed under a scanning electron microscope (SEM; Quanta 250, FEI, Hillsboro, OR, USA) using an ETD detector. The SEM was operated at 5-15KV, with a working distance of 50 µm.

The SEM facility at Department of Nanoscience and Technology, TNAU, Coimbatore, India was utilized for this study.

2.3. Measuring the degree of methyl esterification (DME) of pectin using Fourier Transform-Infrared (FT-IR) spectroscopy

FT-IR spectroscopy was chosen to analyze the pectin properties in papaya fruit. FT-IR works on the principle that, absorption intensity of different functional groups changes at different wave numbers at near infra region, particularly the ionized carboxylate group (COO^-) and ester carbonyl group (C=O) group (Barros et al., 2002; Szymanska-Chargot and Zdunek, 2013), without hydrolyzing the ester linkage (Chatjigakis et al., 1998). Isolation of papaya peel and pulp pectin was done adopting the methods of Koubala *et al.* (2014). The dried powder obtained from the samples after blanching (95°C for 5 min) was used for the isolation of alcohol insoluble residue (AIR) cell-wall components as described in the protocol. Water soluble pectin (WSP) was extracted from AIR and used for the analysis following the procedure described by Manrique and Lajolo, (2002). Approximately, 10 mg of the WSP was dissolved in MilliQ water and pH was adjusted to 6 with 0.1N NaOH (since pH of 6 gives total ionization of carboxylic groups. The samples were then freeze dried for further analysis. The FT-IR (Fourier Transformation- Infrared Analysis) was performed using Jasco FT/IR 6800 model to measure the degree of methyl esterification of pectin in 1-MCP treated and untreated papaya fruit stored under different storage conditions. The standard curve was obtained using pure esterified pectin from Sigma. For DME determination, a sample was placed on sample holder devoid of any air. After loading the sample, it was analyzed in Attenuated Total Reflectance (ATR/FT-IR) and the spectral data were recorded at 64 scans second^{-1} with resolution of 4 cm^{-1} at the spectral range of $400\text{-}4000\text{ cm}^{-1}$. Spectra was changed to absorbance mode for baseline correction and peak splitting was done to correct for any protein interference. The intensity of peaks around $1740\text{-}1745\text{ cm}^{-1}$ indicates ester (C=O) stretching, and peaks around $1630\text{-}1600\text{ cm}^{-1}$ (COO^- stretching) were used as reference to predict the pectin loss or degree of methyl esterification. DME (degree of methyl esterification) was calculated based on Manrique and Lajolo, (2002), using the equation $\text{DME} = 124.7 * R + 2.2013$, where $R = A_{1740} / (A_{1740} + A_{1600-1630})$, $\%A = 2 - \log(\%T)$.

3. RESULTS AND DISCUSSION

3.1. Ultra-structural peel changes in 1-MCP treated and control (untreated) papaya fruit using SEM

SEM images (at 600X and 1200X) revealed notable difference between the 1-MCP treated and control papaya fruit peel (adaxial and abaxial view) modifications (**Fig 1, 2, 3**). At day seven of analysis, remarkable variations were observed among the fruit stored at ambient (AS) and cold conditions (CS). 1-MCP (900 ppb) treated fruit at AS appeared intact with homogenous cell matrix with well-defined/identifiable layers and less degraded than the control fruit. The fruit samples treated with 1-MCP, stored under CS clearly depicted clearly all the cell wall matrix with less intercellular spaces between the pectin matrixes. This clearly shows (from the magnified images) that 1-MCP treated fruit at CS maintained greater peel integrity, while the control fruit stored at cold storage showed higher depolymerization with many intercellular spaces. The control fruit peel (Abaxial view) at AS displayed shrunken, incomplete and

degraded structures (due to highly esterified pectin). These disintegrated, loosened cell wall structures were also visible under untreated fruit kept under CS, but it could be clearly identified to more intact than the ambient stored fruit. Hence, the extent of cell wall modification seemed to be more intact and coherent due to 1-MCP treatment which suppressed the ethylene action, than the storage conditions. The untreated fruit stored at ambient condition, depicted cracked surface with visible fractures, separations of the wall materials due to rapid ripening associated changes contributing to softening, and loss in firmness. The results in the study were similar to the findings of Sanudo-Barajas *et al.* (2009), where depolymerization of pectin polymers in papaya fruit, was due to ethylene associated ripening and softening changes, and 1-MCP application significantly improved the fruit firmness retention. Solubilization and depolymerization of cell wall polysaccharides (pectins; hemicelluloses) causes further dissolution of the cell wall materials, resulting in fruit firmness (Brummell and Harpster, 2001).

SEM images of adaxial view of the peel (**Fig 3.**) depicted that under AS (which were at post-climacteric phase), the untreated fruit peel had thinner layers, with deeper cracks, rough surface, loosened matrix, striated and shrunken compared to the 1-MCP treated fruits. This breakdown of middle lamella and loss of structural unity of cell walls could be presumed to be due to enhanced activity of pectinases enzymes like PME and PGase. Similarly, the untreated fruit under CS had disintegrated and coarse surface with cracks. 1-MCP treated fruit under CS had a smooth homogenous surface with no cracks. At day seven, treated fruit at AS had a coarse skin structure and lesser wall degradation than untreated fruit indicating the role of 1-MCP in decreasing the activity of cell wall hydrolyzing enzymes. The cross-sectional view (**Fig 4.**) of the peel showed that the 1-MCP treated fruit at CS was intact with lesser tissue degradation, indicating a decrease in the degree of polymerization and reduction in the solubilization of the hemicellulosic and pectic cell wall polysaccharides. These structural differences would explain the higher firmness shown by the fruit treated with 1-MCP (recorded in the present investigation, unpublished). Gonçalves and Vitória (2011) had also described the changes in papaya ultrastructural changes due to ripening and physiological disorders. Also, EDAX analysis (**Fig 5a,b.**) shows that 1-MCP treated fruit retained higher levels of potassium (K), calcium (Ca) and magnesium (Mg) when compared to control fruit. Studies on the effect of 1-MCP on K, Ca and Mg contents have not yet been reported but effect of 1-MCP in maintaining higher nutrient levels than the untreated fruits have been reported in few studies including mango (Islas-Osuna *et al.*, 2010) and tomato (Wang *et al.*, 2009 and Pangaribuan and Irving, 2010). Hence, the present SEM examination showed that papaya fruit that received treatment with 1-MCP remained firmer, had displayed higher peel integrity than the untreated fruit.

Fig 1. SEM images of control (a- 600X; b- 1200X) and 1-MCP treated (c- 600X; d-1200X) papaya fruit peel (abaxial view) under CS




Fig 2. SEM images of control (e- 600X; f- 1200X) and 1-MCP treated (g- 600X; h- 1200X) papaya fruit peel (abaxial view) under AS












Fig 3. SEM images of control (i, k at 600X) and 1-MCP treated (j, l at 600X) fruit peel (adaxial view) under AS (i, j) and CS (k, l)




UNDE

Fig4. SEM images of control (m,o at 600X) and 1-MCP treated (n, p at 600X) fruit peel (cross sectional view) under AS (m,n) and CS (o, p)


m.




n.



o.



p.



UNDETERMINED

Fig 5a. EDAX analysis of Mg, Ca and K (%) 1-MCP treated and control sample under ambient storage







Figure 5b. EDAX analysis of Mg, Ca and K (%) 1-MCP treated and control sample under cold storage



3.2. Analysis of pectin FT-IR spectra and Degree of methyl esterification (DME) in 1-MCP treated and untreated papaya fruits

The FT-IR method was used to analyze the methylation level of soluble pectin fractions isolated from the 1-MCP treated and untreated fruit peel as well as pulp, at day seven, when a significant change (climacteric peak) in the ambient storage condition was observed. The spectra of pectin standard showed a distinct peak associated with methylated esterified pectin (Fig 6.) at wavenumber 1730-1745 cm^{-1} (due to stretching of ester carboxyl group (-COOCH₃)) (Fig 6). With the hydrolysis of pectin due to ripening, we could observe two peaks; a peak at wavenumber (cm^{-1}) 1630-1645 = corresponding to asymmetric-COOH stretch and a peak at wavenumber (cm^{-1}) 1420-1425 = corresponding to symmetric -COOH stretch among the untreated fruit (Fig 9. & 10.), however the absence of specific methylated peak confirms the activity of cell wall degrading enzymes. In the untreated papaya peel and pulp spectra, the carbohydrate peaks are visible, which have been reported to have a spectrum between 1200-950 cm^{-1} wavenumber, that is highly specific for each polysaccharide. In 1-MCP treated papaya fruit pulp and peel had displayed peaks at wavenumber 1736 and 1742 cm^{-1} respectively, similar to that of the standard indicating suppressed or lesser pectin hydrolysis (Fig 7. and 8.) DME (degree of methyl-esterification) values were calculated from absorbance spectra of the samples, plotted on a graph. The spectral analysis showed that DME percentage was lower in the pulp and peel of control fruit stored at ambient (AS) and cold storage (CS) conditions, signifying rapid softening process. Higher the DME values, lesser is the rate of cell wall hydrolysis (indicates reduced softening). DME was calculated using the equation $\text{DME} = 124.7 * R + 2.2013$, where $R = A_{1740} / (A_{1740} + A_{1600-1630})$, $\%A = 2 - \log(\%T)$. The degree of methyl esterification (DME) of pectin changes during ripening process, more predominantly in climacteric fruits. Compositionally, pectin is poly α -1-4-galacturonic acids, with varying degree of carboxylic acid residues present as methylated esters (Fischer and Bennet, 1991; Voragen and Pilnik, 1995). Loss of fruit firmness is therefore attributed to an increase in pectin solubility and depolymerization, changes in PG activity during ripening process (Lazan *et al.*, 1995; Zhao and Paull, 1996). Fabi *et al.* (2009) described that upregulation of endo-PGases was directly proportional to the firmness loss in fruit, signifying the crucial role of 1-MCP in downregulation of these endo-PGases and firmness retention in papaya fruit. Application of 1-MCP at 900 ppb significantly delayed the ripening process by suppressing softening, which was evident from the above conducted experiments.




Fig 6. FT-IR spectrum of methylated pure pectin- standard (Sigma)




Fig 7. Pulp of 1-MCP treated fruit showing the characteristic peak similar to standard at around wavenumber 1736 cm^{-1}




Fig 8. Peel of 1-MCP treated fruit showing characteristic peak at wavenumber 1742 cm⁻¹




Fig 9. Pulp of untreated fruit depicting hydrolyzed pectin and more of carbohydrate peaks at around 1200-950 cm⁻¹




Fig 10. Peel of untreated fruit depicting hydrolyzed pectin spectra

Estimating the DME is thus an important indicator which reveals the significant softening and associated changes during fruit ripening, especially in a climacteric fruit like papaya. The lesser DME (%) in 1-MCP treated papaya fruits observed in this study was similar to the findings reported by Thumdee *et al.* (2010). The results from this corroborate with the GCMS profiles conducted during the study (Srividhya and Jeyakumar, 2017) where higher proportion of ester compounds were identified in control compared to 1-MCP treated. Therefore, this investigation further confirms that modifications in the carbohydrate matrix of cell wall, due to the activities of various hydrolytic enzymes are responsible for the tissue softening during fruit ripening (Paull and Chen, 1983; Lazan *et al.*, 2004; Ali *et al.*, 1998; Chen and Paull, 2003; Shiga *et al.*, 2009).

4. CONCLUSION

Application of 1-MCP had proved potential in improving the keeping quality of papaya fruit as revealed from the ultra-structural changes and FT-IR analysis. Studies to examine the papaya ripening associated characteristics using SEM and FT-IR spectroscopic were found to be very effective in studying the cell wall changes. The study further confirms that ethylene induced ripening and associated physiological, metabolic changes can be delayed using 1-Methylcyclopropene.

DISCLAIMER

The 1-MCP product used for this research is and was approved by Environment and Protection Agency, since its use and is predominantly used product in our area of research and country to prevent postharvest losses. There is absolutely no conflict of interest between the authors and producers of the product because we do not intend to use these products as an avenue for any litigation but for the

advancement of knowledge. Also, the research was not funded by the Agrofresh Inc., Mumbai company rather it was funded by personal efforts of the authors.

REFERENCES

1. Ali ZM, Chin LH, Lazan H. A comparative study on wall degrading enzymes, pectin modifications and softening during ripening of selected tropical fruits. *Plant Sci.* 2004; 67:317-327.
2. Ali ZM, Ng S, Othman R, Goh L, Lazan H. Isolation, characterization and significance of papaya β -galactanases to cell wall modification and fruit softening during ripening'. *Physiol. Plant.* 1998. 104: 105-115.
3. Barros AS, Mafra I, Ferreira D, Cardoso S, Reis A, da Silva JAL. et al. Determination of the degree of methylesterification of pectic polysaccharides by FT-IR using an outer product PLS1 regression. *Carbohydrate Polymers.* 2002; 50: 85-94.
4. Brummell DA, Harpster MH. Cell wall metabolism in fruit softening and quality and its manipulation in transgenic plants. *Plant Mol Biol.* 2001; 47: 311-340.
5. Chatjigakis AK, Pappas C, Proxenia N, Kalantzi O, Rodis P, Polissiou M. FT-IR spectroscopic determination of the degree of esterification of cell wall pectins from stored peaches and correlation to textural changes. *Carbohydrate Polymers.* 1998; 37, 395-408.
6. Chen NJ, Paull RE. Endoxylanase expressed during papaya fruit ripening: purification, cloning and characterization, *Funct. Plant Biol.* 2003. 30: 433-441.
7. Dick AJ, Labavitch JM. Cell Wall Metabolism in Ripening Fruit. *Plant Physiol.* 1989; 89: 1394-1400.
8. Fabi JP, Cordenunsi BR, Seymour GB, Lajolo FM, Oliveira do Nascimento JR. Molecular cloning and characterization of a ripening-induced polygalacturonase related to papaya fruit softening. *Plant Physiol. Biochem.* 2009; 47: 1075-1081.
9. Islas-Osuna MA, Norma A, Stephens-Camacho CA, Contreras-Vergara M, Rivera-Dominguez E, Sanchez-Sanchez MA, Villegas-Ochoa A, Gonzalez-Aguilar GA. Novel postharvest treatment reduces ascorbic acid losses in mango (*Mangifera indica* L.) var. Kent. *Am. J. Agric. Biol. Sci.* 2010; 5: 342-349.
10. Jeong J, Huber DJ, Sargent SA. Influence of 1-methylcyclopropene (1-MCP) on ripening and cell-wall matrix polysaccharides of avocado (*Persea americana*) fruit, *Postharvest Biol. Technol.* 2002; 25: 241-256.
11. Koubala BB, Christiaens S, Kansci G, Loey AMV, Hendrickx ME. Isolation and structural characterization of papaya peel pectin. *Food Res. Int.* 2014; 55: 215-221.
12. Lazan H, Ng S, Goh L, Ali ZM. Papaya β -galactosidase/galactanase isoforms in differential cell wall hydrolysis and fruit softening during ripening. *Plant Physiol. Biochem.* 2004; 42: 847-853.
13. Lazan H, Selamat MK, Ali ZM. β -Galactosidase, polygalacturonase and pectin esterase in differential softening and cell wall modification during papaya fruit ripening, *Physiol. Plant.*, 1995; 95: 106-112.

14. Manenoi A, Bayogan ERV, Thumdee S, Paull RE. Utility of 1-methylcyclopropene as a papaya postharvest treatment. *Postharvest Biol Technol.* 2007; 44:55-62.
15. Manrique GD, Lajolo FM. FT-IR spectroscopy as a tool for measuring degree of methyl esterification in pectins isolated from ripening papaya fruit. *Postharvest Biol. Technol.*, 2002; 25: 99-107.
16. Pangaribuan DH, Irving D. The effect of 1-mcp in maintaining the quality of tomato slices. *J. Teknol. dan Industri. Pangan.* 2010; 21: 1-7.
17. Paull RE, Chen NJ. Postharvest variation in cell wall-degrading enzymes of papaya (*Carica papaya* L.) during fruit ripening. *Plant Physiol.* 1983; 72: 382-385.
18. Paull RE, Gross K, Qiu YX. Changes in papaya cell walls during fruit ripening. *Postharvest Biol. Technol.* 1999; 16: 79-89.
19. Fischer RL, Bennett AB. Role of cell wall hydrolases in fruit ripening, *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 1991; 42: 675-703.
20. Sanudo-Barajas JA, Labavitch J, Greve C, Osuna-enciso T, Muy-rangel D, Siller-Cepeda J. Cell wall disassembly during papaya softening: Role of ethylene in changes in composition, pectin-derived oligomers (PDOs) production and wall hydrolases. *Postharvest Biol. Technol.* 2009; 51: 158-167.
21. Shiga TM, Fabi JP, do Nascimento JRO, Petkowicz CLD, Vriesmann LC. Changes in cell wall composition associated to the softening of ripening papaya: evidence of extensive solubilization of large molecular mass galactouronides, *J. Agric. Food Chem.* 2009; 57: 7064-7071.
22. Srividhya S, Jeyakumar P. Effect of 1-methylcyclopropene (1-MCP) on volatile compound production in papaya (*Carica papaya* L.) fruit. *The Pharma Innovation Journal.* 2017; 6(11): 532-536.
23. Szymanska-Chargot M, Zdunek A. Use of FT-IR Spectra and PCA to the Bulk Characterization of Cell Wall Residues of Fruits and Vegetables Along a Fraction Process. *Food Biophysics.* 2013; 8: 29-42.
24. Thumdee S, Manenoi A, Nancy JC, Paull RE. Papaya Fruit Softening: Role of Hydrolases. *Trop. Plant Biol.* 2010; 3: 98-109.
25. Voragen AGJ, Pilnik W. Pectins. In: Stephen AM (Ed.), *Food polysaccharides and their applications*. New York: Marcel Dekker Inc. 1995.; pp: 287-339.
26. Wang CY, Chen CT, Wang SY. Changes of flavonoid content and antioxidant capacity in blueberries after illumination with UV-C. *Food Chem.* 2009.; 117: 426-431.
27. Zhao M, Moy J, Paull RE. Effect of gamma-irradiation on ripening papaya pectin. *Postharvest Biol. Technol.* 1996; 8: 209-222.