

# Studies on antimicrobial effect of botanicals on leaf blight of Indian mustard caused by *Alternaria brassicae*

## ABSTRACT

The present study was conducted at ANDU&T Kumarganj Ayodhya at student instructional form (SIF). As in the investigation it was observed that foliar spray with Carbendazim (0.1%), Neem oil (5%), followed by Neem leaf extract (5%) found to be more effective on growth parameters like Number of branches/plants, number of leaves/ plants, plant height (cm), number of pods/plants,) and increased the yield (1000-seed weight, per plant yield (g), per hectare yield) and also significantly reduced the *Alternaria* blight in Indian mustard. Carbendazim (0.1%), *Trichoderma harzianum* (2%) followed by garlic bulb extract (5%) was found most effective in reducing *Alternaria* blight, recorded maximum disease incidence i.e. percent Disease intensity, percent Leaf area disease (LAD), percent leaf infection, number of spots per pods, seed infection percent respectively.

**Keywords:** *Alternaria brassicae*, Leaf blight, Indian mustard and Botanicals.

## 1. INTRODUCTION

India is one of the major producers in the global oilseeds/ vegetable oil economy. India has the largest area in oilseed crops in the world. However, the productivity is low as it is just 50-60 percent of the world average [1]. *Alternaria* blight is a most dangerous disease of rapeseed mustard crop and this appears all winter session. Crops throughout the crop growing areas of the world. It occurs in various countries like Canada, England, France, Holland, Poland, Spain, Sweden, Australia, USA, and Asian countries. *A. brassicae* causal organism of *Alternaria* blight disease of mustard was reported from whole world. It was infected cruciferous crops and its causing yield loses with no proven main diseases of mustard source of movable fight in several of the crops [2]. This disease is a major limiting factor in reducing the yield of Rapeseed-mustard in India. Which is range from 15-71% due to this disease. The disease cause quantitative loss as well as, seed quality in term of seed size seed colour and oil content are reduced due to loss of chlorophyll in infected plant which also reduce the growth and development of plant [3]. *A. brassicae* is a pathogen necrotrophic and can cause serious loss to the crop. The major symptoms are appears as the form of spots on leaves, stem and pods mustard plant [4,5,6].

## 2. MATERIAL AND METHODS

### 2.1 Collection of botanicals and preparation of plant extract

Botanicals such as leaf of Neem leaf (*Azadirachita indica*), and *Ocimum sanctum* collected from ANDU&T field and Onion (*Allium cepa*), Ginger, Neem oil, and clove of Garlic (*Allium sativum*), *Trichoderma harzianum* and Carbendazim were collected from local market. For preparation of extract, collected plant materials were weighed in an electric balance and then washed in water. Plant material are chopped in required quantity of water (1:1. w/v), boiled for 30 min. and then strain through cheese cloth to obtained standard plant extract solution (100%). The extract is centrifuge at 5000 rpm for 5 min. and the clear supernatant was used as stock solution. All the treatments were used as foliar spray as given in table no.1.

40 **Table no.1. Details of different treatments. (Used as foliar spray).**

Treatments	Name	Dose
T <sup>1</sup>	<i>Trichoderma harzianum</i>	2%
T <sup>2</sup>	Neem oil	5%
T <sup>3</sup>	<i>Alium cepa</i>	5%
T <sup>4</sup>	Ginger rhizome extract	5%
T <sup>5</sup>	Garlic bulb extract	5%
T <sup>6</sup>	<i>Ocimum sanctum</i> extract	5%
T <sup>7</sup>	Neem leaf extract	5%
T <sup>8</sup>	Carbendazim	0.05%
T <sup>9</sup>	Control	Water Spray

41 **2.2 Application of spray solution**

42 The Botanicals and fungicidal solutions were prepared by mixing with required amount of Botanicals  
 43 and fungicide with water. seven botanicals and one fungicidal solution, (concentration @ 1:10) and  
 44 plain water were sprayed with compressed hand sprayer as described in table no.1. Sprays were  
 45 done at 30, 40, 50, 60 days after sowing. Every time the botanicals and fungicide were freshly  
 46 prepared prior to application and the spray tank was thoroughly cleaned before filling with new spray  
 47 solution. Special attention was given to complete coverage of the growing plants with the fungicides.  
 48 Adequate precaution was taken to avoid drifting of spray materials from one plot to the neighbouring  
 49 ones.

51 **3. RESULTS AND DISCUSSION**

52 **3.1 Plant growth Parameters**

53 The effect of treatments on the germination was recorded after the maximum number of  
 54 seeds were germinated. As displayed in table 2 the maximum germination was recorded in T7  
 55 (95.75%) and T5 (95.12%) over the control T9 (92.94%) respectively followed by T3 (95.04%), T4  
 56 (95.0%), T6 (94.8%), T8 (93.88%), T2 (93.63%), T1 (93.60) as all were significantly effective among  
 57 the botanical studies that *Azadirachita indica* is more effective against *Alternaria brassicae* of Indian  
 58 mustard [7]. Many researchers used Neem leaf, Onion extract found more effective in germination [3].  
 59 The data indicates that all treatments were significantly showing the effect as they were increasing the  
 60 number of leaf/plant. The table no.2 is the revealing that T8 (25.00) and T2 (21.13)] were showing the  
 61 best effect of treatment over the control i.e.T9 (14.93) respectively followed by T7 (20.13) T1 (18.93),  
 62 T6 (18.20), T3, T4 (17.27), T5 (16.33).

63 All the treatments were significantly showing the effect as they were increasing the number of  
 64 branches/plant. The table no.2 is the showing that T8 (7.17) and T2 (7.03) were effective over the  
 65 control i.e.T9 (5.17) respectively followed by T7 (6.77), T1 (6.57), T6 (6.37), T3 (6.30), T4 (6.03), T5  
 66 (5.77). Researchers [8] used Neem leaf extract and garlic bulb extract that were significantly effective  
 67 and reduced disease severity and increased yield. The data indicate that all the treatments were  
 68 significantly best over the T9 as they were increasing the plant height. It may be seen from the data  
 69 (table no. 2) that best result found in T8 (79.53 cm), T2 (77.87 cm) and T7 (77.60 cm) followed by T3  
 70 (71.67 cm), T4 (68.27 cm), T5 (65.60 cm). [9] also used several fungicides and botanicals to manage  
 71 the alternaria blight which also were increasing the plant growth.

72 All treatments were significantly showing the effect as they were increasing the number of  
 73 pods/plant. Table No. 2 is displaying that T8 (247.20), T2 (216.27) and T7 (205.93) were showing the  
 74 best effect of treatment over the control i.e.T9 (156.33) followed by T3 (177.07), T4 (163.20), T5  
 75 (159.20). Maximum height found plant from T8 (Carbendazim) plot among the other all treatments.  
 76 Among the other all treatments best result found from T2 (Neem oil) followed by all botanicals and  
 77 bio-agent. Similar observation was recorded by [7] and [10] as he used botanicals' including Garlic  
 78 bulb extract which were effective among the botanicals. 1000-seed weight was found to be significant  
 79 in all the treatments as there were having positive effect. Spraying with T8 (4.50 g) T2 (4.33 g) and T7  
 80 (4.17g) were showing the best effect of treatment over the control i.e., T9 (2.77g) followed by T3 (3.80

81 g), T4 (3.67 g), T5 (3.17 g) respectively the data indicate by (table no.2). Many researchers studied  
82 the same aspect and the study revealed similarity of present investigation [11].


83 Significant variation of different treatments was found on yield per plant (g) and yield quintal  
84 per hectare. Maximum yield per plant (5.60 g) and per hectare (8.97 q) was obtained from T8  
85 (Carbendazim) treated plot followed by T2 (neem oil), and T7 (Neem leaf extract) in both the cases.  
86 The minimum yield per plant (3.07 g) and per hectare (4.70 q) was recorded from T9 (Control) (Table-  
87 1). Carbendazim T8 (Carbendazim) was more effective than other treatment respectively and T2  
88 (Neem oil) was more effective among the botanicals. [8, 12] use Neem leaf extract and garlic bulb  
89 extract were significantly and reduce disease severity and increase yield.

### 90 3.2 Disease incidence Parameters

91 All the treatments were significantly showing the effect as they were increasing the disease  
92 intensity **percent** (%). The table no.3 is the showing that T8 (41.93%) T1 (41.93 %) and T5 (42.40 %)   
93 were showing the best effect of treatment over the control i.e.T9 (48.07) respectively followed by T2  
94 (43.00 %), T6 (43.00 %), T7 (43.53 %). [13, 14] They also used Garlic bulb extract *T. harzianum* and  
95 Neem leaf extract botanicals which were significantly effective in reducing disease incidence. Also [7,  
96 15, 16] revealed that *Trichoderma harzianum* which was best with the botanicals. [6, 17] used garlic  
97 bulb extract, Tulsi and Neem extract at concentration 5%, 10% is more effective on disease intensity.  
98 [18, 19, 20] supported the present investigation by their study on Carbendazim and *Trichoderma*  
99 *harzianum* reducing the disease severity and increasing the grain yield. [8, 12] use Neem leaf extract  
100 and garlic bulb extract which were significantly effective which reduced disease severity and  
101 increased yield.

102 The data indicate that all the treatments were significantly best over the T9 (control) (58.83). It  
103 may be seen from the data (table no.3) that best result found in T8 (32.07), T1 (38.47), T5 (39.33)  
104 followed by T2 (41.67), T6 (41.67), T7 (42.77). [21, 22] also confirm reduced disease intensity the  
105 present investigations finding were also supported by [23, 24, 25]. The **Percent** leaf area disease  
106 (LAD) in data it is indicated that all the treatments were significantly best over the T9 (21.33 %).  
107 It may be seen from the data (table no.3) that best result found in T8 (5.67 %), T1 (8.00 %), T5 (10.0 %)   
108 over the control [T9 (21.33 %)] respectively. followed by T2 (13.00 %), T6 (13.67 %), T7 (14.33 %).  
109 **Percent** pod infection. A perusal of the data indicates that all the treatments were significantly superior  
110 over the T9 [control (41.83 %)] in **Percent** pod infection. It may be seen from the data (table no.3) that  
111 minimum percent pod infection was found in T8 (25.17 %), T1 (25.50 %), T5 (25.87 %) followed by T2  
112 (26.50 %), T6 (26.80 %), T7 (31.17 %). Similar observation was reported by [1] he was also observed  
113 effect of botanicals in reduce disease severity. [14, 19] They were also used Garlic bulb extract *T.*  
114 *harzianum* and Neem leaf extract botanicals which significant effective reduce disease incidence [7].  
115 A perusal of the data indicates that all the treatments were significantly superior over the T9 [control  
116 (2.17)] Number of spot per pods. It may be seen from the data (table no.3) that minimum **percent** pod  
117 infection was found in T8 (0.97), T1 (1.17), T5 (1.17) followed by T2 (1.67), T6 (1.60), T7 (1.67). They  
118 were used botanicals which significant effective reduce disease incidence.

119 After harvest **percent** seed infection was examined. **Percent** seed infection by *Alternaria*  
120 *brassiccae* and *Alternaria brassicicola* of harvested seed received from treated plot with different  
121 botanicals, bio-agent and fungicide was found to be significant. Seed obtained from control treatment  
122 showed the highest **percent** seed infection (19.17%) while seed obtained from carbendazim (T8)  
123 treated plot showed the lowest seed infection (13.37%) preceded by T1 (13.67%), T5 (14 %)   
124 respectively and followed by T2 (14.83)], T6 (14.53)], T7 (16.07). T8 [Carbendazim] was found more  
125 effective for controlling seed borne *Alternaria brassiccae* of harvested mustard seeds. Similar result  
126 was recorded by [19] he used nine botanicals' Garlic bulb extract and found more effective as similar  
127 as [6, 14, 23]. Many researchers also used Garlic bulb extract which was significantly effective in  
128 reducing disease incidence. *T. harzianum* and Neem leaf extract as botanicals [24, 25].



129  
130  
131

**Plate no. 1.** Symptoms of Alternaria blight on mustard leaves (A and D. blighted spots on leaf, B. Symptoms on pods, C. concentric rings).

UNDER PEER REVIEW

132  
133

**Table no.2. Effect of different treatments of growth parameters of crop.**

Treatments	Germination % Mean±SD	Leaves Per Plants Mean±SD	Branches Per Plants Mean±SD	Plant height (cm) Mean±SD	Pods Per Plants Mean±SD	1000 seed weight Mean±SD	Yield Plant Mean±SD	PerYield In(g)hectare Mean±SD	Per in (q) Mean±SD
T <sup>1</sup> <i>Trichoderma harzianum</i>	93.60±0.96	18.93±1.33	6.57±0.12	75.73±1.70	198.47±11.23	4.17±0.76	4.33±0.76	6.83±1.26	
T <sup>2</sup> Neem oil	93.63±2.69	21.13±0.70	7.03±0.42	77.87±1.33	216.27±18.83	4.33±1.53	4.83±1.26	8.17±1.04	
T <sup>3</sup> <i>Alium cepa</i>	95.04±1.80	17.73±2.10	6.30±0.20	71.67±1.53	177.07±3.31	3.80±0.82	4.13±0.81	6.64±1.30	
T <sup>4</sup> Ginger rhizome extract	95.00±0.33	17.27±2.00	6.03±0.50	68.27±1.03	163.20±8.32	3.67±1.53	3.80±1.06	5.50±0.87	
T <sup>5</sup> Garlic bulb extract	95.12±4.00	16.33±2.08	5.77±0.61	65.60±1.74	159.20±10.25	3.17±0.76	3.50±0.50	5.10±1.15	
T <sup>6</sup> <i>Ocimum sanctum</i> extract	94.08±1.46	18.20±2.23	6.37±0.31	74.67±2.14	190.80±7.79	4.00±1.32	4.17±0.76	6.80±0.75	
T <sup>7</sup> Neem leaf extract	95.75±1.69	20.13±1.33	6.77±0.12	77.60±1.44	205.93±8.90	4.17±1.76	4.63±1.31	7.30±0.82	
T <sup>8</sup> Carbendazim	93.88±3.96	25.00±0.53	7.17±0.46	79.53±1.63	247.20±9.10	4.50±1.32	5.60±0.53	8.97±1.05	
T <sup>9</sup> Control	92.94±1.12	14.93±2.05	5.17±0.50	61.40±2.75	156.33±9.19	2.77±0.93	3.07±1.01	4.70±0.82	
SE(in)	1.28	0.47	0.12	1.05	6.15	0.56	0.54	0.60	
CD	3.71	1.37	0.35	3.05	17.88	1.64	1.56	1.75	

134  
135  
136

**Table no.3. Effect of different treatments of disease parameters of crop.**

Treatments	Percent disease intensity Mean±SD	Percent leaf infection Mean±SD	Percent leaf Area Disease (LAD) Mean±SD	Percent Pod infection Mean±SD	Number of spots per Mean±SD	Percent seed infection Mean±SD
T <sup>1</sup> <i>Trichoderma harzianum</i>	41.93±0.12	38.47±2.34	8.00±1.00	25.50±1.32	1.17±0.76	13.67±1.26
T <sup>2</sup> Neem oil	43.00±0.72	41.67±3.21	13.00±3.61	26.50±2.78	1.67±1.15	14.83±2.25
T <sup>3</sup> <i>Alium cepa</i>	42.73±2.77	40.33±3.79	11.67±1.53	26.17±1.04	1.27±0.64	14.33±1.53
T <sup>4</sup> Ginger rhizome extract	42.93±0.76	41.33±3.21	12.67±1.53	26.33±3.79	1.33±0.76	14.50±2.21
T <sup>5</sup> Garlic bulb extract	42.40±2.46	39.33±3.06	10.00±1.00	25.87±2.20	1.17±0.76	14.00±2.00
T <sup>6</sup> <i>Ocimum sacntum</i> extract	43.00±1.04	41.67±4.04	13.67±2.08	26.80±1.59	1.60±0.53	14.53±3.32
T <sup>7</sup> Neem leaf extract	43.53±1.29	42.77±2.54	14.33±1.53	31.17±1.04	1.67±0.58	16.07±2.50
T <sup>8</sup> Carbendazim	41.93±1.14	32.07±2.10	5.67±2.08	25.17±1.44	0.97±0.06	13.37±1.70
T <sup>9</sup> Control	48.07±0.81	58.83±1.76	21.33±1.53	41.83±1.26	2.17±0.76	19.17±1.04
SE (m)	0.81	1.51	1.14	1.21	0.37	1.14
CD	2.35	4.40	3.31	3.52	1.06	3.31

#### 137 4. CONCLUSION

138 As the chemicals have the hazardous effect on environment, mammals and also economical  
139 consideration. Researchers are trying to find eco-friendly bio compounds for plant diseases  
140 management which should be beneficial, cheaper and easily available too. To generate awareness of  
141 harmful effects of fungicides integrated use of bioagents *Trichoderma harzianum* and botanicals will  
142 be the perfect solution as foliar spray against Alternaria blight. In this investigation among the  
143 botanicals and bio agent Neem oil, Neem leaf extract, *Trichoderma harzianum* and Garlic bulb extract  
144 is effective for management of Alternaria blight of mustard caused by *Alternaria brassicae*. So with  
145 this investigation it can be said that many of the botanicals can be used as alternatives of the  
146 chemicals which can be so cheap and good for human and environment too.

#### 147 REFERENCES

- 148 1. Chadar KL, Singh PR, Singh KR, Yadav RR, Mishra KM, Pratap N and Vishnoi KR. Studies  
149 on Alternaria blight of rapeseed-mustard (*brassica juncea* L.) caused by *alternaria brassicae*  
150 (berk.) sacc. and its integrated management. *Plant Archives*. 2016;16 (2): 897-901.
- 151 2. Meena PD, Awasthi RP, Chattopadhyay C, Kolte SJ. and Kumar A. Alternaria blight: a chronic  
152 disease in rapeseed-mustard. *Journal of Oilseed and Brassica*, 2010 (1): 1-11.
- 153 3. Singh KC, Singh PS, Tiwari KN, Tiwari T. and Singh D. bio-efficiency of plant extract,  
154 mancozeb and *Trichoderma viride* to control the Alternaria leaf spot of cabbage *Annals of*  
155 *Plant. and So. Res.* 2015;17 (Special Issue): 219-221.
- 156 4. Prasad R, Khandelwal GL and Jain JP. Morphology, physiology and control of Alternaria  
157 brassicae on taramira. *Indian Phytopathology*, 1970. 23: 105-110.
- 158 5. Chattopadhyay C. Management of diseases of rape-seed mustard with special reference  
159 Indian conditions. In: Sustainable production of oil seeds: rapeseed-mustard technology  
160 (Eds.:A. Kumar, J.S. Chauhan and C. Chattopadhyay). Agrotech Publ. Acad., Udaipur, 2008.  
161 364-388.
- 162 6. Kumar M, Zachariya S. and Lal AA. Management of alternarial blight of mustard (*brassica*  
163 *juncea* L.) By botanicals, *Trichoderma harzianum* and fungicides *Pant Archive*, 2019. 19 (1),  
164 1108-1113.
- 165 7. Sharma P, Singh SD, Rawal P and Sharma P. Antifungal activity of some plant extracts and  
166 oil against seed borne pathogens of pea. *Plant Disease Research. Ludhiana* 2003. 18(1):16-  
167 20.
- 168 8. Ferdous SM, Meah MB and Hossain MM. Comparative effect of plant extracts and fungicide  
169 on the incidence of Alternaria blight of mustard. *Bangladesh Journal Training Dev.* 2002.  
170 15(1-2): 207-210.
- 171 9. Yadav CL, Kumar N and Kumar R. Effect of Seed Treatments with Fungicides, Bio-agents  
172 and Botanicals against Alternaria Leaf Spot in Cabbage (*Brassica oleracea* var. capitata L.)  
173 *Trends in Bioscience*. 2014. 7(23): 3823-3827.
- 174 10. Mahapatra S and Das S. Evaluation of Fungicides and Botanicals Against Alternaria Leaf  
175 Blight of Mustard. *Indian Journal of Plant Protection*. 2013. 41(1) 61-65.
- 176 11. Thakur Y, Zacharia S and Chauhan BS. Efficacy of bio-agents and plant extracts against  
177 Alternaria leaf blight of mustard (*Brassica juncea* L.); *European Journal of Biotechnology and*  
178 *Bioscience*. 2017. 5 (4):29-35
- 179 12. Ansari, La P and Tiwari S. Organic alternative and management leaf blight of mustard.  
180 *Journal of pharmacognosy and Phytochemistry*. 2017. 6 (6) 1463-1465.
- 181 13. Meena PD, Meena RL, Chattopadhyay C and Kumar A. Identification of critical for disease  
182 development and bio control of Alternaria blight of mustard (*Brassica juncea*) *journal of*  
183 *phytopathology*. 2004. 152;204-209.
- 184 14. Thakur Y and Zacharia S. Efficacy of bio-agents and plant extracts against Alternarial leaf  
185 blight of mustard (*Brassica juncea* L.) *International Journal of Chemical Studies*. 2018. 6 (1):  
186 1187-1189.
- 187 15. Meena PD, Jambhulkar SJ, R. Gupta R, Meena HS and Singh D. Rapid screening technique  
188 for Alternaria blight resistance in Indian mustard (*Brassica juncea*) cotyledonary leaf method;  
189 *Journal of plant Pathology*, 2016. 98(3), 463-469.
- 190

- 191 16. Jackson SK and Kumar A. Management of Alternaria leaf spot of Mustard through chemical  
192 and biological agents Article *Plant Cell Biotechnology and Molecular Biology*. 2019. 20  
193 (3&4):162-178.
- 194 17. Yadav JK, Singh HK, Singh SK, Kavita & Singh S. Efficacy of plant extracts against *Alternaria*  
195 *brassicae* under *in- vitro* condition. *Journal of Pharmacognosy. and Phytochemistry*. 2019. 8  
196 (1): 528-532.
- 197 18. Jayapradha C, and Raja Yesu. A Review of eco-friendly Management of *Alternaria* Species  
198 *Indian journal of hill farming*. 2016. 29 (2)106-119.
- 199 19. Singh K, Lal AA, Kumar D, and Meena KN. Evaluation of Selected Bio Agents, Plant Extracts  
200 and Fungicides for the Management of *Alternaria* Leaf Blight of Indian Mustard. *Inter. Journal*  
201 *of current Microbiology and Applied Science*. 2017. 6 (4)26-31.
- 202 20. Rai D, Mishra AK and Pandey A. Integrated Management of *Alternaria* Blight In Indian  
203 Mustard. *The Bioscan*. 2014. 9(3): 1247-1249.
- 204 21. Bhandekar P D, More VS and Aware RG. Efficiency of Biocontrol Agents, Botanicals and  
205 Chemical against *Alternaria porri*. *International Journal of Current Microbiology and Applied*  
206 *Science*. 2019. 8 (2) 384-391.
- 207 22. Waghe KP, Wagh SS, Kuldhar DP and Pawar DV. Evaluation of different fungicides,  
208 bioagents and botanicals against *Alternaria* blight caused by *Alternaria helianthi* (Hansf) of  
209 sunflower. *Academy Journal*. 2014. 10(5):351-358.
- 210 23. Chand H. and Singh S. Effect of plant extracts on *Alternaria* blight of mustard *Alternaria*  
211 *brassicae* (Berk.) Sacc. *Indian Journal of Plant Protection*. 2004. 32(2): 143-144.
- 212 24. Prasad, R. Management of *Alternaria* blight of mustard with combination of chemicals and  
213 botanicals. *Annals Plant Protection Science*. 2006. 14(2): 400-403.
- 214 25. Singh RB and Singh, RN. Spray schedule for the management of *Alternaria* blight and white  
215 rust of Indian mustard (*Brassica juncea*) under different dates of sowing. *Indian Journal of*  
216 *Agriculture Science*. 2006. 76 (9): 575-579.