

ASSESSMENT OF GENOTYPES X ENVIRONMENT INTERACTION OF BLACK GRAM USING MULTIVARIATE ANALYSIS

Comment [EN11]:

ABSTRACT

Twenty black gram genotypes were evaluated at five different locations of middle Gujarat in *kharif* 2016 to assess the genotype x environmental interactions in RBD design with two replications. The data were analyzed according to the AMMI model and AMMI based stability measures (ASV $W_{i(AMMI)}$ and $ASTAB_i$). Analysis of variance on the data pooled over locations and G x E interaction was found significant indicated genotypes performed differently in different locations. IPCA1 and IPCA2 were found significant in AMMI model and both combined accounted for by 78.7% variance of GEI. Environments viz., Devgadbaria, Derol and Dahod were found high yielding environments whereas Vadodara and Jabugam were low yielding environments. Genotypes G3, G18, G16 and G10 gave high yield in environment E3, E1 & E4, E5 and E2, respectively as they were vertex genotypes in polygon of AMMI2. According to AMMI model, ASV, $W_{i(AMMI)}$ and $ASTAB_i$, G19 was found stable and high yielding genotype, whereas G16 was unstable genotypes.

Key-words: Black gram, AMMI, GEI, ASV, $W_{i(AMMI)}$ and $ASTAB_i$

INTRODUCTION

Black gram or urd (*Vignamungo* (L.)_Hepper) is a tropical legume grown widely in India, China, Japan and Brazil mainly for its dry seeds, daal which is an important source of easily digestible protein which supplements the staple rice diet. It is said to be poor man's meat and rich man's vegetable. In Gujarat; area, production and productivity of black gram is 64 thousand hectares, 38 thousand tonnes and 594 kg per hectare, respectively (www.indiastat.com). The nation to become pulse sufficient, productivity level of pulses has to be increased substantially upto 1200 kg per hectare by 2020, which is possible through the development of stable high yielding genotypes using elite parental genotypes. There are different stability methods for studying genotype × environment interaction, viz. parametric method and non-parametric method and multivariate method. Among parametric methods; analysis of variance (ANOVA) models like Eberhart and Russell model is mostly used for studying the G x E interaction, whereas many non-parametric methods (Huehn, 1979 & Nassar and Huehn, 1987), (Thennarasu, 1995), (Sabaghnia, 2015) are commonly used for G x E interaction. But AMMI

model is multivariate technique to interpret GEI using ANOVA and Principal component analysis. The analysis of variance is useful for identifying and testing sources of variability, it provides no insight into the particular pattern of the underlying interaction. The ANOVA model effectively describes the main additive effects, while interaction (residuals from the additive model) is non-additive and requires alternative techniques, such as principal component analysis (PCA) to identify interaction pattern. Thus, ANOVA and PCA models combined to constitute the Additive Main effects and Multiplicative Interaction (AMMI) model (Gauch and Zobel, 1997; Zobel *et al.*, 1988).

The present investigation was carried out to analyze the pattern of genotype x environment interaction (GEI) for yield, to select the stable genotypes of black gram and to compare the stability measures viz., $W_{i(AMMI)}$ (Raju, 2002), ASV (Purchase, 1997) and ASTAB_i (Rao *et al.*, 2004) were calculated based on AMMI analysis for stability of black gram genotypes..

MATERIALS AND METHODS

A set of twenty black gram genotypes [*Vignamungo* (L.) Hepper] was evaluated at five different locations viz. Vadodara (E1), Jabugam (E2), Devgadbaria (E3), Derol (E4) and Dahod (E5) in middle Gujarat, India during the *kharif*-2016 in randomized complete block design with two replications. The yield data of multi-locational trial (Zonal varietal trial) were subjected to stability analysis using multivariate method (AMMI).

Statistical model:

$$\text{ANOVA : } Y_{ij} = \mu + \alpha_g + \beta_e + \alpha\beta_{ge} + \rho_{ij} + \varepsilon_{ijk}$$

$$\text{PCA : } Y_{ij} = \mu + \sum_n \lambda_n \gamma_{gn} \delta_{en} + \rho_{ij} + \varepsilon_{ijk}$$

$$\text{AMMI : } Y_{ij} = \mu + \alpha_g + \beta_e + \sum_n \lambda_n \gamma_{gn} \delta_{en} + \rho_{ij} + \varepsilon_{ijk}$$

Where

μ = grand mean ,

α_g = deviations of genotype(g)

β_e = deviations of environment(e)

λ_n = singular value for Interaction Principal Component Axis n (IPCA)

γ_{gn} = genotype eigenvector for axis n

δ_{en} = environment eigenvector

ρ_{ij} = residual

ε_{ijk} = error term or uncontrolled variation

The bi-plot is a graphical representation from AMMI analysis which is a useful tool to understand more complex specific pattern of genotypes and GEI or both genotypes and

environments. The concept of bi-plot was first developed by Gabriel (1971). It is a scatter plot that graphically displays the genotype (entries) and the environments (testers) of a two-way data and allows visualization of the interrelation among the entries (genotypes) and interaction between entries and testers (environments).

W_{i(AMMI)}

W_{i(AMMI)} a measure of stability is as good as Wricks's covaleance (W_i²) was estimated as under (Raju, 2002).

$$W_{i(AMMI)} = \sum_{m=1}^M \lambda_m^2 \gamma_{mi}^2$$

λ_m^2 = singular value for Interaction Principal Component Axis m (IPCA)

γ_{mi}^2 = genotype eigenvector for axis n

AMMI Stability Value (ASV)

The AMMI model does not make provision for a quantitative stability measure, such a measure is essential in order to quantify and rank genotypes according their yield stability, the following measure proposed by Purchase (1997) was estimated as under.

$$ASV = \sqrt{\left(\frac{IPCA1 \ SS}{IPCA2 \ SS} \times (IPCA1 \ score)^2 \right) + (IPCA2 \ score)^2}$$

AMMI based Selection Indices

Rao *et al.* (2004) proposed a new stability measure and incorporated as a stability component. When more than two axis are retained in AMMI model, the biplot formulation of interaction is failed. When n' of N axis are retained in the AMMI model to explain GEI, then the stability measure of ith variety can be determined as the end point of its vector α_{1i}^* , α_{2i}^* , ..., α_{ni}^* from the origin 0_{n x 1}. This is a squared Euclidean distance and was calculated as under.

$$ASTABi = \alpha_{1i}^{2*} + \alpha_{2i}^{2*} + \dots + \alpha_{ni}^{2*} = \sum_{n=1}^{n'} \alpha_{ni}^{2*} = \sum_{n=1}^{n'} \lambda_n \alpha_{ni}^2$$

A genotype is considered as highly stable when the value of ASTABi is small or closer to zero.

RESULT AND DISCUSSION

The combined analysis of variance (ANOVA) of twenty black gram genotypes over five locations according to AMMI model is presented in Table 1.

Table1: Analysis of variance (ANOVA) according to AMMI model for black gram genotypes.

| Source of Variations | df | Sum of Squares | Mean Squares | F Ratio | % SS |
|----------------------|----|----------------|--------------|---------|-------|
| Trials | 99 | 8.242 | 0.083 | 2.38 | |
| Genotypes | 19 | 0.759 | 0.040 | 1.14 | 9.21 |
| Environments | 4 | 4.827 | 1.207** | 34.53 | 58.57 |
| GxE Interaction | 76 | 2.656 | 0.035** | | 32.23 |
| PCA I | 22 | 1.671 | 0.076** | 10.86 | 62.91 |
| PCA II | 20 | 0.420 | 0.021** | 3.00 | 15.81 |
| PCA III | 18 | 0.358 | 0.020 | 2.86 | 13.48 |
| Residual | 16 | 0.207 | 0.013 | 1.86 | 7.79 |
| Pooled residual | 95 | 1.295 | 0.007 | | |

** Significant at $P < 0.01$.

Additive effects for main effects (genotypes and environments) and multiplicative effects for G x E interaction are considered. The results presented in Table 1 indicated that mean square for genotypes was found non-significant but locations (environments) and GEI were found highly significant indicating the diverse performance of genotypes over locations. The proportion of variance due to locations was the largest (58.57 per cent) followed by the variance due to G x E interaction (32.23 per cent) and genotypes (9.21 per cent). ANOVA provided no insight into the particular pattern of genotypes or environments that gave rise to interactions, but described only main effects effectively. These results confirmed to the observations made by Zobel *et al.* (1988). Since G x E interaction was highly significant (Table 1), ANOVA model was combined with PCA model to further analyze the residuals (GEI) of the ANOVA model.

The GEI was highly significant and was further partitioned into three PCA axes (IPCA) which jointly contributed 92.21 per cent of GEI. The first and second PCA were found highly significant ($P < 0.01$) and contributed 62.9 and 15.8 per cent to the total GEI variance, respectively and both the IPCA jointly 78.73 per cent of GEI. The residual SS accounted 7.79 % of the interaction SS and 21 % of df for GEI. (Gauch and Zobel, 1997).

The results of AMMI analysis can also be easily comprehended with the help of AMMI1 biplot as presented in Fig. 1. The mean performance of genotypes and environments vs IPCA1 score were used to construct the biplot (Table 2). According to AMMI model, the genotypes which are characterized by mean higher than the grand mean and the IPCA scores nearly zero are considered as generally adapted to all environments. However, the genotypes with high mean performance with large value of IPCA scores are considered to have specific adaptability to the

environments (Crossa *et al.* (1991), Pratap *et al.* (2009) and Abraham *et al.* (2013)). Biplot assay (Gabriel, 1971) presented in Fig. 1 identified three high yielding genotypes viz., G19, G18 and G6 having general adaptability, having mean yield >0.770 kg plot⁻¹ and close to IPCA1 = 0 line. Genotypes G3, G9 and G16 were higher yielding and specially adapted to favorable environments. Genotypes G1, G7, G13 and G14 had low mean yield and positioned near to IPCA1 = 0 line indicated that they were stable but they were lower yielding genotypes.

Similar sign of IPCA1 score for both genotypes and environment imply positive interaction and thus it attributed to higher yield of genotype at particular environment (Anandanet *al.*, 2009). IPCA scores of genotypes G3, G8, G9, G19 and G20 and of Devghadbaria (E3) location had positive sign, which indicated that these genotypes attributed higher yield at this location having positive GEI.

Table 2 : IPCA1 and IPCA2 scores of different back gram genotypes and locations.

| Sr. No. | Genotypes | Mean yield(kg plot ⁻¹) | Rank | IPCA1 | IPCA2 |
|---------|---------------------|------------------------------------|------|--------|--------|
| G1 | VUG-14 | 0.627 | 20 | -0.083 | -0.224 |
| G2 | VUG-18 | 0.729 | 14 | 0.297 | 0.075 |
| G3 | VUG-19 | 0.884 | 2 | 0.279 | 0.182 |
| G4 | VUG-23 | 0.722 | 15 | 0.089 | 0.073 |
| G5 | VUG-63 | 0.731 | 13 | -0.312 | -0.176 |
| G6 | VUG-32 | 0.870 | 3 | -0.161 | -0.324 |
| G7 | VUG-35 | 0.645 | 18 | -0.059 | 0.153 |
| G8 | DERUG-16-1 | 0.802 | 9 | 0.258 | 0.100 |
| G9 | DERUG-16-2 | 0.862 | 4 | 0.294 | -0.179 |
| G10 | DERUG-17-1 | 0.744 | 11 | 0.325 | -0.397 |
| G11 | DERUG-17-5 | 0.733 | 12 | 0.184 | 0.099 |
| G12 | DERUG-20-4 | 0.704 | 16 | -0.244 | 0.132 |
| G13 | DERUG-21-4 | 0.666 | 17 | -0.060 | 0.271 |
| G14 | DERUG-27-2 | 0.632 | 19 | -0.010 | -0.301 |
| G15 | DBUGP-2-2 | 0.828 | 7 | -0.293 | 0.070 |
| G16 | DBUGP-2-5 | 0.833 | 6 | -0.406 | -0.256 |
| G17 | DBUGP-6-1 | 0.850 | 5 | -0.175 | 0.125 |
| G18 | DBUGP-6-2 | 0.927 | 1 | -0.153 | 0.465 |
| G19 | T-9 | 0.823 | 8 | 0.062 | 0.209 |
| G20 | GU-1 | 0.784 | 10 | 0.168 | -0.096 |
| | Over all Mean | 0.770 | | | |
| | Environments | | | | |
| E1 | Vadodara | 0.526 | 4 | -0.154 | 0.532 |
| E2 | Jabugam | 0.482 | 5 | 0.750 | -0.352 |
| E3 | Devghadbaria | 1.001 | 1 | 0.154 | 0.061 |
| E4 | Derol | 0.905 | 3 | -0.142 | 0.409 |
| E5 | Dahod | 0.935 | 2 | -0.608 | -0.650 |

Correlation between PC1 and PC2 = 1.25E-11 ns

Environments were widely spread over scatter diagram which indicated high variability was among the locations. The environment E3, E4 and E5 were high yielding potential locations, whereas, E1 and E2 were found low yielding environments (Pratap *et al.*, 2009).

Visualization of which-won-where pattern of MLT data is important for studying the possible existence of different mega-environments. The polygon was drawn by connecting the markers of the genotypes that were far away from the biplot origin such that all other genotypes were contained in the polygon. The rays in Figure 2 were lines that were perpendicular to all the sides of the polygon. These five rays divided the biplot into five sectors and the environments were distributed in the four sectors. The interesting feature of this view of GGE biplot is that the vertex genotype for each sector was the best for the environment fall in the same sector than the others in all environments (Yan, 2002). Thus, environment E1 and E4 fell into sector III delineated by Rays 2 and 3 and the vertex genotypes for this sector was G18 suggesting that G18 was the winner high yielding genotype for Vadodara (E1) and Derol (E4) locations. Similarly, for environment E2 (sector I) the vertex genotypes was G10 which was the best for Jabugam location (E2). Sector IV delineated by Rays 3 and 4 contained the environment E5 and the winning genotypes for Dahod (E5) location was G16, which was higher yielding genotype. The discrepancy between the results of vertex genotype and higher yielding one may be because of the biplot explained 78.7% to variation of GEI and the remaining 22.3% variation was unaccounted (Pratap *et al.*, 2009, Joseph *et al.*, 2015)).

AMMI Stability Value (ASV)

The ASV is the distance from the co-ordinate point to the origin in two dimensions of IPCA1 scores against IPCA2 scores in the AMMI model. Genotype G4 followed by G7, G19 and G1 were the most stable, while genotype G16, G10 and G5 were undesirable in terms of stability (Table 3). Among the stable genotypes, yield of genotype G19 had higher mean than over all mean of grain yield. Similarly in unstable genotypes G16 had higher grain yield while G10 and G5 had lower grain yield than mean grain yield. Thus, present findings are in accordance with those reported by different workers (Joseph *et al.*, 2015; Pratap *et al.*, 2009, Nath and Dasgupta, 2013).

$W_{i(AMMI)}$ measure

The different $W_{i(AMMI)}$ values presented in Table 3 indicated that ranks of $W_{i(AMMI)}$ superimposed on ranks of AVS measures. According to the value of $W_{i(AMMI)}$, G4, G7 and

G19 were the most stable, while genotype G16, G10 and G5 were most unstable. (Raju and Bhatia, 2003).




Figure 1:- AMMI1 biplot for black gram genotypes and locations.




Figure 2:- AMMI2 biplot for black gram genotypes and locations.

AMMI based selection Index (ASTABi)

ASTAB_i (Rao *et al.*, 2004) stability measures (squared Euclidean distance) were calculated using the retained 'n' (3) axis out of 'N' total axis and presented in Table 3. Genotype G19 had the lowest value of ASTAB_i followed by G14 and G7 considered as stable. Among these genotypes, only G19 had higher grain yield than overall mean grain yield. The highest values of ASTAB_i were observed for G16, G10 and G7 which indicated their instability over environments.

Thus, overall results suggested that genotype G19 (T-9) was high yielding stable genotype which can be recommended for all five locations. Genotypes G4 (VUG-23) and G7 (VUG-35) gave low yield but they were stable. Devgadbaria, Derol and Dahod were high yielding whereas Vadodara and Jabugam were low yielding locations.

Table 3: Selection of genotypes based on different indices based on AMMI model for blackgram genotypes.

| Sr. No. | Genotype | Mean yield (kg plot ⁻¹) | Rank | ASV | Rank | W _{i(AMMI)} | Rank | ASATB _i | Rank |
|---------|------------|-------------------------------------|------|-------|------|----------------------|------|--------------------|------|
| G1 | VUG-14 | 0.627 | 20 | 0.279 | 4 | 0.033 | 4 | 0.042 | 5 |
| G2 | VUG-18 | 0.729 | 14 | 0.596 | 16 | 0.149 | 16 | 0.252 | 17 |
| G3 | VUG-19 | 0.884 | 2 | 0.586 | 14 | 0.144 | 14 | 0.232 | 14 |
| G4 | VUG-23 | 0.722 | 15 | 0.192 | 1 | 0.015 | 1 | 0.033 | 4 |
| G5 | VUG-63 | 0.731 | 13 | 0.646 | 18 | 0.175 | 18 | 0.280 | 18 |
| G6 | VUG-32 | 0.870 | 3 | 0.456 | 10 | 0.087 | 10 | 0.091 | 7 |
| G7 | VUG-35 | 0.645 | 18 | 0.192 | 2 | 0.016 | 2 | 0.032 | 3 |
| G8 | DERUG-16-1 | 0.802 | 9 | 0.524 | 12 | 0.115 | 12 | 0.188 | 13 |
| G9 | DERUG-16-2 | 0.862 | 4 | 0.613 | 17 | 0.158 | 17 | 0.250 | 16 |
| G10 | DERUG-17-1 | 0.744 | 11 | 0.760 | 19 | 0.243 | 19 | 0.324 | 19 |
| G11 | DERUG-17-5 | 0.733 | 12 | 0.380 | 9 | 0.061 | 9 | 0.098 | 9 |
| G12 | DERUG-20-4 | 0.704 | 16 | 0.503 | 11 | 0.106 | 11 | 0.170 | 12 |
| G13 | DERUG-21-4 | 0.666 | 17 | 0.296 | 5 | 0.037 | 5 | 0.052 | 6 |
| G14 | DERUG-27-2 | 0.632 | 19 | 0.302 | 6 | 0.038 | 6 | 0.021 | 2 |
| G15 | DBUGP-2-2 | 0.828 | 7 | 0.588 | 15 | 0.145 | 15 | 0.241 | 15 |
| G16 | DBUGP-2-5 | 0.833 | 6 | 0.850 | 20 | 0.304 | 20 | 0.473 | 20 |
| G17 | DBUGP-6-1 | 0.850 | 5 | 0.371 | 8 | 0.058 | 8 | 0.092 | 8 |
| G18 | DBUGP-6-2 | 0.927 | 1 | 0.556 | 13 | 0.130 | 13 | 0.107 | 11 |
| G19 | T-9 | 0.823 | 8 | 0.243 | 3 | 0.025 | 3 | 0.019 | 1 |
| G20 | GU-I | 0.784 | 10 | 0.349 | 7 | 0.051 | 7 | 0.101 | 10 |

REFERENCE

- Abraham, B., Vanaja, M., Raghu Ram Reddy, P., Sivraj, N., Sunil, N., Kamala, V. and Varaprasad, K. S. (2013). Identification of stable and high yielding genotypes in blackgram (*Vignamungo*(L.)Hepper) germplasm. *Indian J. Genet.* **73** (3): 264-269.
- Anandan, A.; Eswaran, R.; Sabesan, T. and Prakash, M. (2009). Additive main effect and multiplicative interactions analysis of yield performances in rice genotypes under coastal saline environments. *Advances in Biological Research*, **3**(1-2):43-47.
- Crossa, J.; Fox, P. N.; Pfeiffer, W. H.; Rajaram, S. and Gauch, H. G. (1991). AMMI adjustment for statistical analysis of an international wheat yield trial. *Theor. Appl. Genet.*, **81**:27-37.
- Gabriel K. R. (1971). The biplot graphic display of matrices with application to principal component analysis. *Biometrika*, **58** (3):453-467.
- Gauch, H. G. and Zobel, R. W. (1997). Identifying mega-environments and targeting genotypes. *Crop Sci.*, **37**:311-326.
- Huehn, M. (1979). Beitrage zur Erfassung der phanotypischen stabilitat. I Vorschlag einiger auf Ranginformationen beruhenden Stabilitatsparameter. *EDO in Medicine and Biology*, **10**:112-117.
- Joseph Jiji, Rose Mary Francis Santhoshkumar A.V Sunil, K.M and Dayalakshmi, E.M. (2015). Stability of blackgram (*Vignamungo*L. Hepper) varieties for seed yield. *Electronic Journal of Plant Breeding*, **6**(4): 899-903.
- Nassar, R. and Huehn, M. (1987). Studies on estimation of phenotypic stability: Tests of significance for non-parametric measures of phenotypic stability. *Biometrics*, **43**:45-53.
- Nath, D. and Dasgupta, T. (2013). Genotype x Environment Interaction and Stability Analysis in Mungbean. *J. of Agric. and Veterinary Sci.* **5**(1): 2319-2372.
- Pervin, M. A., Polash, M. F. M. B., Rahman, S. M. and Deb, A. C. (2007). Study of genetic variability and G x E Interaction of some quantitative traits in black gram (*Vgnamungo*(L.)Hepper). *J. of Biol. Sci.* **7**(1): 169-175.
- Pratap, K., Sudarshanam, A., Satyanarayana Rao, V., Srinivasa Rao, V. and Panduranga Rao, C. (2009). Phenotypic stability analysis in Greengram [*Vigna radiate* (L.)Wilczek] Using Eberhart and Russell and AMMI Models. *The Andhra Agric. J.* **56**(3): 288-297.

Purchase J. L. (1997). Parametric analysis to describe genotype x environment interaction and yield stability in winter wheat. Ph. D. Thesis, Department of Agronomy, Faculty of Agriculture of the University of the Free State, Bloemfontein, South Africa.

Raju, B. M. K. (2002). On some statistical aspects of assessing sensitivity of crop varieties. Unpublished Ph. D. Thesis, P.G. School, IARI, New Delhi.

Raju, B. M. K. and Bhatia, V. K. (2003). Comparison of various measures of stability with respect to ranking ability under varying situations. *J. Ind. Soc. Agric. Statist.*, **56**(3):276-293.

Rao, A. R.; Prabhakaran, V. T. and Singh, A. K. (2004). Development of statistical procedures for selecting genotypes simultaneously for yield and stability. IASRI publication. [URL: http://www.iasri.res.in/ebook/EB_SMAR/index Mod 3.htm](http://www.iasri.res.in/ebook/EB_SMAR/index_Mod_3.htm).

Thangavel P., A. Anandan and R. Eswaran (2011). AMMI analysis to comprehend genotype-by-environment (G×E) interactions in rainfed grown mungbean (*Vignaradiata*L.) *AJCS* **5**(13):1767-1775

Yan, W. (2002). Singular-value partitioning in biplot analysis of multi-environment trial data. *Agron. J.*, **94**:990-996.