

## Original Research Article

### **Study of correlation and path analysis in greengram [*Vignaradiata*(L.) Wilczek]for yield and yield attributing traits**

#### **Abstract**

The present investigation was undertaken to estimate the correlation coefficients among thirteen quantitative traits and to study the direct and indirect effects of various yield contributing traits on seed yield by path analysis in twentygreengram genotypes carried out during *Kharif* 2021-2022 in a randomized block design with three replications. The results of the association study revealed that,seed yield per plant showed a highly significant and positive correlation at both genotypic and phenotypic levels with plant height, days to 50% flowering, days to maturity, number of primary branches, number of clusters per plant, number of pods per plant, no. of seeds per pod, pod length, biological yield per plant, harvest index indicating the possibility of simultaneous improvement for these traits. Seed index at phenotypic levelhad highly significant but negatively correlated with seed yield per plant. Path analysis revealed that,thenumber of seeds per pod, biological yield per plant, harvest index and seed index recorded the highest direct effect at-in the desirable direction. Their association with seed yield was also significant and positive indicating a true and perfect association between these traits. Thereforedirect selection for these characteristicsecharacters would help in isolating high-high-yielding genotypes from highly segregating populations.

**Keywords:** Correlation, path analysis, greengram, seed yield.

#### **Introduction**

Greengram [*Vignaradiata* (L.) Wilczek], also known as mungbean, is an ancient pulse crop widely cultivated in India. It is a diploid species with the chromosomal number  $2n=2x=22$ , a member of the subfamily Papilionaceae of the Leguminosae family, and its botanical name is *Vignaradiata* (L.) Wilczek. Mungbean is a native of South Asia (India), and its likely ancestor is *Vignaradiatavar. sublobata*. It may be produced in a variety of crop rotation strategies due to its short duration, broad adaptability, low water need, and photo insensitivity. India, Pakistan, Bangladesh, Sri Lanka, Nepal, and other Southeast Asian nations are where it is mostly grown (Singh *et al.*, 2015).

Mungbean is one of the essential amino acids, including arginine, leucine, lysine, tryptophan, and valine, that is high in vitamin B among other pulses. Green gram sprouts contain vitamina C and E (Engel, 1978).Greengram is rich in protein (24gm/100gm) which is nearly 2.5 times more than cereals. It is also a good source of carbohydrate (60gm/100gm),

fat (1gm/100gm), minerals (3gm/100gm) and fiber (1gm/100gm). Mungbean seeds contain about 124 mg Calcium/100 gm, 326 mg Phosphorus/100 gm, 1.3% Fat, 7.3 mg Iron/100 gm, 4.1 % Fiber and ~~having~~ 334 Kcal Calorific Value. In our country peoples are predominantly vegetarian and pulses are ~~the~~ main sources of protein, thus its vital importance in daily diet.

Correlation estimates between yield and other ~~characteristic characters~~ are ~~helpful~~ ~~useful~~ in selecting desired plant types in designing an effective breeding programme. Correlation coefficient measures the degree of association, genotypic or phenotypic relationship between two or more ~~characteristic characters, which~~ forming the basis for selection. Path coefficients analysis (Wright, 1921) is an important tool for partitioning the correlation coefficient into direct and indirect effects of variables on dependent variables which can be an added advantage and helps in selection to a greater extent for improvement of yield.

### Materials and Methods

The study of ~~correlation and path analysis~~ ~~(need to be complete the methods)~~ in greengram for ~~yield and yield attributing traits~~ ~~(need to be complete the factors)~~ was carried out under the field condition during *Kharif* 2021-2022 in a randomized block design with three replications at ~~the~~ field experimental center and Seed Testing Laboratory, Department of Genetics and Plant Breeding, Sam Higginbottom Institute of Agriculture, Technology and Sciences Deemed to be University, Allahabad. The experimental area is situated on the left side of Allahabad Rewa road. It is nearly 5.0 km away from Allahabad city and very near to ~~the~~ river Yamuna. The experimental plot had uniform topography with homogenous fertility and this condition was suitable for ~~the~~ cultivation of greengram. The recommended packages of practices were followed for raising a healthy crop and all necessary plant protection measures were taken to control the pest and diseases. The experimental material consisted of 20 diverse lines of greengram (Table 1).

Formatted: Highlight

Formatted: Font color: Red

Formatted: Highlight

Formatted: Font color: Red

**Table: 1 Genotypes of Greengram with their pedigree**

Sno.	Genotypes names	Centre responsible for developing	Pedigree	Year of release
1	PUSA VISHAAL	IARI, New Delhi	Selected from NM92	2002
2	IIPM-99-125	IIPR, Kanpur	PM3 X APM36	2004
3	SML-832	PAU, Ludhiana	SML302 X PUSA BOLD1	2010

4	DGGV-2	UAS,Dharwad	CHINAMUNG X TM-98-50	2014
5	IPMD-604-1-7	IIPR, Kanpur	Germplasm	2016
6	PANT MUNG-5	GBPUAT, Pantnagar	Selected from VC 6368	2002
7	SML-668	PAU , Ludhiana	Selected from NM94	2002
8	PUSA-9531	IARI, New Delhi	Selected from NM9473	2000
9	OBGG-58	OUAT, Berhampur	Mutant of K857	2002
10	IPM-2-3	IIPR, Kanpur	IPM99-125 X PUSA BOLD2	2009
11	MH-3-18	CCSHAU, Hisar	Germplasm	2016
12	PUSA 0672	IARI, New Delhi	11/395 X ML267	2009
13	HUM-16	BHU,Varanasi	PUSA BOLD-1 X HUM8	2006
14	MH-2-15	CCSHAU, Hisar	PDM116 X GUJRAT-1	2007
15	PUSA-9072	IARI,New Delhi	PUSA-106 X 10-215	1995
16	HUM-1	BHU,Varanasi	BHUMI X PANT U -30	1999
17	MH-421	CCSHAU, Hisar	MUSKAN X BDYR2	2014
18	IPM-409-4	IIPR,Kanpur	PDM2881 X IPM 3-1	2020
19	IPM-312-20	IIPR, Kanpur	IPM-1 X SPS 5	2020
20	PANT MOONG -4	GBPUAT, Pantnagar	T44 X UPU-2	1997

The estimation of the correlation coefficient was done using the formula given by Searle (1961) and the test of significance was carried out by the method described by Snedecor and Cochran (1967). The correlation coefficient ~~were~~ ~~was~~ further partitioned into direct and indirect effects with the help of path coefficient analysis as suggested by Wright (1921) and elaborated by Dewey and Lu (1959). Seed yield was considered as a dependent variable as factors assumed to be influenced by the other characters called independent variables as causes (need to be revised).

Formatted: Highlight

Formatted: Font color: Red

## Results and Discussion

Crop improvement programmes depends to a large extent on the availability of sufficient variability and association among different characteristics ~~characters~~ which are the pre-requisite for executing an effective selection programme. Seed yield, being a complex quantitative trait, is dependent on a number of component characteristics ~~characters~~. Therefore, knowledge of the association of different components together with their relative contributions has immense value in selection.

## Correlation analysis

The genotypic and phenotypic correlation coefficients were computed among all characteristic ~~characters under study is~~ presented in Tables 2 and 3. In general, the genotypic correlation coefficients were comparatively higher than the corresponding phenotypic

correlation coefficient. Seed yield per plant (g) was found significant and positively correlated with plant height (0.529, 0.829), days to 50% flowering (0.252, 0.380), days to maturity (0.376, 0.425), number of primary branches (0.377, 0.425), ~~the~~ number of clusters per plant (0.634, 0.651), ~~the~~ number of pods per plant (0.529, 0.376), ~~the number no-~~of seeds per pod (0.327, 0.455), pod length (0.341, 0.446), biological yield per plant (0.901, 0.916), harvest index (0.448, 0.388) at both genotypic and phenotypic levels, respectively and with days to fifty percent pod setting (0.420) at ~~the~~ phenotypic level and showed negative significant correlation with seed index (-0.316) at phenotypic level. ~~A similar Similar~~ finding was earlier reported by Muthuswamy *et al.* (2019), Asari *et al.* (2019), ~~and~~ Ahmad and Belwal (2020). ~~The number~~ Number of seeds per pod had a positive and significant correlation with pod length (0.579, 0.696), biological yield per plant (0.381, 0.534) and seed yield per plant (0.327, 0.455) at both genotypic and phenotypic levels, respectively and ~~the~~ number of clusters per plant was found significant and positively correlated with ~~the~~ number of pods per plant (0.684, 0.762), biological yield per plant (0.593, 0.623) and seed yield per plant (0.634, 0.651) at both genotypic and phenotypic levels, respectively. ~~Similar findings was were~~ earlier reported by Azam *et al.* (2018), Kumar *et al.* (2018), ~~and~~ Muthuswamy *et al.* (2019).

#### **Path analysis**

The correlation coefficient becomes more meaningful when ~~the correlation coefficients correlation coefficient~~ are partitioned into components of direct and indirect effects through path analysis, because correlation coefficients indicate only the ~~inter-inter-~~relationship of the ~~characteristic characters~~ irrespective of cause and effect (Dewey and Lu, 1959). For path analysis, seed yield was taken as the dependent variable and all other ~~12 characteristic characters~~ used for correlation studies were taken as causal variables. The results are presented in tables 4 and 5.

Each factor influenced the yield by a direct contribution and indirect contribution through other variables with which it was correlated. Among all the ~~characteristic characters~~, positive direct ~~s~~ effect on seed yield per plant ~~were-was~~ recorded by no. of seeds per pod (0.0089, 0.0149), biological yield per plant (0.9497, 0.8767), harvest index (0.4131, 0.4344) and seed index (0.0042, 0.0018) at both genotypic and phenotypic levels, respectively. On the other hand, negative direct effects on seed yield per plant were recorded by plant height (-0.0305, -0.0012) and ~~the~~ number of primary branches (-0.0088, -0.0142) at both genotypic

and phenotypic levels, respectively. Similar findings were reported by Pariharet *al.* (2018), [and](#) Muthuswamyet *al.* (2019).

Number of seeds per pod exerted positive indirect effect *via* plant height (0.0053, 0.0077), days to 50% flowering (0.0020, 0.0021), days to fifty percent pod setting (0.0026, 0.0032), days to maturity (0.0018, 0.0035), [the](#) number of primary branches (0.0006, 0.0007), number of clusters per plant (0.0020, 0.0025), [the](#) number of pods per plant (0.0006, 0.0004), pod length (0.0062, 0.0086) and biological yield per plant (0.0047, 0.0057) at both genotypic and phenotypic levels, respectively. Biological yield per plant exerted positive indirect effect *via* plant height (0.7709, 0.4619), days to 50% flowering (0.5458, 0.3339), days to fifty percent pod setting (0.5834, 0.3171), days to maturity (0.5921, 0.4247), number of primary branches (0.3013, 0.2658), [the](#) number of clusters per plant (0.5914, 0.5195), [the](#) number of pods per plant (0.4043, 0.4480), [the number no-](#)of seeds per pod (0.5071, 0.3336), and pod length (0.3596, 0.2558) both at genotypic and phenotypic levels, respectively.

### Conclusion


From [the present investigation](#), ~~present investigation~~ it can be concluded that, [the](#) number of seeds per pod, biological yield per plant, harvest index and seed index were the major yield contributing [characteristics](#) ~~characters~~ which had [a](#) positive and significant association with grain yield per plant and also exhibited [a](#) high direct effect on grain yield per plant. Therefore, due emphasis should be given ~~on to~~ these [characteristic](#) ~~characters~~ in the selection which would help in isolating [high-high](#)-yielding genotypes from highly [segregated populations](#) ~~segregating population~~ to improve [the](#) yield potential of greengram.

### References

- Ahmad, S. and Belwal, V. (2020). Study of correlation and path analysis for yield and yield attributing traits in mungbean (*Vignaradiata*(L.) Wilczek). *I.J.C.S.*, **8**(1): 2140-2143.
- Asari, T., Patel, B.N., Patel, R., Patil, G.B. and Solanki, Chirag (2019). Genetic variability, correlation and path coefficient analysis of yield and yield contributing characters in mungbean (*Vignaradiata*(L.) Wilczek). *Internatiinal Journal of Chemical Studies*, **7**, **7**(4): 383-387.

- Azam, M.G., Hossain, M.A., Alam, M.S., Rahman, K.S. and Hossain, M. (2018).** Genetic variability, heritability and ~~correlation~~ correlation, path analysis in mungbean (*Vignaradiata*(L.) Wilczek). *Bangladesh J. Agril. Res.*, **43**(3): 407-416.
- Dewey, D.R. and Lu, K.H. (1959).** A correlation and path coefficient analysis of components of crested wheat grass seed production. *Agron.J.*,**51**: 515-518.
- Engel, R.H. (1978).** The importance of legumes as a protein source in Asian diet. *Proceeding of first international mungbean symposium, Taiwan*: 35-39.
- Kumar, A., Sharma, N.K., Kumar, R., Sanadya, S.K. and Sahoo, S. (2018).** Correlation and path analysis for seed yield and components traits in mungbean under arid environment. *International Journal of Chemical Studies*,**6**(4): 1679-1681.
- Muthuswamy, A., Jamunarani, M. and Ramakrishnan, P. (2019).** Genetic variability, character association and path analysis studies in mungbean [*Vigna radiate* (L.) Wilczek]. *International Journal of Current Microbiology and Applied Sciences*, **8**(4): 1136-1146.
- Parihar, R., Agrawal, A.P., Sharma, D.J. and Minz, Madhuri Grace. (2018).** Character association and path analysis studies on seed yield and its yield attributing traits in mungbean (*Vignaradiata*(L.) Wilczek).*J.P.P.*, **7**(1): 2148-2150.
- Searle SR.** The value of indirect selection: I. Mass selection. *Biometrics*. 1965; 21:682-707.
- Singh, C., Singh, P. and Singh, R. (2015).** Modern Techniques of Raising Field Crops. *Oxford & IBH Publishing Co. Pvt. Ltd.*, New Delhi: 386 p.
- Snedecor GW, Cochran WG.** Statistical methods. 6th ed. Oxford and IBH Pub. Co. New Delhi, 1967.
- Wright, S. (1921).** Correlation and causation. *Journal of Agriculture Research*, **20**: 557-587.

UNDER PEER REVIEW



**Table 2: Genotypic Correlation Coefficient for Yield and Its Related Traits in 20 Greengram Genotypes**

TRAITS	Plant height (cm)	Days to 50% flowering	Days to 50 % pod sett	Days to maturity	No. of Primary Branches	No. of clusters per plant	No. of pods per plant	No. of seeds/pod	Pod length(cm)	Biological yield /plant	Harvest index (%)	Seed Index (g)	Seed yield/ plant
Plant height(cm)	1.000	0.467**	0.498**	0.353*	0.112	0.617**	0.536**	0.598**	0.692**	0.812**	0.196	-0.153	0.829**
Days to 50% flowering		1.000	0.825**	0.874**	0.077	0.387*	0.398*	0.222	0.386*	0.575**	-0.370*	-0.441**	0.380*
Days to 50% pod sett			1.000	0.861**	0.105	0.385*	0.467**	0.292*	0.431**	0.614**	-0.361*	-0.365*	0.420**
Days to maturity				1.000	0.208	0.303*	0.439**	0.202	0.190	0.624**	-0.336*	-0.436**	0.425**
No. of Primary Branches					1.000	0.490**	0.246	0.062	0.242	0.317*	0.339*	0.096	0.425**
No. of clusters per plant						1.000	0.762**	0.228	0.433**	0.623**	0.184	-0.307*	0.651**
No. of pods per plant							1.000	0.073	0.017	0.426**	-0.022	-0.386*	0.376*
No. of seeds/pod								1.000	0.696**	0.534**	-0.102	-0.059	0.455**
Pod length(cm)									1.000	0.379*	0.225	0.181	0.446**
Biological yield/ plant										1.000	-0.017	-0.417**	0.916**
Harvest index (%)											1.000	0.165	0.388*
Seed Index (g)												1.000	-0.316*
Seed yield per plant													1.000

\*5%Level of significance

\*\*1%Level of significance

**Table 3: Phenotypic Correlation Coefficient for Yield and Its Related Traits in 20 Greengram Genotypes**

TRAITS	Plant height (cm)	Days to 50% flowering	Days to 50 % pod sett	Days to maturity	No. of Primary Branches	No. of clusters per plant	No. of pods per plant	No. of seeds/pod	Pod length	Biologic al yield /plant	Harvest index (%)	Seed Index (g)	Seed yield/ plant
Plant height(cm)	1.000	0.326*	0.414*	0.274*	0.094	0.209	0.304*	0.517**	0.592**	0.527**	0.125	-0.143	0.529**
Days to 50% flowering		1.000	0.941**	0.619**	0.078	0.232	0.237	0.140	0.271*	0.381*	-0.238	-0.345*	0.252
Days to 50% pod sett			1.000	0.596**	0.111	0.214	0.267*	0.215	0.299*	0.362*	-0.253	-0.294*	0.226
Days to maturity				1.000	0.169	0.252	0.381*	0.235	0.136	0.485**	-0.162	-0.400*	0.376*
No. of Primary Branches					1.000	0.338*	0.221	0.047	0.206	0.303*	0.278*	0.091	0.377*
No. of clusters per plant						1.000	0.684**	0.168	0.201	0.593**	0.242	-0.198	0.634**
No. of pods per plant							1.000	0.030	0.013	0.511**	0.156	-0.314*	0.529**
No. of seeds/pod								1.000	0.579**	0.381*	-0.051	-0.048	0.327*
Pod length(cm)									1.000	0.292*	0.176	0.163	0.341*
Biological yield/ plant										1.000	0.023	-0.346*	0.901**
Harvest index (%)											1.000	0.143	0.448**
Seed Index (g)												1.000	-0.253
Seed yield per plant													1.000

\*5%Level of significance

\*\*1%Level of significance

UNDER PEER REVIEW

**Table 4: Genotypic Path Coefficient for Yield and Its Related Traits in 20 Greengram Genotypes**

	Plant height (cm)	Days to 50% flowering	Days to 50 % pod sett	Days to maturity	No. of Primary Branches	No. of clusters per plant	No. of pods per plant	No. of seeds/pod	Pod length	Biological yield /plant	Harvest index (%)	Seed Index (g)	Seed yield/ plant
<b>Plant height(cm)</b>	<b>-0.0305</b>	-0.0143	-0.0152	-0.0108	-0.0034	-0.0188	-0.0164	-0.0183	-0.0211	-0.0248	-0.0060	0.0047	0.829**
<b>Days to 50% flowering</b>	-0.0013	<b>-0.0028</b>	-0.0028	-0.0024	-0.0002	-0.0011	-0.0011	-0.0006	-0.0011	-0.0016	0.0010	0.0012	0.380*
<b>Days to 50% pod sett</b>	0.0324	0.0655	<b>0.0650</b>	0.0559	0.0068	0.0250	0.0304	0.0190	0.0280	0.0399	-0.0235	-0.0237	0.420**
<b>Days to maturity</b>	-0.0236	-0.0583	-0.0574	<b>-0.0667</b>	-0.0138	-0.0202	-0.0293	-0.0135	-0.0127	-0.0416	0.0224	0.0290	0.425**
<b>No. of Primary Branches</b>	-0.0010	-0.0007	-0.0009	-0.0018	<b>-0.0088</b>	-0.0043	-0.0022	-0.0005	-0.0021	-0.0028	-0.0030	-0.0008	0.425**
<b>No. of clusters per plant</b>	0.0098	0.0061	0.0061	0.0048	0.0078	<b>0.0158</b>	0.0121	0.0036	0.0069	0.0099	0.0029	-0.0049	0.651**
<b>Number of pods per plant</b>	-0.0060	-0.0045	-0.0053	-0.0049	-0.0028	-0.0086	<b>-0.0113</b>	-0.0008	-0.0002	-0.0048	0.0002	0.0043	0.376*
<b>No. of seeds/pod</b>	0.0053	0.0020	0.0026	0.0018	0.0006	0.0020	0.0006	<b>0.0089</b>	0.0062	0.0047	-0.0009	-0.0005	0.455**
<b>Pod length(cm)</b>	-0.0076	-0.0042	-0.0047	-0.0021	-0.0026	-0.0047	-0.0002	-0.0076	<b>-0.0109</b>	-0.0041	-0.0025	-0.0020	0.446**
<b>Biological yield per plant</b>	0.7709	0.5458	0.5834	0.5921	0.3013	0.5914	0.4043	0.5071	0.3596	<b>0.9497</b>	-0.0163	-0.3956	0.916**
<b>Harvest index (%)</b>	0.0809	-0.1526	-0.1492	-0.1387	0.1402	0.0759	-0.0092	-0.0420	0.0928	-0.0071	<b>0.4131</b>	0.0679	0.388*
<b>Seed Index (g)</b>	-0.0006	-0.0018	-0.0015	-0.0018	0.0004	-0.0013	-0.0016	-0.0002	0.0008	-0.0017	0.0007	<b>0.0042</b>	-0.316*
<b>Seed yield per plant (g)</b>	0.829**	0.380*	0.420**	0.425**	0.425**	0.651**	0.376*	0.455**	0.446**	0.916**	0.388*	-0.316*	1.0000
<b>Partial R<sup>2</sup></b>	-0.0253	-0.0011	0.0273	-0.0284	-0.0037	0.0103	-0.0042	0.0040	-0.0049	0.8697	0.1604	-0.0013	

\*5%Level of significance



\*\*1%Level of significance

**Table 5: Phenotypic Path Coefficient for Yield and Its Related Traits in 20 Greengram Genotypes**

	Plant height (cm)	Days to 50% flowering	Days to 50 % pod sett	Days to maturity	No. of Primary Branches	No. of clusters per plant	No. of pods per plant	No. of seeds/pod	Pod length	Biological yield /plant	Harvest index (%)	Seed Index (g)	Seed yield/ plant
<b>Plant height(cm)</b>	<b>-0.0012</b>	-0.0004	-0.0005	-0.0003	-0.0001	-0.0003	-0.0004	-0.0006	-0.0007	-0.0006	-0.0002	0.0002	0.529**
<b>Days to 50% flowering</b>	0.0108	<b>0.0331</b>	0.0311	0.0205	0.0026	0.0077	0.0078	0.0046	0.0090	0.0126	-0.0079	-0.0114	0.2515
<b>Days to 50% pod sett</b>	-0.0092	-0.0209	<b>-0.0222</b>	-0.0132	-0.0025	-0.0047	-0.0059	-0.0048	-0.0066	-0.0080	0.0056	0.0065	0.2255
<b>Days to maturity</b>	0.0027	0.0060	0.0058	<b>0.0097</b>	0.0016	0.0024	0.0037	0.0023	0.0013	0.0047	-0.0016	-0.0039	0.376*
<b>No. of Primary Branches</b>	-0.0013	-0.0011	-0.0016	-0.0024	<b>-0.0142</b>	-0.0048	-0.0031	-0.0007	-0.0029	-0.0043	-0.0039	-0.0013	0.377*
<b>No. of clusters per plant</b>	-0.0002	-0.0002	-0.0002	-0.0002	-0.0003	<b>-0.0009</b>	-0.0006	-0.0001	-0.0002	-0.0005	-0.0002	0.0002	0.634**
<b>Number of pods per plant</b>	0.0035	0.0027	0.0031	0.0044	0.0026	0.0079	<b>0.0116</b>	0.0003	0.0001	0.0059	0.0018	-0.0036	0.529**
<b>No. of seeds/pod</b>	0.0077	0.0021	0.0032	0.0035	0.0007	0.0025	0.0004	<b>0.0149</b>	0.0086	0.0057	-0.0008	-0.0007	0.327*
<b>Pod length(cm)</b>	0.0001	0.0001	0.0001	0.0000	0.0000	0.0000	0.0000	0.0001	<b>0.0002</b>	0.0001	0.0000	0.0000	0.341*
<b>Biological yield per plant</b>	0.4619	0.3339	0.3171	0.4247	0.2658	0.5195	0.4480	0.3336	0.2558	<b>0.8767</b>	0.0198	-0.3030	0.901**
<b>Harvest index (%)</b>	0.0544	-0.1032	-0.1099	-0.0705	0.1209	0.1049	0.0679	-0.0221	0.0765	0.0098	<b>0.4344</b>	0.0621	0.448**
<b>Seed Index (g)</b>	-0.0003	-0.0006	-0.0005	-0.0007	0.0002	-0.0004	-0.0006	-0.0001	0.0003	-0.0006	0.0003	<b>0.0018</b>	-0.2531
<b>Seed yield per plant (g)</b>	0.529**	0.2515	0.2255	0.376*	0.377*	0.634**	0.529**	0.327*	0.341*	0.901**	0.448**	-0.2531	1.0000
<b>Partial R<sup>2</sup></b>	-0.0006	0.0083	-0.0050	0.0036	-0.0053	-0.0005	0.0061	0.0049	0.0001	0.7901	0.1944	-0.0005	

\*5%Level of significance

\*\*1%Level of significance



**Fig. 1: Genotypic path diagram for seed yield per plant** **Fig. 2: Phenotypic path diagrams for seed yield per plant** (explain it in text)

Formatted: Highlight

Formatted: Font color: Red