

Original Research Article

Carriage of group B Streptococcus in pregnant women in Dakar, Senegal: prevalence and associated factors.

Abstract

Background: Beta-hemolytic group B streptococcus (GBS) is a germ responsible for vaginal carriage which can be responsible for serious maternal-fetal disorders. The frequency, intermittency and consequences of this carriage are the main reasons for its research in pregnant women. The objective of this study was to assess this carriage and identify associated factors.

Method: A vaginal sample was taken from 158 pregnant women and inoculated on GRP.B STREP_DIFF.AGAR (Granada) chromogenic medium from Beckton Dickinson (BD) and then incubated in an oven at 37 °C. The reading took place at 24 and 48 hours. Group B Streptococcus appears in orange colonies. The data were collected, entered by Excel (version 2010) then analyzed by the SPSS software. 20.0. Chi-square test was used to analyze contingency tables with a significant difference if $p < 0.05$. **Results:** In sum, 158 pregnant women were included with a median age of 32 years. The carry rate was 25.7%. This carriage was variable according to the age of the patients (28.5% between 20 and 29 years and 21.0% after 40 years), the gestational age (27.4% in the second trimester and 27.2% in the third trimester), flora balance (39.3% intermediate flora and 24.1% unbalanced flora), parity (25.2% in nulliparas and primiparas), vulvovaginal candidiasis (25.5%) and a history of vaginal infections (26.0%). However, no significant difference was found between carrying GBS and these latter situations.

Conclusion: The intermittent nature and the potential complications of this carriage of GBS require its screening at the end of pregnancy in order to prevent any maternal-fetal disease.

Keywords: Group B streptococcus, vaginal carriage, prevalence.

1. Introduction

Group B beta-hemolytic streptococcus (GBS) or *Streptococcus agalactiae* is an encapsulated Gram-positive cocci whose human reservoir is digestive. It is also found in the vagina and this often-asymptomatic vaginal carriage can occasionally be responsible for genital infections of varying severity [1].

Indeed, it can be responsible for chorio-amnionitis, endometritis and urinary tract infection in pregnant or postpartum women [2].

It is also implicated in pregnant women in premature rupture of membranes and low birth weight [3,4].

In addition, its role in neonatal mortality and morbidity as well as in serious neonatal infections such as sepsis, meningitis and pneumonia are the reasons for its screening in pregnant women [1,5].

The prevalence of GBS colonization in pregnant women is variable around the world and approximately 40 to 75% of these mothers will pass the germ to their newborn [6].

However, although the incidence of neonatal infection with group B streptococcus-hemolytic disease remains at 1.35 to 5.4 newborns per 1000 live births, it remains serious for newborns [7].

Given the importance of maternal colonization and the pathogenicity of this bacterium, screening, prevention and treatment strategies must be implemented in each country [8].

Thus in Senegal, few data concerning the prevalence of carriage of *S. agalactiae* in pregnant women is available.

It is in this context that we initiated this work with the aim to determine prevalence of carriage of GBS in pregnant women and identify contributing factors.

2. Methodology

This was a descriptive cross-sectional retrospective study carried out at the Laboratory of Medical Biology (LBM) of the Pasteur Institute in Dakar between September and November 2020.

- Study population

It consisted of pregnant women who had come to the LBM of the Pasteur Institute in Dakar for a prenatal check-up.

- Taking of the samples

In women in a gynecological position respecting the conditions for taking a vaginal sample, a speculum was placed and samples from the exocervix and endocervix were taken.

For the exocervix sample, a vaginal swab was carried out for the detection of group B streptococcus. Also, another swab was carried out and then soaked in physiological water for the search for other pathogens and for the typing of the bacterial flora.

For the endocervix sample, swabs were taken for the detection of *Neisseria. gonorrhoeae*, *Chlamydia trachomatis* and mycoplasmas.

- Isolation and identification of *Streptococcus agalactiae*

Samples taken were immediately sent to the laboratory and immediately the swab taken from the vagina was inoculated on chromogenic medium GRP.B STREP_DIFF.AGAR (Beckton

Dickinson, unit kingdom) then incubated in the oven at 37 ° C for 18 to 24 hours.

The identification of *Streptococcus agalactiae* was made based on the orange color of the colonies on the medium but also by studying their morphological, cultural and biochemical characters.

Also the other swab taken from the exocervix was used for the typing of the bacterial forebore at the vaginal level but also for the search for other agents with high infectious risk.

Data analysis: The data (age, type of flora, carriage of Streptococcus B, associated infections, pregnancy, parity, history of infections, antibiotic treatment) were collected, entered by Excel (version 2010) then analyzed by the SPSS software. 20.0. Chi-square test was used to analyze contingency tables with a significant difference if $p < 0.05$.

3. Results

The study was conducted among 158 pregnant women with a median age of 32 years. The age group [30-39 years] was the most represented with 52.5% followed by that [20-29 years] with 35.4%.

Women in the 3rd trimester of pregnancy were in the majority in our population with a prevalence of 48.7% followed by those in the 2nd trimester with 32.3%.

Our study population consisted mostly of women with less than 2 children (nulliparous or first-time mothers) with 65.2% (**Table I**).

Table I : Socio-demographics characteristics

Socio-demographics characteristics	Number	percentage %
Age group (years)		
[20-29]	56	35,4
[30-39]	83	52,5
>=40	19	12,0
Parity		
≤ 1	103	65,2
≥ 2	55	34,8
Age of pregnancy		
1st trimester	30	19,0
2 nd trimester	51	32,3
3rd trimester	77	48,7
History of infections		
No	35	22,2
Yes	123	77,8
Treatment Antibiotics		
No	148	93,7
Yes	10	6,3

We looked for pathologies such as candidiasis and bacterial vaginosis which were 32.3% and 18.4% respectively.

Notions of a history of infections and antibiotic treatment were reported in 77.8% and 6.3% respectively.

The carriage of Group B Streptococcus was found in our population at 25.7% (**Table II**).

Table II: Vaginal pathologies

Pathologies	Number	percentage %
Carriage of group B Streptococcus		
No	119	74,3
Yes	39	25,7
Candidiasis		
No	107	67,7
Yes	51	32,3
Type of flora (Nugent's score)		
Normal (0-3)	101	63,9
Intermediate (4-6)	28	17,7
Vaginosis (7-10)	29	18,4

Depending on the age groups, this carry was estimated at 28.5% for the age group [20-29 years] while it was respectively 22.9% and 21.0% for the [30- 39 years] and those over 40 years old. We did not observe a statistically significant difference between the carriage of Streptococcus B and the age of the patients ($p = 0.693$).

This carriage was noted almost similarly in multiparous women ($P > 2$) and nulliparas or first-time mothers ($P < 2$) without any significant difference ($p = 0.823$).

We noted that carry occurred almost in the second and third quarter with prevalence of 27.4% and 27.2% respectively. We did not find a significant difference between the age of pregnancy and carriage of Streptococcus B ($p = 0.277$).

Concerning the balance of the vaginal flora, the carriage of Streptococcus B occurred more on an intermediate or unbalanced flora (vaginosis) with prevalences of 39.3% and 24.1% respectively without significant difference ($p = 0.133$).

Streptococcus B carriage was associated with candidiasis in 25.5% of cases, antibiotic

treatment in 40.0% and history of infections in 26.0% of cases. There was no significant difference between this carry and these different situations (**Table III**).

Table III: Streptococcus B vaginal carriage and associated factors

UNDER PEER REVIEW

Carriage of Group B Streptococcus			
Characteristics	n/N	%	p-value
Age group (years)			
[20-29]	16/56	28,5	<i>0,693</i>
[30 - 39]	19/83	22,9	
≥ 40	4/19	21,0	
Parity			
≤ 1	26/103	25,2	<i>0,823</i>
≥ 2	13/55	23,6	
Type of flora			
Normale	21/101	20,8	<i>0,133</i>
Intermediate	11/28	39,3	
Unbalanced (vaginosis)	7/29	24,1	
Age of pregnancy			
1st trimester	4/30	13,3	<i>0,277</i>
2 nd trimester	14/51	27,4	
3rd trimester	21/77	27,2	
Candidiasis			
Yes	13/51	25,5	<i>0,871</i>
No	26/107	24,3	
Treatment antibiotics			
Yes	4/10	40,0	<i>0,246</i>
No	35/148	23,6	
Antécédents d'infections			
Yes	32/123	26,0	<i>0,466</i>
No	7/35	20,0	

4. Discussion

Our study showed a prevalence of *Streptococcus agalactiae* carriage of 25.7%. Similar prevalences have been reported in other countries such as Morocco and Tanzania with respectively 20.2% and 23% [9,10]. In the United States, asymptomatic vaginal carriage of GBS has been estimated in 20-30% of pregnant women in late pregnancy[11].

However, low prevalence has been reported in other countries including Cameroon, Nigeria and Ethiopia with respectively 6.70%, 2.3% and 7.2% [12, 13,14].

The variation in the prevalence of *Streptococcus agalactiae* in the different countries could be due to differences in sampling size but also to differences observed in the sampling or diagnostic technique used.

Concerning the sampling technique, it plays an important role in the isolation of *Streptococcus agalactiae*. Indeed, the most important colonization of GBS being at the level of the lower third, the use of the speculum will mask the anterior and posterior surface of the vagina thus reducing the surface studied.

Thus, during the sample, it is recommended to stress the importance of scanning the interior of the vagina to the vestibule and the vulva as recommended by the National Agency for Accreditation and Assessment in Health (ANAES) hence the technique of introducing the swab without a speculum [15].

Also, a rectal sample would increase the sensitivity of the screening [16,17].

The rate of carriage is also correlated with the bacteriological technique employed. In fact, a vaginal sample seeded without selective enrichment, as recommended by ANAES, makes it possible to find GBS in 10% of pregnant women. On the other hand, inoculation with selective enrichment increases the positivity rate reaching 15% [18].

In the present study GBS was isolated more frequently from women of age group 20-29 (28.5%). High isolation frequencies were shown in women at this age in Nigeria in 2015 and Tanzania in 2013 with respectively 50% and 43.6% [19,20].

In our study, this carriage was found almost similarly in multiparous women ($P > 2$) and nulliparas or first-time mothers ($P < 2$) with an average of 24.4%.

In Nigeria, a prevalence quite similar to ours was observed in first-time mothers by Ezeonu et al. (25.5%) [21].

Musa Mohammed found in Ethiopia in 2010 a higher colonization rate among multiparas (22%) compared to first-time mothers (17.9%) [22].

However, in India Hajare reported in 2016 a higher prevalence among first-time mothers (46.7%) without any statistically significant difference as we observed in our study [23].

Our study also showed that carriage occurred almost in the second and third trimester with

prevalence of around 27%.

An almost similar prevalence was observed by Rabiei in a multicenter study conducted in 2006 where the prevalence of *Streptococcus B* carriage was 26.65% in women after the 20th week of amenorrhea (WA) [24].

On the other hand, Rohi had reported in 2011 in Ardabil (Iran) a carrying level estimated at 18.0% in women between 8 and 40 weeks of amenorrhea [25].

Some countries have adopted universal screening of all pregnant women by rectovaginal culture between 35 and 37 weeks of gestation, which has made it possible to reduce the incidence of *Streptococcus agalactiae* by 80% [26]. On the other hand, other countries such as France, Denmark, the Netherlands and Australia do not recommend universal screening between 35 and 37 weeks because it can be effective in terms of cost and impact on the carrying of *Streptococcus agalactiae* [27, 28].

Regarding conditions such as vaginosis and candidiasis, no connection with the carriage of *Streptococcus B* was found in our study.

Vaginosis, which is an imbalance of the vaginal flora, often creates an environment favorable to the multiplication of other bacteria.

We investigated other potential associated factors such as history of vaginal infections, treatment with antibiotics. However, we found no statistically significant difference.

5. Conclusion

The carriage of *Streptococcus agalactiae* is relatively high in pregnant women in our study. Thus, due to these formidable complications in the newborn and somewhere in the mother, effective screening and management strategies should be put in place and harmonized in order to limit any pathological situation that may be linked to this germ during pregnancy.

Data availability

The data used to support the conclusions of this study are included in the manuscript and are available from the corresponding author on reasonable request.

Conflicts of interest

The authors declare that they have no conflict of interest.

Thanks

We would like to thank all colleagues in the medical biology laboratory for their unwavering investment in carrying out this work, particularly the bacteriology unit. We would also like to thank all of the study participants.

Special mention to employees of the bacteriology virology department of Aristide Le Dantec Hospital.

References

1. **Bevilacqua G.** Prevention of perinatal infection caused by group B α -hemolytic streptococcus. *Acta Biomed Ateneo Parmense* 1999; 70 (5-6):87-94.
2. **Dani C, Martelli E, Rubaltelli FF.** Prophylaxis of group B beta-hemolytic streptococcal infections. *Acta Biomed Ateneo Parmense* 2000; 71 (1):541-5.
3. **Regan JA, Klebanoff MA, Nugent RP, Eschenbach DA, Blackwelder WC, Lou Y et al.** Colonization with group B streptococci in pregnancy and adverse outcome. VIP Study Group. *Am J Obstet Gynecol* 1996; 174:1354–60.
4. **Regan JA, Klebanoff MA, Nugent RP.** The epidemiology of group B streptococcal colonization in pregnancy. Vaginal Infections and Prematurity Study Group. *Obstet Gynecol* 1991; 77: 604-610.
5. **Mc Kenna DS, Iams JD.** Group B streptococcal infections. *Semin Perinatol* 1998 Aug; 22 (4):267-76.
6. **Clay LS.** Group B streptococcus in the perinatal period. A review. *J Nurse Midwifery* 1996 Sep Oct; 41(5):355-63.
7. **Beri R, Lourwood DL.** Chemoprophylaxis for group B streptococcus transmission in neonates. *Ann Pharmacother* 1997 Jan; 31(1):110-2.
8. **Jaureguy F, Carton M, Teboul J, Butel MJ, Panel P, Ghnassia JC.** Facteurs de risque et stratégie de dépistage de la colonisation par le streptocoque du groupe B chez la femme enceinte: résultats d'une étude prospective. *J Gynecol Obstet Biol Reprod (Paris)*. 2003 Apr;32(2):132-8.
9. **Ahlam Bassir, Hanane Dhibou, Majdi Farah, Lharmis Mohamed, Addebous Amal, Souraa Nabila et al.** Portage vaginal du streptocoque du groupe B chez la femme enceinte au niveau de la région de Marrakech. *Pan African Medical Journal*. 2016; 23:107
10. **Joachim A, Matee MI, Massawe FA, Lyamuya EF.** Maternal and neonatal colonisation of group B streptococcus at Muhimbili National Hospital in Dares Salaam, Tanzania: prevalence, risk factors and antimicrobial resistance. *BMC Public Health*.2009; 9:437.
11. **Gibbs RS, Schrag S, Schuchat A.** Perinatal infections due to group B streptococci. *Obstet Gynecol*. 2004 Nov;104(5 Pt 1):1062-76.
12. **Foumane P, Mboudou E, Dohbit JS, Nkemyim DC, Tchokoteu PE, Doh AS.** Streptocoque Beta Hemolytique du groupe B et conséquences materno-foetales observées à l'hôpital général de Yaoundé: Etude descriptive. *Clin Mother Child Health* 2009; Vol 6, N° 1 :995-1001

- 13. Sharmila V, Joseph NM, ArunBabu T, Chaturvedula L, Sistla S.** Genital tract group B Streptococcal colonization in pregnant women : a South Indian perspective. *J Infect Dev Ctries* 2011, 5(8):592-5.
- 14. Lakew ZW, Gebreegziabher TT, Teklu SM, Yigeremu MG.** The prevalence of Group B Streptococcus recto-vaginal colonization and antimicrobial susceptibility pattern in pregnant mothers at two hospitals of Addis Ababa, Ethiopia. *Rep Health.* 2014; 11:80.
- 15. Loulergue J, Couhé C, Grasmick C, Laudat P, Quentin R.** Sensibilité aux antibiotiques des souches de streptocoque du groupe B de portage vaginal isolées en France, 2003. *BEH.* 2004; 18: 69-70 Google Scholar
- 16. Jerbi M, Hidar S, Hannachi N, El Moueddeb S, Djebbari H, Boukadida J et al.** Facteurs de risque du portage du streptocoque du groupe B chez la femme enceinte à terme : étude prospective à propos de 294 cas. *Gynecol Obstet Fertil.* 2007 Apr;35(4):312-6. PubMed | Google Scholar
- 17. Mahmoud M, Yahyaoui G, Benseddik N.** Dépistage de streptocoque du groupe B au cours du troisième trimestre de grossesse au CHU Hassan II de Fès. *Revue Tunisienne d'Infectiologie.* Janvier 2011; 5(1): 12- 15. Google Scholar
- 18. Benitz WE, Gould JB, Druzin ML.** Preventing early-onset group B streptococcal sepsis: strategy development using decision analysis. *Pediatrics.* 1999 Jun;103(6):e76. PubMed | Google Scholar
- 19. Idih EE, Ezem BU, Onyegbule OA, Ododo NA, Onumajuru CC.** Prevalence of vaginal group-B-Streptococcus antibiotic and antigen sensitivity amongst parturients at the federal medical center owerri, Nigeria. *Nigerian Journal of Medicine,* Vol. 28 No. 1, January - March, 2019, ISSN 1115-2613
- 20. Matee, M. I., Massawe, F. A., & Lyamuya, E. F.** Maternal and neonatal colonisation of group B streptococcus at Muhimbili National Hospital in Dar es Salaam, Tanzania: prevalence, risk factors and antimicrobial resistance. *BMC public Health,* 2009 ;9(1), 1-7.
- 21. Ezeonu IM, Agbo MC.** Incidence and antimicrobial resistance profile of Group B Streptococcus (GBS) infection in pregnant women in Nsukka, Enugu State, Nigeria. *Afr J Microbiol Res* 2014;8:91-95.
- 22. Musa M, Daniel A, Yimtubezinash W, Demissie A.** Prevalence of group B Streptococcus colonization among pregnant women attending antenatal clinic of Hawassa Health Center, Hawassa, Ethiopia. *Ethiop J. Health Dev.* Janvier 2012; 26 (1): 36-42

- 23. Hajare V, Madhari LH, Singh HKG.** Antibiogram of Group B Streptococci isolated from the vagina of pregnant women in third trimester of pregnancy. *Peop J SciRes* 2012;5(2):22-26.
- 24. Rabiei S, Arab M, Yousefifi Mashouf R.** Epidemiologic pattern of vaginal colonization by group B Streptococcus in pregnant women in Hamadan, Central west of Iran. *Iran J Med Sci* 2006; 31: 106–108.
- 25. Rohi E, Ghasemi K, Agdam FK.** Incidence of nongonococcal infection in childbearing and pregnant women in ardabil. *Int J Mol Clin Microbiol* 2011; 1: 71–76.
- 26. Verani JR, Spina NL, Lynfield R, Schaffner W, Harrison LH, Holst A et al.** Early-onset group B streptococcal disease in the United States : potential for further reduction. *Obstet Gynecol.* 2014;123(4):828-37.
- 27.** Prevention of Early-onset Neonatal Group B Streptococcal Disease: Greentop Guideline No. 36. *BJOG.* 2017;124(12):e280-e305.
- 28. Melin P.** Neonatal group B streptococcal disease: from pathogenesis to preventive strategies. *Clin Microbiol Infect.* 2011;17(9):1294-303.