

Letter to Editor

Prevalence of Dengue Virus Serotypes of 2021 Dengue Outbreak in Dhaka, Bangladesh

Sir,

Dengue is a self-limiting febrile illness caused by four serotypes of dengue viruses (DENVs)-DENV1-4¹. Bangladesh being a tropical country is hyperendemic for dengue as it serves as an ideal habitat for the female *Aedes aegypti* and *A. albopictus*– mosquito vectors for DENVs, and thus, vulnerable to annual dengue outbreaks during the monsoon season which provides perfect egg-laying shallow, still-water habitats^{2,3,4}. While it mostly leads to silent or mild-moderate symptoms, disease severity is often increased with reemergence of a serotype of DENV following years of absence, especially in cases of heterologous secondary dengue⁵. To prevent a potential epidemic with surge of cases with increased severity and mortality rate, nationwide preparedness through identification of circulating DENV serotypes is instrumental. As such, this study aimed to identify the prevalence of circulating DENV serotypes in the 2021 dengue outbreak in Dhaka, Bangladesh.

it (VISASURE, CerTest Biotec SL, Spain) and Multiplex real-time RT-PCR with Dengue differentiation kit (Fast-track Diagnostics, Sliema, Malta) was later carried out with the RNA positive samples for serotyping the DENVs in the samples. Anti-dengue IgG and IgM testing was done using SD BIOLINE Dengue IgG/IgM WB Test kit (Standard Diagnostics, Inc., Gyeonggi-do, Republic of Korea).

Among the NS1-positive samples, DENV RNA was detected in all (100%) patients. Among them, 88 (97.8%) samples were positive for DENV3, while 2 (2.2%) were DENV2-positive. Only DENV3 was detected in all 60 samples from June 2021 to September 2021 while in October-November 2021, 28 samples (93.3%) were DENV3 positive, and 2 (6.7%) samples tested positive for DENV2 (figure 1). This study also revealed that sixty-one (67.80%) of the dengue-patients had secondary infections with 59 (67.05%) cases being DENV3 and 2 (100%) being DENV2. The result, however, was not statistically significant (Chi-square, $P>0.05$).




Figure 1: Monthly DENV serotype findings from June to November, 2021

This study, thus, identified DENV3 as the predominant DENV serotype in the 2021 dengue outbreak in Dhaka, Bangladesh with DENV2 being in co-circulation while observing the absence of DENV1 and DENV4. DENV1 was absent from circulation for the first time in the last decade^{6,7}. The dwindling prevalence of DENV1 and DENV2, and absence of DENV4 were substantiated by the findings of Institute of Epidemiology, Disease Control & Research (IEDCR), Bangladesh, which showed that while DENV1's prevalence had decreased from 50% in 2015 to 0% in 2021, DENV2's prevalence had also declined from 76% in 2016 to 0% in 2021⁷. Henceforth, while the surveillance reports of IEDCR had not reported any DENV2 cases in Bangladesh in 2021, this study found two DENV2 cases in October-November study period (Figure 1)⁷. Dengue being hyperendemic in Bangladesh explains the predominance of secondary dengue infections among the study population, as a significant percentage of the Bangladeshi population had prior exposures to dengue, particularly DENV2 which had been in circulation for the last few decades and DENV3 which has been prevalent since its reemergence in late 2017⁴.

This study concluded that DENV2 and DENV3 were the circulating DENV serotypes in the 2021 dengue outbreak in Dhaka, Bangladesh with DENV3 being predominant. It was also revealed that majority of the cases were secondary dengue infections. Since majority of the cases were secondary infections which may have increased disease severity, studies to ascertain the circulating DENV serotypes must be carried out annually in Bangladesh to prevent unexpected reemergence of DENV1 or DENV4 after possible prolonged absence, as such an unforeseen circumstance without nationwide preparedness could lead to a massive outbreak with significant casualties.

Reference.

1. Simmons CP, Farrar JJ, van Vinh Chau N, Wills B. Dengue. *New England Journal of Medicine*. 2012 Apr 12;366(15):1423-32.
2. Guzman MG, Gubler DJ, Izquierdo A, Martinez E, Halstead SB. Dengue infection. *Nature reviews Disease primers*. 2016 Aug 18;2(1):1-25.
3. Prapty CN, Rahmat R, Araf Y, Shounak SK, Rahaman TI, Hosen MJ, Zheng C, Hossain MG. SARS-CoV-2 and dengue virus co-infection: Epidemiology, pathogenesis, diagnosis, treatment, and management. *Reviews in Medical Virology*. 2022 Mar 3:e2340.
4. Shirin T, Muraduzzaman AK, Alam AN, Sultana S, Siddiqua M, Khan MH, Akram A, Sharif AR, Hossain S, Flora MS. Largest dengue outbreak of the decade with high fatality may be due to reemergence of DEN-3 serotype in Dhaka, Bangladesh, necessitating immediate public health attention. *New microbes and new infections*. 2019 May;29.
5. Soo KM, Khalid B, Ching SM, Chee HY. Meta-analysis of dengue severity during infection by different dengue virus serotypes in primary and secondary infections. *PloS one*. 2016 May 23;11(5):e0154760.
6. Akther T, Muraduzzaman AK, Parvin SM, Tabssum S, Munshi SU. Molecular & serological study of dengue virus-infected patients attending a tertiary hospital of Dhaka city, Bangladesh (2013 to 2016). *The Indian Journal of Medical Research*. 2019 Jul;150(1):96.
7. Dengue surveillance. IEDCR 2021. Available from: <https://iedcr.gov.bd/surveillances/f364364f-c826-43db-9dc6-68421983bce7>, accessed on December 20, 2021.