

***Plasmodium falciparum* infection and Associated Risk Factors in Pregnant Woman attending Bafang District Hospital, Semi-Rural area, West Region of Cameroon: A cross-sectional study**

ABSTRACT

Background: Malaria is a major public health problem in countries with limited resources. This morbid and fatal infection mainly affects vulnerable groups with weak immunity, such as children and pregnant women. In pregnant women, malaria leads to the occurrence of anaemia with the consequences of abortion, low birth weight (LBW) of the child and death of the baby. This study aimed to determine the *Plasmodium falciparum* (*P. falciparum*) infection and associated risk factors among pregnant women attending the Bafang district hospital (BDH).

Methods: A hospital based quantitative cross-sectional and descriptive study were conducted from May 30th, to June 24th, 2022 among 156 pregnant women attending the BDH, West Region of Cameroon. All pregnant women consented to participate in the study without any constraint. Blood samples were collected, thick film and stained blood smear examined for Plasmodium trophozoites and gametocytes respectively. Statistical analyses were performed using Epi Info v. 7 software with any p-value<0.05 considered statistically significant.

Results: The age of the study participants ranged from 14 to 48 years old (mean age of 27.41 ± 6.67 years). Of the 156 pregnant women tested, 25% (95% CI: 18.42%-32.55%) were infected with *P. falciparum*, a species found in the blood. *P. falciparum* infection was high in pregnant women aged < 30 years old (66.67%), in non-salaried (82.05%), in pregnant women in the second trimester of pregnancy (48.72%) with no difference statistically significant (p-value<0.05). Furthermore, the use of mosquito nets (OR= 3.88, 95% CI: 1.78-8.43, p-value=0.0003), taking intermittent preventive treatment (IPT) (OR= 2, 95, 95% CI: 1.31-6.59, p-value=0.006) and the number of IPT doses (p-value=0.0006) were risk factors statistically associated with *P. falciparum* infection.

Conclusion: This study revealed a high prevalence of *P. falciparum* among pregnant women attending BDH. The use of mosquito nets, the taking of IPT were the factors statistically associated with the infection.

Keywords: *P. falciparum*, infection, risk factors, pregnant women, BDH

ABBREVIATIONS

ANC: Antenatal care

BDH: Bafang district hospital

CI: Confidence interval

IQR: Interquartile range

IPT: Intermittent preventive treatment

IPT-SP: Intermittent preventive treatment with sulfadoxine pyrimethamine

LBW: Low birth weight

LLIMN: Long lasting impregnated mosquito net

OR: Odds ratio

SSA: sub Saharan African

WHO: World Health Organization

1. INTRODUCTION

Malaria is the greatest widespread parasitic disease worldwide mainly in sub-Saharan African (SSA) such as Cameroon. The disease is caused by intracellular erythrocyte parasites of the genus *Plasmodium* and transmitted to human by the bite of female *Anopheles* mosquitoes [1]. Of note, there are five different human malaria species worldwide such as *P. falciparum*, *P. vivax*, *P. malariae*, *P. knowlesi* and *P. ovale* with *P. falciparum* as main species found in SSA [2]. Low and middle-income countries of Asia, Western pacific and SSA bear the brunt of the disease with high prevalence [1]. Of the 91 countries recording malaria cases worldwide, around 80% of the total cases were from SSA countries [3]. An estimated about 229 million cases and 409,000 deaths worldwide in 2019 [4]. Malaria leads to the occurrence of anaemia which can have serious consequences in pregnant women, the foetus and the new-born [5]. This anaemia which could lead to hypoxia in the tissues, LBW with risk of neonatal death. Moreover, the anaemia could lead to a decrease in immunity, abortion, intrauterine growth retardation, premature delivery and increase the risk of death in the mother [5-6]. According to the WHO, about 11.6 million (34%) pregnancies exposed to malaria infection in the WHO African Region, which resulted in 900,000 to LBW, 25,000 maternal deaths, and 100,000 neonatal deaths [7]. Confronted with this situation, the intermittent preventive treatment of malaria with sulfadoxine pyrimethamine (IPT-SP) in all pregnant women attending the antenatal care (ANC) for the first time, the use of the long-lasting impregnated mosquito net (LLIMN) and the hygiene of the surroundings of the endemic areas such as Cameroon [7].

Despite all its prevention measures and the national malaria control program set up, the prevalence of malaria in Cameroon has been reported at 30.3% with more than one million cases and about 10,000 deaths recorded each year in health facilities [8]. Malaria is the first reason for consultation, the main cause of death among children and pregnant women in health facilities in Cameroon. According to the 2018 world report, the WHO estimates 6,228,154 cases and Cameroon is among the eleven SSA countries most affected by malaria [4]. Moreover, several studies have found high prevalence of malaria in pregnant women attending ANC in the North-West region in 2020 by Calvin et al., [9] and by Anchang et al. [10] in the South-West region with 18.0% and 16.0% respectively. However, data on the characteristics of malaria in pregnancy are sparse, particularly in rural areas, and it is an obstacle for adequate control strategies planning and implementation. The aim of this study was to determine the *P. falciparum* infection and associated risk factors among pregnant women attending the BDH in the semi-rural area West region of Cameroon.

3. RESULTS

3.1 Characteristics of study population

This study involved consecutively 156 pregnant women consulting the BDH. The age of the participants ranged from 14 to 48 years old, with a mean age of 27.41 years (Standard Deviation \pm 6.67), and a median age of 28 years (IQR: 23-32). The majority of participants, were under 30 years old (64.10%), were single (40.38%), were self-employed (76.92%), had a secondary level (82.05%), were in the second trimester of pregnancy (41.03%) and 76.28% had a secondary sector of activity.

Table 1. Characteristics of Study population

Variables	Total (N=156)	Percentage (%)
Age (min-max: 14-48 years)		
< 30 years	100	64.10
\geq 30 years	56	35.90
Matrimonial status		
Single	63	40.38
Cohabiting	44	28.21
Divorced	2	1.28
Married	47	30.13
Economic conditions		
Salary	36	23.08
No salary	120	76.92
Level of education		
No formal education	2	1.28
Primary	4	2.56
Secondary	128	82.05
Higher	22	14.10
Age of pregnancy		
First trimester	54	34.62
Second trimester	64	41.03
Third trimester	38	24.36
Activity sector		
Primary	13	8.33
Secondary	119	76.28
Tertiary	24	15.38

3.2 Distribution of associated risk factors among study population

Regarding the associated risk factors, the majority of participants were not on intermittent preventive treatment (55.77%), 56.42% had no IPT dose, 73.08% did not use a mosquito net (**Table 2**).

Table 2. Distribution of associated risk factors among study population

Variables	Total (N=156)	Percentage (%)
Intermittent Preventive Treatment (IPT)		
No	87	55.77
Yes	69	44.23
Number of doses (IPT)		
No dose	88	56.42
One dose	63	40.38
Two doses	4	2.56
Three doses	1	0.64
Use of the mosquito net		
No	42	26.92
Yes	114	73.08
Stagnant water around the house		
No	98	62.82
Yes	58	37.18

?: Percentage

3.3 Distribution of *P. falciparum* prevalence among study population

Out of 156 pregnant women included in this study, 25% (n=39, 95%CI: 18.42-32.55) were infected (figure 2).




Fig 2. Prevalence of *P. falciparum* among study population

3.4 Distribution of *P. falciparum* infection among study population

Table 3 shows a high prevalence of *P. falciparum* infection in pregnant women aged < 30 years (66.67%) vs. 33.33% for those aged 30 and over (p-value=0.69). Regarding economic conditions, the prevalence of *P. falciparum* was four times higher in non-salaried pregnant women (82.05%) vs. 17.95% for employees (p-value=0.51). Pregnant women with a level of education (secondary) were more infected with 87.18%, followed by those with a higher level (10.26%), p-value=0.71. In addition,

participants in the first second trimester of pregnancy were more infected (48.72%) vs. 33.33% and 17.95% for those in the first and third trimester respectively (p-value=0.44).

Table 3. Distribution of *P. falciparum* infection among study population

Variables			<i>P. falciparum</i> Infection				Khi-square	p-value
	Total		Yes		No			
	N=156	%	n= 39	%	n=117	%		
Age (min-max: 14-48 years)								
< 30 years	100	64.10	26	66.67	74	63.25	0.14	0.69
≥ 30 years	56	35.90	13	33.33	43	36.75		
Marital status								
Single	63	40.38	13	33.33	50	42.74	1.97	0.58
Cohabiting	44	28.21	13	33.33	31	26.50		
Divorced	2	1.28	0	0	2	1.71		
Married	47	30.13	13	33.33	34	29.05		
Economic conditions								
Salary	36	23.08	7	17.95	29	24.79	0.77	0.38
No salary	120	76.92	32	82.05	88	75.21		
Level of education								
No formal education	2	1.28	0	0	2	1.71	1.38	0.71
Primary	4	2.56	1	2.56	3	2.56		
Secondary	128	82.05	34	87.18	94	80.34		
Higher	22	14.10	4	10.26	18	15.38		
Age of pregnancy								
First trimester	54	34.61	13	33.33	41	35.04	1.65	0.44
second trimester	64	41.03	19	48.72	45	38.46		
Third trimester	38	24.36	7	17.95	31	26.5		
Activity sector								
Primary	13	8.33	3	7.69	10	8.55	2.40	0.29
Secondary	119	76.28	33	84.62	86	73.50		
Tertiary	24	15.38	3	7.69	21	17.95		

#: Percentage

3.5 Distribution of *P. falciparum* infestation and associated risk factors

It appears from this distribution that pregnant women on intermittent preventive treatment (IPT) were less infected (25.64%) vs. 74.36% for those who did not take IPT (OR=2.95; 95% CI: 1.31-6.59, p-value=0.006) (Table 4). *P. falciparum* infection varied significantly depending on the number of IPT doses, going from 76.92% for zero doses to 2.57% for those who took three doses (p-value=0.0001). Strangely, the infestation was slightly high among participants who used the mosquito net with 51.28% vs. 48.72% for those who did not use (OR=3.88; 95% CI: 1.78-8.43, p-value=0.0003) (Table 4).

Table 4. Distribution of *P. falciparum* infestation and associated risk factors

Variables	<i>P. Falciparum</i> Infestation						Khi-square	p-value
	Total		Yes		No			
	N=156	%	n= 39	%	n=117	%		
Intermittent Preventive Treatment (IPT)								
No	87	55.77	29	74.36	58	49.57	7.28	0.006
Yes	69	44.23	10	25.64	59	50.43		
Number of doses (IPT)								
No dose	88	56.42	30	76.92	58	49.57	21.99	0.0001
One dose	63	40.38	5	12.82	58	49.57		
Two doses	4	2.56	3	7.69	1	0.86		
Three doses	1	0.64	1	2.57	0	0		
Use of the mosquito net								
No	42	26.92	19	48.72	23	19.66	12.55	0.0003
Yes	114	73.08	20	51.28	94	80.34		
Stagnant water around the house								
No	98	62.82	24	61.54	74	63.25	0.03	0.84
Yes	58	37.18	15	38.46	43	36.75		

4. DISCUSSION

In the current study, the prevalence of *P. falciparum* infection among pregnant women attending the BDH was 25% (95%CI: 18.42-32.55). Compared with previous studies conducted in Cameroon, this result is higher than that reported in the South-West Region (16.0%) [10], in Ndop health district, North-West Region (19.3%) [12], and Sanaga-Maritime (22.9 %) [13]. Meanwhile, this finding is lower than that observed in Fouban Subdivision, West Region (39.4%) [14], in Manfe (53.4%) [15] and Mbakong Health Centre, North-West Region (29.30%) [16]. This difference may be due to the study period, the type of study, the sample size and the environmental conditions. In addition, this result could be justified by the geolocation of Cameroon, classified as a zone of high malaria endemicity [8]. Regarding the prevalence of *P. falciparum* infection according to socio-demographic characteristics, the study revealed that none have been associated. These results are contrary than that observed by [10, 17] who found that age, age of pregnancy, economic factors and marital status were associated with *P. falciparum* infection. This difference could be due to the small size of our sample. In this study, pregnant women in the first and second trimester of pregnancy were more infected than those in the third trimester. This result corroborates with studies conducted in the Republic of Ethiopia [17, 18, 19] who found that pregnant women in the first trimester of pregnancy had an increased risk of having malaria than women in the third trimester.

The present study showed a statistically significant association between IPT and *P. falciparum* infection. Thus, women on IPT were less infected (OR= 2.95; 95% CI: 1.31-6.59, p-value=0.006) meaning that pregnant women who did not take IPT were three times more likely to be infected. In addition, the incomplete dose of IPT was found to be associated with the development of *P. falciparum* infection. Our results showed that the infection with *P. falciparum* decreased drastically according to the number of doses thus going from 76.92% in pregnant women without doses of IPT to 2.57% in pregnant women who were on their third dose of IPT (p-value=0.0001) meaning that IPT has a positive impact on reducing malaria in pregnant women. Similar results were reported in a clinical study that the prevalence of malaria in pregnant women was associated with non-use of Fansidar during pregnancy for IPT [20, 21,22, 23].

Alarming fact, the use of the mosquito net was statistically associated with the infection but with a high prevalence among pregnant women who declared using the mosquito net, with 51.28% vs. 48.72% for those who declared not using the mosquito net. In the first hand, this result could be justified by the type of study (cross-sectional study) and the statements given by the pregnant women during the study. Secondly, this result could be explained by the use of mosquito nets by these pregnant women. Of note, the most widely used interventions to prevent malaria in pregnancy are insecticide-treated bed nets, including long-lasting insecticide-treated bed nets (LLINs) and IPT in pregnancy. These interventions have shown a considerable reduction in morbidity and mortality due to malaria in children and pregnant women. Furthermore, these interventions were associated with a decrease in maternal parasitaemia and a reduction in maternal anaemia. [24, 25, 26, 27, 28].

5. CONCLUSION

Malaria remains a real public health problem in Cameroon despite the measures put in place to fight against the disease. Vulnerable population such as pregnant women and children are groups at risk. The present study reported a *P. falciparum* prevalence of 25% (95%CI: 18.42-32.55) among pregnant women attending the BDH. This prevalence was high in pregnant women aged under 30 years old, in the self-employed, in those with a secondary education, in those in the second trimester of pregnancy and in those in the secondary sector, with no statistically significant difference. However, the risk factors statistically associated with the development of malaria during pregnancy were the administration of intermittent preventive treatment as methods of malaria prevention and the use of mosquito nets.

CONSENT AND ETHICAL APPROVAL

Research proposal was approved by the Ethics Committee of the Regional Delegation of Public Health for the Centre Region, Yaounde-Cameroon and Ethic clearance was issued (Identification: CEN°451CRERSHC/2022 from 15 April 2022). Both official's language (French and English) were used to explain the objectives of the study. Only volunteer pregnant women who signed an informed consent form for their participation and that of their newborn were enrolled. Participants found infected were referred to clinicians of the health facility concerned for appropriate treatment.

REFERENCES

1. WHO. World Health Organisation. World malaria report 2020: 20 years of global progress and challenges. Geneva: Licence: CC BY-NC-SA 3.0 IGO. World Health Organization 2020; 2020:18–34.

2. Kwenti TE, Kwenti TDB, Njunda LA, Latz A, Tufon KA, Nkuo-Akenji T. Identification of the Plasmodium species in clinical samples from children residing in five epidemiological strata of malaria in Cameroon. Trop Med Health. 2017; 45 (14). <https://doi.org/10.1186/s41182-017-0058-5>
3. WHO. World Health Organisation. World malaria report 2016. Accessed May 23th, 2022.
4. WHO. World Health Organization. World malaria report 2019 World health organization 2019.
5. Bakken L, Iversen PO. The impact of malaria during pregnancy on low birth weight in East-Africa: a topical review. Malaria Journal. 2021 ; 20 : 348. <https://doi.org/10.1186/s12936-021-03883-z>
6. Dombrowski JG, de Souza RM, Lima FA, Bandeira CL, Murillo O, de Sousa Costa D, et al. Association of malaria infection during pregnancy with head circumference of newborns in the Brazilian Amazon. JAMA. 2019; 2(5):193–300. <https://doi.org/10.1001/jamanetworkopen.2019.3300> PMID: 31050780
7. WHO. World Health Organisation. World malaria report 2021. <https://www.who.int/publications/i/item/9789240040496> (2021). Accessed March 14th, 2022.
8. WHO. World Health Organisation. World malaria report 2018. Geneva: World Health Organization; 2018. Licence:CCBY-NC-SA3.0
9. Bisong EC, Etindele EFN, Kwende-Tanjong LO, Musoro JZ, Yamssi C, and Kuokuo KH. Plasmodium falciparum Parasitaemia during Pregnancy and the Use of Malaria Prevention Methods by Women Attending Antenatal Consultation at the Regional Hospital Bamenda, Northwest Cameroon. *International Journal of Tropical Disease & Health*. 2020;41(20): 46-56, Article no. IJTDH.63874 ISSN: 2278–1005, NLM ID: 101632866
10. Anchang-Kimbi JK, Ngenwie NV, Ngum NH, Apinjoh TO, Bantar TR, Fru Chi H, et al. Plasmodium falciparum parasitaemia and malaria among pregnant women at first clinic visit in the mount Cameroon. BMC Infectious Diseases. 2015:439 DOI 10.1186/s12879-015-1211-6
11. <https://mapcarta.com/fr/W894942019/Carte> accessed September 10th, 2022
12. Sumo L, Mbah NE, Nana-Djeunga CH. Malaria in pregnancy in the Ndop Health District (North West Region, Cameroon): results from retrospective and prospective surveys. Journal of Parasitology and Vector Biology. 2015;7(9):177-181.
13. Tonga C, Kimbi HK, Anchang-Kimbi JK, Nyabeyeu HN, Bissemou ZB, Lehman LG. Malaria risk factors in women on intermittent preventive treatment at delivery and their effects on pregnancy outcome in Sanaga-Maritime, Cameroon. PLoS ONE. 2013;8:e65876.
14. Ngouyamsa NAS, Vincent KP, Yamssi C, Noumedem ACN. "Effect of Impregnated Mosquito Bed Nets on the Prevalence of Malaria among Pregnant Women in Fouban Subdivision, West Region of Cameroon", Journal of Parasitology Research, vol. 2020, Article ID 7438317, 10 pages, 2020. <https://doi.org/10.1155/2020/7438317>
15. Elime FA, Nkenyi NR, Ako-Egbe Luis, Njunda A, and Nsagha D. Malaria in Pregnancy: Prevalence and Risk Factors in the Mamfe Health District, Cameroon. Journal of Advances in Medicine and Medical Research. 2019; 30(1): 1-11. Article no. JAMMR.49667 ISSN: 2456-8899
16. Ndong IC, Van RM, Boakye DA, Mbacham WF, Grobler AF. Trends in malaria admissions at the Mbakong Health Centre of the North West Region of Cameroon: a retrospective study. Malaria Journal. 2014; 13:328.

17. Gontie GB, Wolde HF. & Baraki AG. Prevalence and associated factors of malaria among pregnant women in Sherkole district, Benishangul Gumuz regional state, West Ethiopia. *BMC Infect Dis.* 2020; 20, 573. <https://doi.org/10.1186/s12879-020-05289-9>
18. Almaw A, Yimer M, Alemu M, Tegegne B. Prevalence of malaria and associated factors among symptomatic pregnant women attending antenatal care at three health centers in north-west Ethiopia. *PLoS ONE.* 2022; 17(4): e0266477. <https://doi.org/10.1371/journal.pone.0266477>
19. Tilahun A, Yimer M, Gelaye W, Tegegne B. Prevalence of asymptomatic Plasmodium species infection and associated factors among pregnant women attending antenatal care at Fendeka town health facilities, Jawi District, North west Ethiopia: A cross-sectional study. *PLoS ONE* 2020;15(4):0231477. pmid:32315341
20. Azubike K, Lucky O, Chukwuemeka A, Chukwudi R, Nwabunike E. Adherence to intermittent preventive treatment for malaria with sulphadoxine-pyrimethamine and outcome of pregnancy among parturients in South East Nigeria. *Patient Preference and Adherence.* 2014; 8:447– 51.
21. Vincenz C, Dolo Z, Saye S, Lovett JL, Strassmann BI. Risk factors for placental malaria, sulfadoxine-pyrimethamine doses, and birth outcomes in a rural to urban prospective cohort study on the Bandiagara Escarpment and Bamako, Mali. *Malar Journal.* 2022; 21:110. <https://doi.org/10.1186/s12936-022-04125-6>.
22. COSMIC Consortium, Community-based Malaria Screening and Treatment for Pregnant Women Receiving Standard Intermittent Preventive Treatment With Sulfadoxine-Pyrimethamine: A Multicenter (The Gambia, Burkina Faso, and Benin) Cluster-randomized Controlled Trial, *Clinical Infectious Diseases*, Volume 68, Issue 4, 15 February 2019, Pages 586–596, <https://doi.org/10.1093/cid/ciy522>
23. Dongang NRR, Hawadak J, Kojom FLP, Kumar A, Chaudhry S, Arya A. Intermittent preventive treatment with Sulfadoxine pyrimethamine for malaria: a global overview and challenges affecting optimal drug uptake in pregnant women, *Pathogens and Global Health.* 2022. DOI: 10.1080/20477724.2022.2128563
24. Jayasooriya S, Hislop A, Peng Y, Croom-Carter D, Jankey Y, Bell A, Dong T, Rowland-Jones S, Rickinson A, Walther M, and Whittle H, Revisiting the effect of acute *P. falciparum* malaria on Epstein-Barr virus: host balance in the setting of reduced malaria endemicity. *PLoS ONE.* 2012; 7(2): e31142. <http://dx.doi.org/10.1371/journal.pone.0031142> PMID:22347443
PMCID:3275582
25. Mwangu LM, Mapuroma R, & Ibisomi L. Factors associated with non-use of insecticide-treated bed nets among pregnant women in Zambia. *Malaria Journal.* 2022; 21:290. <https://doi.org/10.1186/s12936-022-04313-4>
26. Hill J, Hoyt J, van Eijk AM, ter Kuile FO, Webster J, Steketee RW. Prioritizing Pregnant Women for Long-Lasting Insecticide Treated Nets through Antenatal Care Clinics. *PLoS Med.* 2014; 11(9): e1001717. <https://doi.org/10.1371/journal.pmed.1001717>
27. Aung PL, Win KM, Show KL. Utilization of insecticide-treated bed nets among pregnant women in Myanmar—analysis of the 2015–2016 Demographic and Health Survey. *PLoS ONE.* 2022; 17(3): e0265262. <https://doi.org/10.1371/journal.pone.0265262>

28. Duut TB, Alhassan AR. Factors associated with ownership of insecticide-treated nets for malaria prevention among pregnant women in Ghana. *Public Health and Toxicology*. 2022;2(3):13. doi:10.18332/pht/152624.

UNDER PEER REVIEW