

**Immuno-Protective Effect of Aqueous Extract of *Ocimum gratissimum* Leaf in Cyclophosphamide-Induced Myelotoxicity in Wistar Rat.**

**ABSTRACT**

The bone marrow is the manufacturing center of blood cells; the suppression of its activity causes a deficiency of blood cells. This condition can rapidly lead to life-threatening infections. *Ocimum gratissimum* plays a vital role in preventing some disease conditions, but its hematological effects on the bone marrow have not been documented much. This study is designed to investigate the immuno-protective activity of the aqueous leaf extract of *Ocimum gratissimum* in cyclophosphamide-induced myelotoxicity. Twenty-four Wistar rats aged two to three months, weighing 170-200 g were used for the study. The rats were divided into six groups of four rats each, labeled A to F. The rats in the control group (group A) were given feed and water. Group B was administered with 3 mg/kg of cyclophosphamide intraperitoneally daily for seven days to induce myelotoxicity. Groups C and D were treated daily with 100 mg/kg and 200 mg/kg of the extract for fourteen (14) days respectively. Groups E and F were treated with the extract for seven (7) days and then induced and treated with myelotoxicity and extract simultaneously for the next seven (7) days. On the 15<sup>th</sup> day, blood samples (3.0 ml) were collected from each rat through the retro-orbital plexus. Also, the bone marrow was harvested and analyzed using a standard histological procedure. Data were analyzed with Pearson's correlation test and multivariate analysis of variance using Statistical Package for Social Sciences (SPSS) version 21 and results were expressed as mean  $\pm$  SD. Myelotoxicity was achieved in Group B rats. Group E and F rats showed a significant increase in hemoglobin (Hb), hematocrit (Hct), and total white blood cell count (TWBC) compared with Group B. The Group C and D rats revealed a significant increase in Hb, Hct, and TWBC when compared with group A. Aqueous leaf extract of *Ocimum gratissimum* may possess immuno-protective properties when orally administered in cyclophosphamide-induced bone marrow suppression.

**Keywords;** *Ocimum gratissimum*, cyclophosphamide, hematological parameters, leaf extract, myelosuppression.

**INTRODUCTION**

Cancer and its treatment have over the years posed great difficulties in the field of medicine. Some of the treatment options like chemotherapy have discouraging side effects which on their own pose a threat to the life of the oncology patient. Myelosuppression and liver injury is the most common dose-limiting side effect of chemotherapy <sup>[1]</sup>. The bone marrow is the spongy tissue in the center of the bones. It is where red blood cells (RBCs), white blood cells (WBCs), and platelets are made. With bone marrow suppression, the bone marrow doesn't make normal numbers of blood cells. It leads to lower levels of one or more types of blood cells. Bone marrow suppression is also called myelosuppression.

The medicinal plants are rich in secondary metabolites (which are potential sources of drugs) and essential oils of therapeutic importance. The important advantages claimed for therapeutic uses of medicinal plants in various ailments besides being economical, effective, and easily

available, are their safety<sup>[2]</sup>. Because of these advantages, medicinal plants have been widely used by traditional medical practitioners in their day-to-day practice. According to a survey (1993) by World Health Organization (WHO), the practitioners of the traditional system of medicine treat about 80% of patients in India, 85% in Burma, and 90% in Bangladesh<sup>[3]</sup>.

Herbal medications are used widely in developing countries for the treatment of various diseases and ailments. This is because they are seen as alternatives to orthodox medicines in terms of costs and perceived side effects<sup>[4]</sup>. One of the main problems found in the field of medicinal plants is the lack of pharmacological, toxicological, and clinical evidence. In many cases, this practice is commonly associated with their traditional use. Only a small fraction of thousands of medicinal plants used in the world have been rigorously tested in controlled studies, therefore the evidence of toxicity risks and verification of efficacy is even lower<sup>[5]</sup>.

***Ocimum gratissimum***, also known as **Clove Basil**, **African Basil**, and in Hawaii as **Wild Basil**, is a specie of *Ocimum*. It is native to Africa, Madagascar, Southern Asia, and the Bismarck Archipelago, and naturalized in Polynesia, Hawaii, Mexico, Panama, West Indies, Brazil, and Bolivia. *Ocimum gratissimum* is one of the leafy vegetables consumed by Nigerians. It is usually seen as a small shrub with many branches and simple oval leaves. The leaves are used as food additives for their believed nutritive and medicinal values<sup>[6]</sup>. It also adds flavor or aroma to the food. It is used mainly as a spice<sup>[7][8]</sup>. It is a medicinal plant that can be used for therapeutic purposes or as a precursor for the synthesis of curative drugs<sup>[9]</sup>. It can be used as a flavoring agent in a pharmaceutical product. The *Ocimum gratissimum* oil could be used to mask the bitter taste of most drugs, particularly the anti-malaria tablets<sup>[10]</sup>. Although much has been documented on the medicinal properties of this plant, this work, however, is designed to evaluate the myelo-protective property of the extract of this plant to know the best extract to use in the treatment or amelioration of myelotoxicity in folk medicine.

## MATERIAL AND METHODS

### Plant Material Collection

Fresh plant leaves of *Ocimum gratissimum* were harvested from a garden in the University of Nigeria, Enugu Campus. The plant was authenticated by a taxonomist at the Department of Botany, University of Nigeria, Nsukka.

### Plant Extraction and Phytochemical Analysis

Fresh leaves of *Ocimum gratissimum* were washed in fresh water and air dried in the laboratory at room temperature. The air-dried leaves were milled into fine powder in an electric blender and the fine powder was soaked in 2 liters of distilled water for 24 hours, filtered with Whatman No.1 filter paper (150 mm), and evaporated with a water bath / rotary evaporator at 50°C to crude extract of *Ocimum gratissimum*. Crude extract residue was weighed and dissolved in distilled water for use on each day of our experiment<sup>[11]</sup>. 10 g of crude extract was dissolved in 100 ml of normal saline to get the stock solution of extract used for daily administration.

Phytochemical screening was carried out to qualitatively assess the secondary metabolites in the plant leaf extract using the standard laboratory procedures of precipitation and coloration reaction as described by Sofowara<sup>[6]</sup> to identify the secondary metabolites present.

### Animal Procurement

Twenty-four (24) male (170g-200g) adult Wistar rats were used for the experiment. After being purchased from the animal house of College of Medicine, University of Nigeria, Enugu Campus, they were housed in aluminum cages, placed in a well-ventilated house with optimum condition (temperature 30°C photoperiod; 12hours natural light and 12hours dark; humidity is 40-50%). The animals were fed growers mash manufactured by Top Feed Nigeria Limited and allowed water *ad*

*libitum*. They were allowed an acclimatization period of 2 weeks and throughout the experimental period; the animals were handled according to the guidelines for animal research in the National Institute of Health guidelines for care and use of laboratory animals. The study was carried out per the principles of laboratory animal care and standard experimental procedure.

### Ethical Approval

Ethical clearance was sought for and approved by the Research and Ethical Clearance committee, Faculty of Basic Medical Sciences, College of Medicine, Enugu State University of Science and Technology (ESUCOM/FBMS/ETR/2022/013).

**TABLE 1: EXPERIMENTAL DESIGN AND ADMINISTRATION**

EXPERIMENTAL GROUPS	RATS	ADMINISTRATION	TREATMENT
<b>GROUP A (-ve control)</b>	4	CONTROL	Distilled water and food pellets were only given for 14 days.
<b>GROUP B</b>	4	Cyclophosphamide (3mg/kg)	3mg/kg body weight of cyclophosphamide was given without treatment per day for 14 days.
<b>GROUP C (Extract LD)</b>	4	OC (100mg/kg)	100mg/kg of the extract was only given daily for 14 days.
<b>GROUP D (Extract HD)</b>	4	OC (200mg/kg)	200mg/kg of the extract was only given daily for 14 days.
<b>GROUP E (Protective)</b>	4	Cyclophosphamide 3mg/kg+OC 100mg/kg	100mg/kg of the extract was given per day for 7 days before myelotoxicity was induced through the administration of 3mg/kg of cyclophosphamide per day for the next seven days.
<b>GROUP F (Protective)</b>	4	Cyclophosphamide 3mg/kg + OC 200mg/kg	200mg/kg of the extract was given per day for 7 days before myelotoxicity was induced through the administration of 3mg/kg of cyclophosphamide per day for the next seven days.

\*OC = *Ocimum gratissimum*.

HD = High dose; LD = Low dose

### Body Weight Measurement and Animal Sacrifice

Measurement of the body weight of all the animals in each group

The body weights of all the animals in each group were measured and recorded on the first and last days of study before sacrifice. The measurements were done using a digital electronic scale and recorded to the nearest gram. At the end of the 14 days treatment period, the animals were sacrificed with an anesthetic agent called thiopental at a dose of 50mg/kg body weight which was injected intraperitoneally 24 hours post-treatment.

### Blood Analysis

On the 15<sup>th</sup> day, Blood samples were collected from each of the groups with the aid of capillary tubes through the medial optical vessels for evaluation of blood parameters (complete blood

count). Complete blood count was performed with an automated hematology system (Sysmex KX-2N hematology analyzer, Sysmex Incorporation Kobe, Japan).

### Histological Study

The histological analyses of the tissues were done according to the method of Drury [12]. The stained tissues were micrographed and interpreted by a pathologist at the University of Nigeria, Nsukka.

### Statistical Analysis

Results of hematological analysis (Total blood count) were analyzed using SPSS (version 21), expressed as Mean  $\pm$  Standard Error of Mean (SEM). Statistical differences in the means between groups were analyzed using one-way ANOVA (Analysis of variance). P-values less than 0.05 were considered to be statistically significant.

## RESULTS

### Result of Phytochemical Analysis

The phytochemical analysis of the aqueous leaf extract of *Ocimum gratissimum* revealed the presence of tannins, steroids, terpenoids, flavonoids, and cardiac glycosides. Anthraquinones were also detected in the aqueous extract while steroids were absent (Table 2).

**Table 2: Phytochemicals in aqueous leaf extract of *Ocimum gratissimum*.**

S/No	Parameter	
1	Alkaloids	+++
2	Saponins	++
3	Tannins	+
4	Phlobatannins	+
5	Anthraquinones	+
6	Steroids	-
7	Terpenoids	+
8	Flavonoids	+++
9	Cardiac glycosides	
	With steroidal ring	++
	With deoxy-sugar	++
10	Essential oil	+++

+++ = Present in high concentration

++ = Present in moderate concentration

+ = Slightly or sparingly present

- = Absent

### Result of the Effect of Aqueous Leaf Extract of *Ocimum gratissimum* on Body Weight of the Wistar Rats

**Table 3:** Effect of *Ocimum gratissimum* extracts on the body weight of the Wistar rats.

Groups	Initial Weight	Final Weight	% change in weight
<b>A (normal control)</b>	293.3 $\pm$ 19.4	306.4 $\pm$ 18.8	4.5%

<b>B (CYC control)</b>	237.0±8.4	218.9±42.8 <sup>a</sup>	-7.6%
<b>C (100mg kg<sup>-1</sup> extract)</b>	211.6±4.4	216.2±26.8 <sup>b</sup>	4.6%
<b>D (200mg kg<sup>-1</sup> extract)</b>	163.3±4.0	197.8±9.2	21.1%
<b>E (Protective I)</b>	188.1±3.9	207.1±26.1 <sup>ab</sup>	10.1%
<b>F (Protective II)</b>	168.6±6.8	183.5±18.6 <sup>ab</sup>	8.8%

<sup>a</sup>*P* < 0.05 (statistically significant compared with control), <sup>b</sup>*P* < 0.05 (statistically significant compared with myelosuppressed rats).

### Result of the Effect of *Ocimum gratissimum* on the Haematological Parameters of the Treated Rats

Bone marrow suppression occurred in group B rats that were treated with only cyclophosphamide. When compared to group A (negative control) there was no significant (*p*>0.05) increase in PCV, TWBC, RBC, and Hb in group C and D rats which received only 100 mg/kg and 200 mg/kg of the extract respectively. However, there was a significant increase (*p*<0.05) in the above-mentioned blood parameters of the group C and D rats when compared to group B which was treated with only cyclophosphamide. The increase was moderately dose-dependent.

The administration of both the extract and cyclophosphamide brought about a protective activity. This was evident in the group E and F rats which had no significant decrease (*p*>0.05) in HCT, Hb, RBC, and TWBC when compared to the control group. The group E and F rats received 100mg/kg and 200mg/kg extract respectively.

The mean differences in RBC, TWBC, HCT, Hb, and platelet of group E were slightly lower when compared to the normal control group. Meanwhile, the HCT, Hb, RBC, TWBC, and platelet amounts are significantly higher (*p*<0.05) than that of the cyclophosphamide group.

**Table 4:** Effect of aqueous extract of *Ocimum gratissimum* leaf on some hematological indices of cyclophosphamide treated rats.

Groups	PCV (%)	WBC (×10 <sup>9</sup> L <sup>-1</sup> )	RBC (×10 <sup>9</sup> L <sup>-1</sup> )	Hb (g/dL <sup>-1</sup> )
<b>A (normal control)</b>	49.0±2.65	6666.7±231.0	245.0±5.00	17.7±0.91
<b>B (CYC control)</b>	23.3±3.50	4000.0±400.0 <sup>a</sup>	216.7±15.3 <sup>a</sup>	14.7±2.88 <sup>a</sup>
<b>C (100mg kg<sup>-1</sup> OG)</b>	45.0±1.00 <sup>b</sup>	7333.3±1222.0 <sup>b</sup>	236.7±25.2 <sup>b</sup>	15.3±0.37 <sup>b</sup>
<b>D (200mg kg<sup>-1</sup> OG)</b>	51.7±6.51 <sup>b</sup>	8533.3±2023.2 <sup>b</sup>	236.7±15.3 <sup>b</sup>	17.6±2.21 <sup>b</sup>
<b>E (Protective I)</b>	35.7±6.14 <sup>b</sup>	5400.0±229.2 <sup>ab</sup>	235.0±8.67 <sup>ab</sup>	18.9±2.76 <sup>b</sup>
<b>F (Protective II)</b>	39.0±7.14	6266.7±1205.5	246.7±15.3 <sup>b</sup>	15.0±2.70

<sup>a</sup>*P* < 0.05 (statistically significant compared with control), <sup>b</sup>*P* < 0.05 (statistically significant compared with myelosuppressed rats).

**OG** = *Ocimum gratissimum*

**CYC** = Cyclophosphamide

**Table 4:** Effect of aqueous extract of *Ocimum gratissimum* leaf on some hematological indices of cyclophosphamide treated rats continued.

Groups	MCV (fl)	MCH (pg)	MCHC (g/dL <sup>-1</sup> )	PLATELETS (×10 <sup>9</sup> L <sup>-1</sup> )
<b>A (normal control)</b>	0.20±0.01	0.68±0.04	0.28±0.02	243.3±5.77
<b>B (CYC control)</b>	0.20±0.05	0.69±0.18	0.37±0.23	150.0±20.0
<b>C (100mg kg<sup>-1</sup> OG)</b>	0.19±0.02	0.65±0.08	0.26±0.01	220.0±20.0
<b>D (200mg kg<sup>-1</sup> OG)</b>	0.22±0.03	0.75±0.12	0.26±0.02	240.0±17.3

<b>E (Protective I)</b>	0.22±0.03	0.74±0.09	0.31±0.02	230.0±30.0 <sup>b</sup>
<b>F (Protective II)</b>	0.18±0.04	0.61±0.14	0.30±0.01 <sup>b</sup>	240.0±10.0 <sup>b</sup>

<sup>a</sup>*P* < 0.05 (statistically significant compared with control), <sup>b</sup>*P* < 0.05 (statistically significant compared with myelosuppressed rats).


**OG** = *Ocimum gratissimum*

**CYC** = Cyclophosphamide


UNDER PEER REVIEW

**Effect of Aqueous Extract of *Ocimum gratissimum* leaf on Histo-Architecture of Cyclophosphamide treated Wistar Rats.**

The bone marrow micrograph of the Wistar rats in each of the groups was examined and the following were observed:



**Plate 1: Photomicrograph of the bone marrow of Wistar rats in group A showing clustered normal cells (circles) with clearly visible connective tissues. [H & E. mag. 400x].**



**Plate 2: Photomicrograph of the atrophied bone marrow of Wistar rats in group B showing myelodysplastic syndrome (yellow star) together with a clear appearance of tumor cell (red star) (lymphoid progenitor cells) which usually have a yellow appearance. [H & E mag. 400x].**




Plate 3: Photomicrograph of the bone marrow of Wistar rat in group C showing normal clustered cells with the presence of megakaryocytes and hematopoietic cords. [H & E. mag 400x].






Plate 4: Photomicrograph of the bone marrow of Wistar rat in group D showing normal clustered cells with the presence of megakaryocytes and hematopoietic cords. [H& E. mag. 400x].



**Plate 5: Photomicrograph of the bone marrow of Wistar rats in group E showing the destruction of the blood-forming cells in the bone marrow. This condition can be referred to as myelodysplastic syndrome mainly of secondary origin. [H&E. mag. 400x].**



**Plate 6: Photomicrograph of the bone marrow of group F rats showing apparent normocellular (circles) bone marrow with slight loss of some hematopoietic cells which appears quite necrotized (star). [H& E. mag. 400x].**

## DISCUSSION

In this study, the phytochemical analysis of the aqueous leaf extract of *Ocimum gratissimum* revealed the presence of tannins, steroids, terpenoids, flavonoids, and cardiac glycosides. Anthraquinones were also detected in the aqueous leaf extract while steroids were absent (Table 2). A similar result to this was observed by Okoye<sup>[13]</sup> and Afolabi<sup>[14]</sup>.

de Vasconcelos<sup>[15]</sup> reported that the phytochemical constituents of a plant are largely affected by the type of soil and environment in which the plant is grown, this could therefore be the reason why in contrast to the result of this study, Akinmoladun<sup>[16]</sup> reported the presence of steroids.

The high content of flavonoids and alkaloids in the leaf extract could be the reason for the protective activity observed in the leaf extract. Alkaloids are known to play some metabolic roles and control development in a living system. They also have a protective role on animals<sup>[8]</sup>. Flavonoids are free radical scavengers, super antioxidants, and potent water soluble which prevent oxidative cell damage and have strong anticancer activity<sup>[17]</sup>. The presence of alkaloids and flavonoids in this study therefore could be the reason for the protective actions of *Ocimum gratissimum*. Also, flavonoids could have been the reason for the reduction in the oxidative cell damage reported in this study. This is in agreement with the studies carried out by Thabrew<sup>[18]</sup>, Halliwell, and Gutteridge<sup>[19]</sup>.

Histological results in the Normal control group showed normal cells with clearly visible connective tissues and it is therefore in agreement with the study made by Travlos<sup>[20]</sup>. The atrophied bone marrow seen in the rats induced with bone marrow suppression is indicative of myelodysplastic syndrome. This is a result of the myelotoxicity activity of cyclophosphamide and its ability to cause bone marrow lesions, as also recorded by Travlos<sup>[20]</sup>. The normal clustered cells in the rats that were treated with only extracts imply that aqueous extract of *Ocimum gratissimum* does not have any toxic effect on the bone marrow cells of Wistar rats.

Administering the rats with both the leaf extract and cyclophosphamide brought about a protective effect against the cyclophosphamide-induced myelotoxicity in the Wistar rats. This was evident in the treated rats which showed reduced cell death in their histo-architecture. This might be a result of the presence of alkaloids and flavonoids in the extract. According to Onwueyiagba<sup>[17]</sup>, McMahon<sup>[21]</sup>, and Okoye<sup>[13]</sup>, flavonoids and alkaloids are free radical scavengers, super antioxidants, and potent water-soluble agents which prevent oxidative cell damage and have strong anticancer activity.

Chemotherapy-induced anemia is one of the most common side effects experienced by cancer patients, occurring in approximately 70-90% of those undergoing treatment for the disease<sup>[22]</sup>. This is because chemotherapeutic drugs kill rapidly dividing cells in the body including cancer cells and normal cells which include red blood cells at the same time suppress bone marrow's ability to produce new ones, resulting in a decrease in blood Hb level<sup>[11]</sup>. In this study, it was observed that gavage of the animals with *Ocimum gratissimum* extract enhanced the Hb level, platelet count, TWBC, PCV, and RBC count which was depleted by cyclophosphamide treatment in the cyclophosphamide control group. The effect of this plant extract on the hematological parameters may be due to free radical scavenging activity or interference with the formation of the active metabolite of cyclophosphamide through the inhibition of the cytochrome p-450 enzyme system<sup>[11]</sup>.

## CONCLUSION

Aqueous extracts of *Ocimum gratissimum* leaf possess a protective effect against cyclophosphamide-induced bone marrow toxicity and consequently justify the use of this extract as a tonic in traditional medicine.

## CONSENT

It is not applicable.

## REFERENCES

1. Neboh, E. E., & Ufelle, S. A. (2015). Myeloprotective activity of crude methanolic leaf extract of *Cassia occidentalis* in cyclophosphamide-induced bone marrow suppression in Wistar rats. *Advanced biomedical research*, 4, 5. <https://doi.org/10.4103/2277-9175.148285>.
2. Atal, C. K., Zutshi, U., & Rao, P. G. (1981). Scientific evidence on the role of Ayurvedic herbals on bioavailability of drugs. *Journal of ethnopharmacology*, 4(2), 229–232.
3. Siddiqui, M. S., & Amy, G. L. (1993). Factors affecting DBP formation during ozone–bromide reactions. *Journal-American Water Works Association*, 85(1), 63-72.
4. Okochi, V. I., Kazeem, A. A., Gbenle, G. O., Fagbenro-Beyoku, A. F., Dare, A., & Arukwu, U. (2001). Effect of *Allium sativum* (Garlic) water extract on *Trypanosoma brucei* infection in laboratory rat. *Biokemistri*, 11, 9-15.
5. De Smet, P. A. G. M. (2002). Drug therapy: Herbal remedies. *The New England Journal of Medicine*, 347(25), 2046–2056. <https://doi.org/10.1056/NEJMra020398>
6. Sofowora A. (1993). Medicinal Plants and Traditional Medicinal in Africa. 2nd Ed. *Sunshine House, Ibadan, Nigeria: Spectrum Books Ltd. Screening Plants for Bioactive Agents*; pp. 134–156.
7. Grayer, R. J., & Kite, A. C. (2000). Characteristics of flavonoids from *Ocimum gratissimum* (pp. 257-267). *Richmond, U. K.*
8. Edeoga, H. O., & Eriata, D. O. (2001). Alkaloids, Tannins and Saponins contents of some Nigeria Medicinal Plants. *Journal of Medicinal Aromatic and Plant Science*, 33, 344-349.
9. Anyanwu, M. & Okoye, Rosemary. (2017). Antimicrobial activity of Nigerian medicinal plants. *Journal of Intercultural Ethnopharmacology*. 6. 240-259. [10.5455/jice.20170106073231](https://doi.org/10.5455/jice.20170106073231).
10. Randall, R.P. (1973): Mode of action of Herbicides, Willey, New York p.220.
11. Okwuosa, C. N., Achukwu, P. U. O., Azubike, N. C., & Abah, A. I. E. (2012). Protective Effect of the Leaf Extracts of *Combretum racemosum* and *P.Beauv (combretaceae)* on Cyclophosphamide Induced Pancytopenia and Liver Injury in Male Rats. *Research Journal of Pharmacology*; 6(2): 30-34.
12. Drury, R.A.B., Wallington, E.A., & Cameron, R. (1967). Carleton's Histological Techniques: 4th ed., Oxford University Press NY. USA, pp. 279-280.
13. Okoye, N. N., Ajaghaku, D. L., Okeke, H. N., Ilodigwe, E. E., Nworu, C. S., & Okoye, F. B. (2014). beta-Amyrin and alpha-amyrin acetate isolated from the stem bark of *Alstonia boonei* display profound anti-inflammatory activity. *Pharmaceutical biology*, 52(11), 1478–1486. <https://doi.org/10.3109/13880209.2014.898078>
14. Afolabi, C., Akinmoladun, E. O., Ibukun, E. A., Obuotor, E. M., & Farombi, E. O (2007). Phytochemical constituent and antioxidant activity of extract from the leaves of *O. gratissimum*, *Scientific Research and Essay*. 2 (5); 163-166
15. de Vasconcelos Silva M, Craveiro A, Abreu Matos F, Machado M, & Alencar J (1999) Chemical variation during the daytime of constituents of the essential oil of *Ocimum gratissimum* leaves. *Fitoterapia* 70:32–34
16. Akinmoladun, C., A., Ibukun, E. O., Afor, E., Obuotor, E. M., & Farombi, E. O. (2007). Phytochemical constituent and antioxidant activity of extract from the leaves of *Ocimum gratissimum*. *Scientific Research and Essay*, 2(5): 163-166.
17. Onwueyiagba, E. O. (2001). Evaluation of the seed of *M. pruriens* as a feed ingredient in poultry and pig diets. Ph.D. thesis in Animal Nutrition, p.44.

18. Thabrew, M. I., Hughes, R. D., & McFarlane, I. G. (1998). Antioxidant activity of *Osbeckia aspera*. *Phytotherapy Research: An International Journal Devoted to Pharmacological and Toxicological Evaluation of Natural Product Derivatives*, 12(4), 288-290.
19. Halliwell, B., Gutteridge, J. M., & Cross, C. E. (1992). Free radicals, antioxidants, and human disease: where are we now? *The Journal of laboratory and clinical medicine*, 119(6), 598–620.
20. Travlos, G. S. (2006). Normal structure, function, and histology of the bone marrow. *Toxicologic Pathology*; 34(5), 548–565.
21. McMahon, M., Itoh, K., Yamamoto, M., Chanas, S. A., Henderson, C. J., McLellan, L. I., Wolf, C. R., Cavin, C., & Hayes, J. D. (2001). The Cap'n'Collar basic leucine zipper transcription factor Nrf2 (NF-E2 p45-related factor 2) controls both constitutive and inducible expression of intestinal detoxification and glutathione biosynthetic enzymes. *Cancer Research*, 61(8), 3299–3307.
22. Groopman, J. E., & Itri, L. M. (1999). Chemotherapy-induced anemia in adults: incidence and treatment. *Journal of the National Cancer Institute*, 91(19), 1616–1634. <https://doi.org/10.1093/jnci/91.19.1616>
23. Abbah, J., Amos, S., Chindoc, B., Ngazalc, I., Vongtaue, H. O., Adzue, B., Faridad, I., odutolad, A. A., Wambebec, C., & Gamaniel, K. S. (2009). Pharmacological evidence favouring the use of *Nauclea Latifolia* in malaria ethnopharmacy; effect against nociception, inflammation, and Pyrexia in rat and mice. *Journal of Ethnopharmacology*; 127: 85-90.
24. Atuboyedia, W. O., Jonah, S. A., and Chinagoro, T. O. E. (2010). Anti-fertility effects of aqueous crude extract of *Ocimum gratissimum* L. leaves in male mice. *Journal of Medicinal Plants Research*; 4(9): 809-816. DOI: 10.5897/JMPR10.099.
25. Bonadonna G., Valagussa P., Moliterni A., Zambetti M., & Brambilla C. (1995). Adjuvant cyclophosphamide, methotrexate, and fluorouracil in node-positive breast cancer – the result of 20 years of follow-up. *New England Journal of Medicine* 332 (14), 901 – 906.
26. Hu, W., Sung, T., Jessen, B. A., Thibault, S., Finkelstein, M. B., Khan, N. K., & Sacca, A. I. (2016). Mechanistic Investigation of Bone Marrow Suppression Associated With Palbociclib and its Differentiation from Cytotoxic Chemotherapies. *Clinical Cancer Research*; 22(8): 2000-8. Doi: 10.1158/1078-0432.CCR-15-1421.
27. Ivy, A. W. H., Han, C. T., Wai, H. N., Yuan, L. T., Chang, M. G., Kam, M. H., & Paula, Y. P. L. (2013). Human Bone Marrow-Derived Mesenchymal Stem Cells Suppress Human Glioma Growth Through Inhibition of Angiogenesis. *Stem Cells Journals*; 31(1): 146-155. DOI: 10.1002/stem.1247.
28. Johanna, K. M., Maciej, K., Keith, L. K., Harry, D. B., & Masoud, H. M. (2010). GM-CSF is one of the main breast tumor-derived soluble factors involved in the differentiation of CD11b-Gr1-bone marrow progenitor cells into myeloid-derived suppressor cells. *Breast Cancer Research and Treatment*; 123: 39. Doi:10.1007/s10549-009-0622-8.
29. Joshi, R. K. (2013). Chemical composition, In vitro antimicrobial and antioxidant activities of the essential oils of *Ocimum gratissimum*, *O. Sanctum*, and their major constituents. *Indian Journal of Pharmaceutical Sciences*; 75(4): 457-462. DOI: 10.4103/0250-474X.119834.
30. Kosowski, A. (2014). Whole Herbs vs Standardized Herbal Extracts: Which are better? Retrieved from: <http://www.nowfoods.com/Quality/Do-supplements-Work/M043723.htm>.
31. Lorke, D. A. (1983). A new approach to practical acute toxicity testing. *Archives in Toxicology*; 53: 275-289.

32. Luciane, A. G., Aline de Moraes, P., & Mirtes, C. (2010). Biological effects of *Ocimum gratissimum* L. are due to synergic action among multiple compounds present in essential oil. *Journal of Natural Medicines*; 64(4): 436-441. DOI: 10.1007/s11418-010-0429-2.
33. Mbakwem-Aniebo, C., Onianwa, O., & Okonko, I. O. (2012). Effects of *Ocimum gratissimum* Leaves on Common Dermatophytes and Causative Agent of Pityriasis Versicolor in Rivers State, Nigeria. *Journal of Microbiology Research*; 2(4): 108-113. DOI: 10.5923/j.microbiology.20120204.08.
34. Momoh M. A., Adikwu M. U., & Oyi A. R. (2010). *Vernonia amygdalina* extract and CD4<sup>+</sup> cell counts. An immune study. *Global journal of biotechnology and biochemistry*; 5(2): 9296.
35. Nakamura, C. V., Ueda-Nakamura, T., Bando, E., Melo, A. F., Cortez, D. A., & Dias Filho, B. P. (1999). Antibacterial activity of *Ocimum gratissimum* L. essential oil. *Memorias do Instituto Oswaldo Cruz*, 94(5), 675–678. <https://doi.org/10.1590/S0074-02761999000500022>
36. Nwosu, M. O., & Okafor, J. J. (1995). Preliminary studies of the antifungal activities of some medicinal plants against *Basidiobolus* and some pathogenic fungi. *Mycoses*;38: 191-195.
37. Ofem O. E, Ani, E. J., &Eno, A. E. (2012). Effect of aqueous leaf extracts of *Ocimum gratissimum* on hematological parameters in rats. *International Journal of Applied and Basic Medical Research*; 2(1): 38-42. DOI: 10.4103/2229-516X.96807.
38. Oguanobi, N. I., Chioli, P. C., & Ghasi, S. I. (2012). Effects of aqueous leaf extract of *Ocimum gratissimum* on oral glucose tolerance test in type-2 model diabetic rats. *African Journal of Pharmacy and Pharmacology*; 6(9): 630-635. DOI: 10.5897/AJPP11.811.
39. Okoli, C. O., Ezike, A. C., Agwagah, O. C., & Akah, P. A. (2010). Anticonvulsant and anxiolytic evaluation of leaf extracts of *Ocimum gratissimum*, a culinary herb. *Pharmacognosy Research*, 2(1), 36–40. <https://doi.org/10.4103/0974-8490.60580>.
40. Onajobi F. D. (1986). Smooth muscle contracting lipidic soluble principle in chromatographic functions of *Ocimum gratissimum*. *Journal of Ethnopharmacology*. 18: 3-11.
41. Oparaocha, E. T., Iwu, I., &Ahanaku, J. E. (2010). Preliminary study on mosquito repellent and mosquitocidal activities of *Ocimum gratissimum* (L.) grown in eastern Nigeria. *Journal of Vector-Borne Diseases*; 47(1): 45-50.
42. Saha, S., Tarak, N. D., Sengupta, C., &Ghosh, P. (2013). Biological Activities of Essential Oils and Methanol Extracts of Five *Ocimum* species against Pathogenic Bacteria. *Czech Journal of Food Science*; 31(2): 194-202.
43. Subbaram, N. R. (1997). Medicinal plants and intellectual proper rights. International workshop on medicinal plants. Their bioavailability screening and evaluation (lecturer 10) Lucknow, India.
44. Tilburt, J. C., and Kaptchuk, T. J. (2008). Herbal medicine research and global health: an ethical analysis. *Bulletin of the World Health Organization*; 86: 594- 599.
45. Vashishtha, V. M., John, T. J., & Kumar, A. (2009). Clinical & pathological features of acute toxicity due to *Cassia occidentalis* in vertebrates. *The Indian journal of medical research*, 130(1), 23–30.
46. Yung-Wei, C., Hung-Jen, L., Hsin-Yu, H., Pei-Yu, C., Jin-Ming, H., Pei-Yun, H., Shyh-Jer, H., Jer-Yuh, L., &Te-Jen, L. (2013). The antioxidant and cytoprotective activity of *Ocimum gratissimum* extracts against hydrogen peroxide-induced toxicity in human HepG2 cells. *Journal of Food and Drug Analysis*; 21(3): 253-260. DOI:10.1016/j.fda.2013.002.