

Original Research Article

Weight Change Before and After 30 Days of Intermittent Fasting Among University Students in Pakistan

ABSTRACT

INTRODUCTION: Fasting is a practice of abstaining from certain or all kinds of food and beverages depending on the cultural and religious values of the individual. The objective of this study was to observe the effects of intermittent fasting on the weight of Muslim university students in Karachi, Pakistan.

METHODS: A longitudinal study was undertaken at the public sector university in Karachi, Sindh, Pakistan between April 2022 and August 2022. Using a predefined questionnaire at the beginning and end of **one month of** intermittent fasting, **personal data and weight changes were** recorded for a total of 95 participants.

RESULTS: The **sample was based on** primary medical students with a predominance of 73.7% females (70). Most of them (67.4%) were in the age range between 20 and 29 years. About 44 (62.90%) were under the normal BMI of 18.5. When comparing the participants' weight, the mean weight pre-fasting was 58.66 ± 12.12 kg, and the weight after one month of fasting was 58.76 ± 12.67 kg showing a difference of 0.10 kg and a non-significant p-value of 0.73. 17.9% of the subjects had gained over 1 kilogram, 23.2% had lost over 1 kilogram, and 22.1% maintained their weight, after fasting.

CONCLUSION: The study indicated a weight change in most individuals enrolled. Although the weight change was not significant, an increase was seen among most of the individuals. Factors such as decreased physical activity due to COVID 19 quarantine practices, consumption of high-calorie dense foods, and reduced metabolism during fasting may contribute to the results.

Keywords: Diet, Fasting, Intermittent Fasting, Metabolism, Weight Loss, Young Adults

INTRODUCTION:

Fasting is achieved by minimal or no consumption of food or beverages ranging from a period of a few hours or multiple weeks [1]. Religious fasting can be observed in Christianity, Buddhism, Jainism, Taoism, and Hinduism [2]. During fasting, an individual is required to abstain from eating for a certain number of hours. If it is performed during a particular time of the day, it is called intermittent fasting. The length of the fast varies depending upon the type of fasting. A religious fast can last up to 18 hours in tropical countries [3-5].

Over recent years intermittent fasting has gained popularity as a diet trend for weight loss and other benefits such as improving sleep, gut microbiome, and metabolism [6]. Intermittent fasting raises the question of how effective it is in losing weight [7]. Many studies have shown fluctuating data on the weight change during the month of Ramadan which is one of the models for intermittent fasting. Previously, a study conducted in a mosque in London comparing the weight before the beginning of the holy month with the weight of participants at the end of the month showed a reduction of less than one kilogram (kg) of weight for 46% of the participants. Also, those who continuously fasted every day throughout the whole month could lose more weight. However, all participants regained their weight one month later [8].

Another study conducted about the differences in the metabolism when fasting in university students concluded that participants had a decrease in glucose and BMI but not a significant difference in cholesterol [9]. Similarly, a study in Jordan monitoring students' lipid and glucose levels after fasting found a weight decrease in males of 2.2 kg and females of 1.2 kgs with the high-density-lipoprotein increase and low-density-lipoprotein decline throughout Ramadan [10]. A non-systematic review tabulated the nutrient intake and weight changes from available studies stating no robust data on weight change [11]. As of now, no study has been conducted in Pakistan assessing the weight change of the youth because of fasting during the coronavirus pandemic.

The aim of this study was to assess and monitor the effects of fasting during the 30 days of intermittent fasting among university students from Karachi, Pakistan.

METHODOLOGY:

A longitudinal study was undertaken at the public sector university in Karachi, Sindh, Pakistan between April 2022 and August 2022. All authors hereby affirm that all procedures followed the ethical guidelines outlined in the 1964 Declaration of Helsinki after being reviewed and approved by the institutional ethics committee (Reference # S212).

The sampling technique used was non-probable and convenient. The sample size was calculated through OpenEpi Version 3. The population size was 1000000, the anticipated frequency was 7.4, and the confidence interval was 95%. The inclusion criteria consisted of participants aged 18 years or older and university-going students who were either normal weight, overweight, or obese. All those who were underweight, who failed to fast for more than 15 days or did not fill the post-intermittent fasting questionnaire were excluded from the study. Informed written consent was documented and requested from each patient prior to their inclusion in the study.

The data collection procedure was carried out as follows. A predefined proforma was distributed among the students using the inclusion criteria mentioned above. All participants were requested to give informed consent and fill in their socio-demographic information, including age, gender, weight, height, university, comorbidities, and lifestyle. Body mass index was reported by dividing the square of height in meters with weight of the individual in kilograms according to the following categories:

If a patient's BMI was $<18.5 \text{ kg/m}^2$, he/she was considered underweight. If a patient's BMI was between 18.5 to 24.9, he was considered normal weight. If a patient's BMI was between 25 to 29.9, he was considered overweight. If a patient's BMI was > 30 , he was considered obese.

The students who agreed to observe intermittent fasting during the study were guided by the authors about the schedule of intermittent fasting. All individuals were requested to skip breakfast then have the first meal of the day at 12 pm then have the last meal at 8 pm. In total 16 hours of fasting was observed by the participants. The participants were requested to maintain a diary where any deviations from the schedule must be observed. If individuals could not keep fasting due to weakness, they were requested to **have** breakfast immediately and consume food to prevent hypoglycemia. Water, coffee, tea and other non-caloric beverages were allowed during the intermittent fasting.

After 30 days, all individuals were requested to visit the university health center and they were **weighed** with their shoes off, using a weighing machine. The weight change was recorded and body mass index was documented. Anonymity was assured as the results were analyzed confidentially. Agreement to be a part of this study was mentioned on the questionnaire, and participants were allowed to withdraw at any time by filling out the forms.

All data were collected by the researchers. Further statistical analysis was done through the Statistical Package for the Social Sciences (SPSS) software version 24, observing the correlation between fasting on weight and BMI. Other variables

included in the demographics were also inspected for the multifactorial variance seen among the weight of the individuals.

RESULTS:

The results discussed below are of only those 95 participants who had completed both questionnaires provided. Most of the participants were students from Dow University of Health Sciences, 69(72.6%) (Figure 1). The gender distribution was 70(73.7%) female and 25(26.3%) male (Table 1). The mean age was 18 ± 3.31 , the bulk being between 20 to 29 years. The study consisted of largely unmarried individuals being 85(89.5%), whereas married individuals were 10(10.5%). The vast majority of the participants in the study were in the range of a normal BMI between 18.5 - 24.9 %, with females being 44(62.90%), and males 16(64.00%) (Table 2).

When comparing the baseline weight of the 95 students with the weight after intermittent fasting, the mean weight difference in kilograms was 0.10 kg and a non-significant value p-value of 0.73. BMI baseline was also compared to the increased BMI after the fasting month yielding a mean difference of 0.06 kg/m² with a p-value of 0.59, also considered non-significant (Table 3).

The mean weight change was calculated as an average gain of 0.10 ± 2.63 kgs (Table 4). Most of the participants, 23(24.2%), gained between 0.5 to 1 kg, whereas many, 22(23.2%) lost over 1 kg, About 21(22.1%) maintained their weight with a marginal gain or decrease of 0 - 0.5 kgs, and few individuals, 17(17.9%), gained over 1kg. The least number of participants, 12(12.6%), lost over 0.5 - 1 kgs. Only one obese female participant with a BMI of >30.0 had lost enough weight after the fasting month to be categorized in the lower overweight BMI category of 25.0 - 29.9 after the 30 days had ended (Table 2).

Figure 1: University Distribution

Distribution of Participants According To Universities




Table 1. Demographics

Demographics	N(%)
Gender	
Female	70 (73.7)
Male	25 (26.3)
Marital Status	
Married	10 (10.5)
Unmarried	85 (89.5)
Age	18±3.31
<20 yrs	30 (31.6)
20-29 yrs	64 (67.4)
30-39yrs	1 (1.1)

Table 2: Chi-square evaluation of gender and age with BMI before and after intermittent fasting.

	BMI pre-fasting	Significance

	<18.5 Underweight	18.5- 24.9 Healthy	25.0- 29.9 Overweight	>30.0 Obese	
Gender					0.118
Female	17(24.3%)	44(62.9%)	6(8.6%)	3(4.3%)	
Male	3(12%)	16(64%)	6(24%)	0(0%)	
	BMI post-fasting				
Gender					0.197
Female	17(24.3%)	44(62.9%)	7(10%)	2(2.9%)	
Male	3(12%)	16(64%)	6(24%)	0(0%)	
	BMI pre-fasting				
	<18.5 Underweight	18.5- 24.9 Healthy	25.0- 29.9 Overweight	>30.0 Obese	
Age					<0.0001
>20 yrs	10(33.3%)	16(53.3%)	3(10.%)	1(3.3%)	
20-29 yrs	10(15.6%)	44(68.8%)	9(14.1%)	1(1.6%)	
30-39yrs	0(0%)	0(0%)	0(0%)	1(100%)	
	BMI post-fasting				
Age					<0.0001
>20 yrs	10(33.3%)	16(53.3%)	4(13.3%)	0(0%)	
20-29 yrs	10(15.6%)	44(68.8%)	9(14.1%)	1(1.6%)	
30-39yrs	0(0%)	0(0%)	0(0%)	1(100%)	

Table 3: Comparison of Weight Change and BMI Change Before and After Fasting

Weight pre-fasting	Weight post-fasting	Difference	P-Value
58.66±12.12	58.76±12.67	0.1	0.73

BMI pre-fasting	BMI post-fasting	Difference	P-Value
21.52±4.043	21.58±4.44	0.06	0.59

Table 4: Association of BMI and Gender Before and After intermittent fasting

	BMI for Male	BMI for Female	p-value
Pre-intermittent fasting	22.47	21.18	0.17
Post-intermittent fasting	22.62	21.2	0.16

DISCUSSION

One of the notable physical changes due to fasting is weight difference among Muslim individuals from Karachi. This study has evaluated weight variation and BMI patterns among medical students as gain, loss, or no change. It has been founded that weight was gained among most of the samples. However, the difference is statistically insignificant, both for weight variables as well as BMI. While there have been factors such as lack of physical activity and changes in mental health etc., that may conflict with the results; the subsequent literature studies support women's weight al. conducted an observational study of 34 volunteers where weight change ($p=0.894$) and BMI ($p=0.764$) proved insignificant [12].

Another study done in Turkey investigated metabolic, biochemical, and psychiatric alterations among 39 subjects. It was stated that BMI change was from 64.2 ± 12.2 kg/m^2 to 62.8 ± 11.8 kg/m^2 which too is seen as insignificant [13]. A meta-analysis of 21 studies by Kul et al. observed no significant changes in women's weight ($p=0.620$); however, weight reduction was seen as slightly significant in men ($p=0.001$) [14]. Harder-Lauridsen et al. conducted a nonrandomized, crossover intervention study where a decrease of 1.7 kg of total body mass was seen among ten healthy lean men. This change, too, was seen as insignificant ($p=0.05$) [15].

There have been recent research papers that state significant differences in weight and BMI among healthy populations. In contrast to our study, Fernando et al. highlighted in their meta-analysis of 70 publications that participants with average weight had a significant decrease of 1.34 kg, which remained after follow-up ($p < 0.001$) [16]. Syam et al. reported that their study involving 43 participants who fasted for 28 days revealed a significant change in baseline weight from 59.82 ± 11.252 to 58.95 ± 11.201 kgs ($p < 0.001$) [17]. Sadeghirad et al. stated in their meta-analysis that weight reduction was seen majorly among 35 studies where African populations had the most significant decrease of an average of 1.13 kg ($P = 0.001$). This study also mentioned that most weight losses was recovered after two weeks as dietary

patterns and energy utilization resumed before intermittent fasting [18]. Compared to the contrasting studies done in recent years, our study was carried out during the onset of COVID-19 when quarantine was strictly practiced. Lack of physical activity, energy expenditure, altered mental status, limited food choices, etc., may be significant reasons for the difference in findings. Studies have also reported that weight reduction is seen more in men than among women. In most Middle Eastern and South Asian countries, females have comparatively lower metabolic rates and physical activity [19].

Further research investigation on the commonly practiced Intermittent fasting by young adults and its effects on body weight and health is recommended. Nocturnal activity is high during fasting and is now increasingly practiced. This may cause changes in body metabolism. A study states that sleep quality significantly improved after fasting due to resumed dietary patterns [20]. The participants in this study had no severe comorbidities, and their food choices were primarily high in carbohydrates and fat. This can be an alerting topic to discuss if consuming lipid-rich food after hours of fasting may lead to cardiovascular diseases, hypertension, and obesity. Abstaining from calorie-rich diets has improved blood lipid profile, so this can hypothetically prove that diet has a significant effect on improving or worsening health during fasting [21].

The limitations of the study are acknowledged. Only students were included in the study from which a small minority was overweight. Also, the study was in Karachi, Pakistan, and no participants from different ethnicities and geographical locations were present. The research was conducted during the COVID-19 pandemic and may not represent normal circumstances of individuals due to quarantine practices. Lastly, data collection was through self-filled questionnaires, which could lead to the risk of bias.

CONCLUSION:

The study aimed to observe the weight changes caused by ~~fasting in~~ intermittent fasting. After comparing the participants' weight after intermittent fasting from their baseline weight, most had gained weight, although the results were not statistically significant. The study sheds light on standard intermittent fasting methods used for weight loss and the impact of fasting without food and water for long intervals on the weight of individuals. It also implies the need for future research to be conducted on intermittent fasting, focusing more on obese individuals and including participants from all ages, professional backgrounds, and diverse ethnicities.

REFERENCES:

1. Longo VD, Mattson MP. Fasting: Molecular Mechanisms and Clinical Applications. *Cell Metab.* 2014;19(2):181–192. <https://doi.org/10.1016/j.cmet.2013.12.008>
2. Venegas-Borsellino C, Sonikpreet, Martindale RG. From Religion to Secularism: the Benefits of Fasting. *Curr Nutr Rep.* 2018;7(3):131–138. <https://doi.org/10.1007/s13668-018-0233-2>
3. Azizi F. **Medical aspects of islamic fasting**. *Iran MJIRI.* 1996;10(3):241–246.
4. Kettani H. World Muslim Population:1950–2020. *Int J Environ Sci Dev.* 2010;1(2):1–42.
5. 10 countries with the largest Muslim populations, 2015 and 2060 | Pew Research Center. Accessed 2022 Apr 2. Available from: https://www.pewresearch.org/fact-tank/2019/04/01/the-countries-with-the-10-largest-christian-populations-and-the-10-largest-muslim-populations/ft_19-03-29_muslimchristianpopulations_muslim/
6. Patterson RE, Sears DD. Metabolic Effects of Intermittent Fasting. *Ann Rev Nutr.* 2017. [at https://doi.org/10.1146/annurev-nutr-071816-064634](https://doi.org/10.1146/annurev-nutr-071816-064634)
7. Lessan N, Ali T. Energy Metabolism and Intermittent Fasting: The Ramadan Perspective. *Nutr* 2019;11:1192. <https://doi.org/10.3390/nu11051192>
8. Hajek P, Myers K, Dhanji A-R, West O, McRobbie H. Weight change during and after Ramadan fasting. *J Pub Health (Bangkok).* 2012;34(3):377–381.
9. Ziaee V, Razaee M, Ahmadinejad Z, Shaikh H, Yousefi R, Yarmohammadi L, et al. The changes of metabolic profile and weight during Ramadan fasting. *Singapore Med J.* 2006;47(5):409.
10. Mansi KMS. Study the effects of Ramadan fasting on the serum glucose and lipid profile among healthy Jordanian students. *Am J Appl Sci.* 2007;4(8):565–569.
11. Trepanowski JF, Bloomer RJ. The impact of religious fasting on human health. *Nutr J.* 2010;9(1):1–9.
12. Yucel A, Degirmenci B, Acar M, Albayrak R, Haktanir A. The Effect of Fasting Month of Ramadan on the Abdominal Fat Distribution: Assessment by Computed Tomography. *Tohoku J Exp Med.* 2004;204(3):179–187.
13. Furuncuoglu Y, Ender K, Aras Ş, Arif Y. Metabolic, Biochemical and Psychiatric Alterations in Healthy Subjects During Ramadan. *Pakistan J Nutr.* 2007;6.
14. Kul S, Savaş E, Öztürk ZA, Karadağ G. Does Ramadan Fasting Alter Body Weight and Blood Lipids and Fasting Blood Glucose in a Healthy Population? A Meta-analysis. *J Relig Health.* 2014;53(3):929–942. Available from: <https://doi.org/10.1007/s10943-013-9687-0>
15. Harder-Lauridsen NM, Rosenberg A, Benatti FB, Damm JA, Thomsen C, Mortensen EL, et al. Ramadan model of intermittent fasting for 28 d had no major effect on body composition, glucose metabolism, or cognitive functions in healthy lean men. *Nutrition.* 2017;37:92–103. <https://doi.org/10.1016/j.nut.2016.12.015>

16. Fernando HA, Zibellini J, Harris RA, Seimon R V, Sainsbury A. Effect of Ramadan Fasting on Weight and Body Composition in Healthy Non-Athlete Adults: A Systematic Review and Meta-Analysis. *Nutrients*. 2019;11(1):1-11. doi:10.3390/nut11010011.
17. FahrialSyam A, SuryaniSobur C, Abdullah M, Makmun D. Ramadan Fasting Decreases Body Fat but Not Protein Mass. *Int J Endocrinol Metab*. 2016;14(1):e29687–e29687. doi:10.1515/ijem-2016-0087.
18. Sadeghirad B, Motaghipisheh S, Kolahdooz F, Zahedi MJ, Haghdoost AA. Islamic fasting and weight loss: a systematic review and meta-analysis. *Public Health Nutr*. 2014;17(2):396–406. doi:10.1017/S1368857513001808.
19. Norouzy A, Salehi M, Philippou E, Arabi H, Shiva F, Mehrnoosh S, et al. Effect of fasting in Ramadan on body composition and nutritional intake: A prospective study. *J Hum Nutr Diet*. 2013;26(suppl.1):97–104. doi:10.1111/jhn.12001.
20. Hsouna H, Abdessalem R, Boukhris O, Trabelsi K, Chtourou L, Tahri N, et al. Short-term maximal performance, alertness, dietary intake, sleep pattern and mood states of physically active young men before, during and after Ramadan observance. *PLoS One*. 2019;14(6):e0217851. doi:10.1371/journal.pone.0217851.
21. Mattson MP, Longo VD, Harvie M. Impact of intermittent fasting on health and disease processes. *Ageing Res Rev*. 2017;39:46–58. doi:10.1016/j.arr.2016.10.005.