

1 **Enhancing common bean tolerance to short-term droughts at the reproductive stage using a soil**
2 **fertility management approach**
3

4 **Authors' contributions**
5

6 All authors contributed to the conception and implementation of this research. Research data collection
7 and analyses were performed by author MBB. The first draft of the manuscript was prepared by author
8 MBB and all authors reviewed previous versions of the manuscript. All authors read and approved the
9 final manuscript.

10 **Abstract**
11

12 **Aims:** This study was conducted to enhance the tolerance of common beans to drought events occurring
13 at the reproductive stage, from a soil improvement perspective.

14 **Study design:** Split plot completely randomized design was used.

15 **Place and duration of study:** Study was conducted in a screen-house at the Legumes and Oil Seeds
16 Division of CSIR-Crops Research Institute, Ghana, from September 2021 to January 2022.

17 **Methodology:** Municipal Solid Waste Compost and inorganic fertilizer combinations were applied to
18 common beans in a pot experiment. They included control, full rate compost (FRAC), full rate fertilizer
19 (NPK 5:30:30 kg/ha) (FRG), FRG + half rate compost (HRAC) and FRG + FRAC. All soils were
20 maintained at 80% field capacity (FC) from the start of the experiment. At flowering, two groups of plants
21 were water stressed till 40 and 16% FC and returned to 80% FC till physiological maturity, while one group
22 maintained 80% FC throughout study. Forty-five soil samples each and plant data were collected at 3, 7
23 and 10 weeks after planting. Samples were analyzed for soil organic matter (SOM) and water retention,
24 soil nutrients, crop growth, yield and nutrient uptake. Water and nitrogen use efficiencies (W/NUE) were
25 calculated after harvest.

26 **Results:** During the growing period, highest soil moisture ($6-9 \text{ cm}^3/\text{cm}^3$) was retained by FRG and
27 FRG+HRAC, FRG+FRAC; 20-38% more than FRAC and control but was not influenced by SOM. While
28 FRG influenced the highest yield and WUE, combining it with compost rates reduced yield by 56-84% and
29 WUE by 55-64%. WUE correlated positively with NUE.

30 **Conclusion:** Antagonistic effect observed with integrating compost with FRG is likely because compost
31 was not properly cured and immobilized soil nitrogen. Farmers can mitigate short-term drought effects on
32 common beans with adequate nutrient supply through fertilizer application; however, fertilizer should only
33 be integrated with compost after compost quality analysis.

34 **Keywords:** compost, mineral fertilizer, water stress, soil organic matter, soil water retention, water use
35 efficiency, climate change

36 1. INTRODUCTION

37 Legumes account for 27% of global primary crop production and 33% of global protein requirement [1].
38 They are major cash crops for more than 700 million smallholder farmers in developing countries and can
39 be grown in a variety of climates and soil types[1].They fix atmospheric nitrogen in the soil and may
40 reduce the required amounts of chemical nitrogen fertilizers needed per application. Hence they are one
41 of the most promising crops to promote climate smart agriculture [2].

42 Common beans (*Phaseolus vulgaris* L.), the most important food legume for direct
43 consumption,contribute about 8.8% to the global annual total legume value of 31 billion USD[3]. Though
44 an important legume, about 60% of common bean production occurs under short-term or terminal drought
45 stresses [4]. In Ghana, legumes (common bean included) are widely cultivated in the
46 Savannahagroecological zonesof the country where short and long term droughts are common
47 occurrences [5, 6].Drought stress is a major constraint to common bean production in Ghana and many
48 other countries and results in about 10% to 100% yield losses globally [7].A70% reduction in common
49 bean yield due to drought stress in Colombia was observed by [8].An80% decrease in common bean
50 seed yield at very severe drought (drought intensity index of 0.8) was also reported by [9].As a result,
51 drought coping mechanisms have become key traits for common beangermplasm selectionand for
52 improving productivity of the crop [10]. Plants may use various mechanisms to cope with drought stress.
53 These mechanisms may be grouped into drought tolerance, drought avoidance, and drought escape. The
54 drought tolerance mechanism allows plants to adjust cell osmosis, plasticity and size and produce organic
55 solutes like proline to protect cells from damage caused by water stress [11, 12]. In drought avoidance,
56 plants maintain relatively higher tissue water potential even when surface soil moisture decreases below
57 optimal levels. They may achieve this through deep rooting systems, reduction of radiation absorption in
58 leaves and reduction in hydraulic conductance [13]. The drought escape mechanism involves an
59 accelerated plant cycle through flowering and maturity[7].It is the ability of the crop to rapidly allocate
60 photosynthates to reproductive structures before the onset of a drought [14].

Comment [MB1]: I would suggest adding the name(s) of the author(s) whose work you are citing. That should go throughout the manuscript.

61 Environmental and genetic factors interact to confer drought resistance on plants [15], and one or both
62 factors could be manipulated to enhance any of the afore-mentioned mechanisms in common beans. Soil
63 is a common environmental factor that affects the drought resistance of common beans. The soil's
64 available water capacity (AWC) is an important control on the amount and length of time it can retain
65 water for plant use, and is an effective soil property to manage crop drought resistance, especially in
66 short-term droughts [16, 17]. An increase in soil organic matter (SOM) may increase soil water retention
67 at field capacity and relatively increase AWC [18, 19]. A relatively higher soil water retention capacity
68 implies that crops would have relatively longer access to water for growth.

69 **Poor soil fertility is another major soil constraint to legume production and common bean drought**
70 **tolerance in Ghana and sub-Saharan Africa [20, 21].** In many small-holder farms, legumes are cultivated
71 without external inputs [22, 23]. Though common bean fixes between 2-28 kg/ha nitrogen annually
72 through biological nitrogen fixation (BNF) [24], a proper crop growth requires adequate supply of all other
73 essential nutrients. Previous studies have found up to 80% improvements in common bean yield with
74 phosphorus and potassium fertilizer applications [25, 26, 27]. An adequate supply of nutrients may
75 enhance the drought tolerance of common beans because water-nutrient interactions impact water use
76 and productivity at all levels of crop growth [28, 29, 30]. Crops with adequate nutrient supply often show
77 higher drought tolerance [31] because of the increase in water use efficiency (WUE) [32]. **Water use**
78 **efficiency is the amount of biomass or grain produced per unit water transpired or applied in irrigation.**

79 Thus, when soil moisture and nutrients are adequately supplied, water aids mass flow and transport of
80 nutrients to roots. Water uptake by the roots to meet transpiration needs simultaneously takes up soil
81 nutrients [33].

82 In this study, improving soil organic matter (which controls soil water retention) and nutrient
83 concentrations were the focal points to manage drought resistance of common beans. To address these
84 problems, we explored the integrated use of fertilizer and compost. Integrated fertilizer and organic
85 soil amendment use has been **recommended by [34]** because of the ability of the two resources to jointly
86 supply soil nutrients and improve soil physical properties. Hence the objective of the study was to supply
87 essential nutrients to common beans through fertilizer application while compost improved soil organic
88 matter and consequently soil water retention to mitigate the effects of drought on common beans. The

89 study imposed drought at the flowering to pod-setting period of common bean growth because that is the
90 most sensitive period to drought stress [35, 36]. Many drought tolerant studies have confirmed 80% field
91 capacity (FC) moisture as the optimum moisture level for common bean production [37, 38, 10]. Drought
92 stress up to 16% FC, reported in a previous study [10], was followed to avoid bringing common bean plants
93 to permanent wilting points. Periodic soil moisture, soil nutrients, plant growth and yield data were
94 collected over time to achieve the objective of the study.

95 **2. MATERIALS AND METHODS**

96 **2.1 Study site**

97 The research was conducted in a greenhouse at the Legumes and Oil Seeds Division of the CSIR-Crops
98 Research Institute (CRI), Fumesua in the Ashanti Region of Ghana from September 2021 to January
99 2022. CSIR-CRI is situated at Latitude 6.7 109°N, Longitude 1.5172°W, and 800 m above sea level. It is
100 in the semi-deciduous forest agro-ecological zone (CSIR-CRI weather station). The area has a bi-modal
101 rainfall pattern, with a mean annual rainfall of about 1550 mm. The major rainy season starts from April to
102 the end of July, followed by a dry spell in August, while the minor rainy season continues from September
103 to November every year. Annual temperatures range from a minimum of 21.1°C to a maximum of 32.7°C
104 and a mean of 31.6°C.

105 **2.2 Activities before experimental set-up**

106 The soil's bulk density was determined on a field previously planted to legumes. Soil was collected from
107 this field, sterilized and its field capacity moisture determined before the experimental set-up. Soil
108 samples and compost samples were taken for initial analyses and characterization. Five holes of about
109 2cm diameter were perforated at the bottom of the buckets. Buckets were filled with sterilized soil. The
110 procedures outlined below were followed:

111 **2.2.1 Soil bulk density determination**

112 Three core samplers (cylindrical in shape) were used to collect soil from the field. The core samplers with
113 the soil were weighed and put in an oven at 105°C for two days. After two days, the dried soil samples
114 with the core samplers were weighed. Bulk density was calculated by the formula below:

115 Bulk density = $\frac{\text{mass of dry soil}}{\text{volume of core sampler}}$ equation 1 [39]

116 Mass of dry soil = *weight of core sampler with oven-dried soil (after cooling down)– weight of core*
117 *sampler* equation 2

118 The volume of a core sampler was determined by the formula of the volume of cylinder as follows:

119 Volume of core sampler = $\pi r^2 h$ equation 3

120 Where $\pi = 22/7$; r is the radius of the circular end of the core sampler; h is the height of the core sampler.

121 The average bulk density of soil in the three core samplers was determined.

122 **2.2.2 Soil sterilization**

123 Field soil was collected, thoroughly mixed, filled into barrels and heated over an LPG flame while covered
124 with jute sacks and a lid. The temperature of the soil was monitored with a thermometer on the top of the
125 soil until it reached 100°C. The soil was left to heat on the flame after the 100°C point for three more
126 hours. The prescribed sterilization method [40] was done to combat nematodes and other soil-borne
127 pathogens.

128 **2.2.3 Field capacity moisture determination**

129 Three polyvinyl chloride (PVC) pipes of 25 cm length and 11 cm diameter were marked at 15 cm length.
130 They were taken to the field where soil was collected and pushed down carefully to the 15 cm mark (thus
131 a soil depth of 15 cm was collected). Circular trenches were dug around the pipes to 15 cm depth to
132 enable us carefully carry the PVC pipe with the full depth of soil at the bottom of the pipe with a hand
133 trowel. This was done to ensure that the bulk density of the soil is not altered. The bottom ends of the PVC
134 pipes with soil were covered with plastic netting material and sent to a greenhouse. The soil was flooded
135 with 1 L of tap water from the other open end. The water drained through the net after about 30 seconds
136 of pouring it. The PVC pipes were left on a greenhouse bench for two days with the covered net side
137 raised on two slabs of wood sitting on the greenhouse bench. Gravimetric soil moisture determination
138 was done after two days when no water was visibly draining from the soil through the net. Volumetric soil
139 moisture determination was done with an instant moisture meter. Ten grams of the soil from each pipe
140 was oven dried at 105°C for two days to determine gravimetric soil moisture as follows:

141 Gravimetric soil moisture (g/g) = $\frac{\text{weight of fresh soil (g)} - \text{weight of dry soil (g)}}{\text{weight of dry soil (g)}}$ equation 4 [41]

142 **2.2.4 Filling buckets with soil to simulate field bulk density**

143 Gravimetric soil moisture of the sterilized soil was determined. The sterilized soil was used to fill the
 144 buckets to 15 cm depth. The buckets measured 18 cm deep and 20 cm in diameter (buckets were
 145 cylindrical in shape).The weight of dry soil to fill up to the 15 cm mark was calculated to simulate the field
 146 bulk density. The filling depth and radius of the buckets were used to calculate the filling volume. Mass of
 147 soil used to fill bucket was calculated as follows:

148 Dry mass of soil (g) = Bulk density of field soil (g/cm³)× filling volume of the bucket (cm³)equation 5[39]

149 Filling volume of the bucket = $\pi r^2 h$ equation 6

150 Where $\pi = 22/7$; r is the radius of the circular end of the bucket; h is the filling height (15cm) of soil.

151 To account for moisture content of the sterilized soil in order to fill the exact dry soil weight:

152 Fresh sterilized soil weight (g) = $\frac{\text{dry mass of soil (g)} \times 100\%}{(100\% - \text{gravimetric soil moisture}\%)}$ equation 7

153 The soil was pressed to the 15 cm mark after filling and left to settle for two weeks while buckets were
 154 covered with lids. The buckets were arranged on the screenhouse floor.

155 2.3 Initial soil and compost sampling and analyses

156 Three samples were collected from the sterilized soil (about 100g each) for initial analyses. Three
 157 Municipal Solid Waste (MSW) Compost samples were also analyzed and characterized. The soil was
 158 analyzed for pH[42], organic carbon/matter (OC/M)[43], mineral nitrogen (NO₃⁻, NH₄⁺)[44],Bray P-1
 159 phosphorus (P)[45] and particle size distribution (texture) [46]. Compost was also analyzed for organic
 160 carbon [43], total N[47], P[48] and K[49], spelt out in Table 1 below.

161 Table 1. Characteristics of soil and MSW compost before the start of the experiment

	pH	NO ₃ ⁻	NH ₄ ⁺ mg/kg	Avail. P	OC	OM	Total N %	Total P	Total K	Texture
Soil	6.21	71.95	48.57	33.72	1.79	3.09	-	-	-	Loamy sand
Compost	5.48	-	-	-	38.17	66.96	1.65	0.86	0.79	

162 C:N ratio of compost is 23.13

163 2.4 Experimental design and treatments

164 Split plot in completely randomized design was used in this study. Treatments were moisture regimes (the
 165 main plot factors) and fertility treatments(the sub-plot factors). The levels of moisture regime/drought
 166 stress included D1- 80% FC throughout the growth period till physiological maturity; D2 - 80% FC from
 167 sowing till flowering; water stress from flowering till 40% FC and re-wetting to 80% FC till physiological

168 maturity; and D3 - 80% FC from sowing till flowering; water stress from flowering till 16% FC and re-
169 wetting to 80% FC till physiological maturity. (The only exception to the moisture regimes happened a day
170 before drought imposition, when all the buckets were saturated with water (methodology adopted from
171 [10]).The levels of the fertility treatments were control, full rate glycine mix NPK legume fertilizer (FRG),
172 full rate compost (FRAC), full rate glycine mix + half rate compost (FRG + HRAC) and full rate glycine mix
173 + full rate compost (FRG + FRAC). The compost used in this study was made from a collection of
174 municipal solid waste. There were 15 treatments in total. The treatments were replicated thrice to make a
175 total of 45 buckets.

176 **2.4.1 Application of treatments**

177 Buckets were labelled with their designated treatments after randomizing them on the screen house floor.
178 Compost was applied at 4 t/ha, one month before planting. Compost was weighed and mixed with a
179 gardener's fork to about 4 cm depth. Weights of compost to apply were calculated as follows:

$$180 \text{ Weight of compost} = \frac{4 \text{ t} \times (\text{top surface area of soil in bucket}) \text{ m}^2}{10000 \text{ m}^2} \text{ equation 8}$$

181 Where 4t represents the rate per hectare; 10000 m² is the area of a hectare.

$$182 \text{ Surface area of the soil} = \pi r^2 \text{ equation 9}$$

183 Where $\pi=22/7$; r is the radius of the circular open end of the bucket.

184 Fertilizer was applied at 4 g/plant into two splits. Two grams per plant was applied two weeks after
185 planting and the other 2 g/plant at pod initiation (48 days after planting, thus after returning from drought
186 imposition).The fertilizer contains a proportion of 5:30:30 kg/ha N:P₂O₅:K₂O. The fertilizer was applied by
187 band placement at 3 cm depth and a distance of 5 cm away from the plant, and well covered with soil.

188 **2.4.2 Planting of common beans**

189 Before planting, 100g of *Enepa* common bean variety (a white seeded common bean variety released by
190 the Legumes and Oil Seeds Division of CSIR-Crops Research Institute in Kumasi-Ghana, in 2016) seeds
191 were soaked with tap water for an hour. The seeds were inoculated with 5g Sarifix *Rhizobium* inoculum.
192 Seeds were planted at three per pot with the hand to about 3 cm depth and later thinned to two per pot.
193 Each pot was watered with 500 ml of water at planting. Gloves were worn to prevent cross contamination
194 of the soil.

195 **2.4.2.1 Watering regime and drought imposition**

196 Soil moisture was maintained at 80% FC for all the pots from the start of the experiment. An instant
197 moisture meter was used to estimate volumetric soil moisture to determine how much water to top-up to
198 80% field capacity. After five moisture readings and topping up water every two days, it was determined
199 that an average of 125 ml of water was needed to bring the soil to 80% FC every two days.

200 At the first flower stage (R1 stage), thus 30 days after planting, drought imposition was implemented. A
201 day before drought imposition, soil in all the pots was saturated with 1L of water (adopted from [10]).
202 From that day, soil in pots receiving treatment D1 continued to be maintained at 80% FC. Soil moisture in
203 pots receiving D2 was monitored from the day of saturation till 40% FC. It took 8 days to reach this FC
204 and 80% FC was returned until physiological maturity. Pots receiving treatment D3 was monitored till 16%
205 FC and then returned to 80% FC till harvest. It took 15 days to reach 16% FC.

206 **2.4.2.2 Data collection**

207 Data was collected on volumetric soil moisture, plant height, leaf number, leaf area, Soil Plant Analysis
208 Development (SPAD) chlorophyll (surrogate) concentration of leaves at 3, 7 and 10 weeks after planting
209 (WAP). Data on number of pods, pod weight, number of seeds per pod, seed weight, biomass and soil
210 samples were collected at harvest (71 days after planting). Biomass and soil nutrient statuses were
211 analyzed in the laboratory after harvest. The biomass was analyzed for total nitrogen (N), phosphorus (P)
212 and potassium (K). The soil was analyzed for pH, organic carbon/matter (OC/M), nitrates (NO_3^-) and
213 ammonium (NH_4^+).

214 **2.4.2.3 Harvest**

215 All pods were picked from the plants in the pots into labeled envelopes when the plants were at
216 physiological maturity, 71 days after planting (DAP). The remaining above-ground biomass was cut at
217 root level into labeled envelopes. The samples were oven-dried at 60°C for two days. The pods were
218 weighed and shelled. The seeds were also weighed as g/surface area of soil in the bucket and
219 extrapolated to kg/ha.

220 **2.4.2.4 Nutrient and water use efficiencies**

221 Nitrogen and water use efficiencies were calculated by the following formulae:

222 Nitrogen use efficiency =
$$\frac{\text{nitrogen uptake in grain yield } \left(\frac{\text{kg}}{\text{ha}}\right)}{[(\text{initial nutrient} + \text{fertilizer nutrient}) - \text{residual nutrient after harvest}] \text{ kg/ha}}$$
 equation 10 [50]

223 Water use efficiency = $\frac{\text{grain yield (kg/ha)}}{\text{amount of irrigation water applied (mm)}}$ equation 11 [32]


224 2.5 Statistical analyses

225 Analyses of variances in the data conferred by the fertility treatments and drought imposition were
226 determined using IBM SPSS statistics 2.0 package. Statistically significant treatment means were
227 separated with Fisher's least significant difference (LSD) at 5% probability. Regression analysis was used
228 to determine relationship between nitrogen and water use efficiencies in Excel.

229 3. RESULTS

231 3.1 Periodic soil moisture measurements

232 Soil moisture was not affected ($P > 0.05$) by drought imposition or its interaction with fertilizer treatments
233 on any of the sampling days. However, the fertility treatments significantly affected soil moisture at 3 WAP
234 ($P < 0.001$), 7 WAP ($P < 0.001$) and 10 WAP ($P = 0.007$). On all the sampling days, FRG, FRG + HRAF and
235 FRG + FRAC affected the highest soil moisture on average (6.9 - 9 cm^3/cm^3) and between 20 to 38%
236 more than compost alone and the control (Fig. 1).




237 Fig. 1. Volumetric soil moisture (cm^3/cm^3) affected by fertility treatments at 3 WAP (before drought
238 imposition) and at 7 and 10 WAP (after drought imposition). Error bars represent standard errors of the
239 means. Different lower case letters on top of the bars mean significant differences between the treatment
240 means. Different lower case letters on top of the bars mean significant differences between the treatment
241 means.

242 3.2 Common bean growth parameters

243 Plant height was not influenced ($P>0.05$) by the fertility treatments, drought imposition or their interactions
 244 in any of the sampling days.

245 Fertility treatments affected ($P<0.001$) leaf number at 3 WAP (before drought imposition) (Fig. 2A). Plants
 246 applied with FRG alone had the most number of leaves (12.7) about 10 to 35% more than other
 247 amendments. There was 10-24% reduction in leaf number with the addition of half and full rates of the
 248 compost to FRG. Interaction between fertility treatments and drought imposition affected ($P= 0.01$) leaf
 249 number at 7WAP (after drought imposition) (Fig. 2B). Plants applied with FRG x D2 at flowering had the
 250 most number of leaves (40) while plants applied with FRACx D3 at flowering had the least number of
 251 leaves (10). The latter was similar to the number of leaves affected by control x D1; control x D3; FRAC x
 252 D1; FRAC x D3; FRG x D3; (FRG + HRAC)x D2. The average number of leaves affected by fertility
 253 treatments alone was in the order $FRG \geq (FRG + FRAC) > (FRG + HRAC) > FRAC = Control$. Leaf number
 254 was not influenced ($P>0.05$) by the fertility treatments, drought imposition or their interactions at 10 WAP.

255
 256
 257
 258
 259
 260




261
 262
 263
 264

265
266
267

268 Fig. 2. Leaf number affected by fertility treatments at 3 WAP (before drought imposition) (A); leaf number
269 affected by the interaction between fertility treatments and drought stress at 7 WAP (after drought
270 imposition) (B). Error bars represent standard errors of the means. Different lower case letters on top of
271 the bars mean significant differences between the treatment interaction means. Upper case letters on top
272 of the bars represent significant differences between corresponding fertility treatment means.


273
274 Leaf area was affected ($P<0.001$) by fertility treatments at 3 WAP (Fig. 2). FRG affected the highest leaf
275 area between 19-68% higher than the other amendments. The addition of half or a full rate of the
276 compost to FRG reduced leaf area by 19 – 43%. At 7 WAP (after drought imposition), FRG+FRAC
277 affected ($P<0.001$) the largest leaf area which was similar to that affected by FRG. The application of
278 FRG+HRAC, FRAC and control affected the smallest leaf area (Fig. 3). Leaf area at 10 WAP was not
279 affected ($P>0.05$) by fertility treatments, drought stress regimes or their interactions.



280
281 Fig. 3. Leaf area (cm²) affected by fertility treatments at 3 WAP (before drought imposition) and 7 WAP
282 (after drought imposition). Error bars represent standard errors of the means. Different lower case letters
283 on top of the bars mean significant differences between the treatment means.

284 SPAD chlorophyll content was affected ($P<0.001$) by fertility treatments at 3 WAP (Fig. 4A). FRG affected
285 the highest leaf chlorophyll concentration (47 SPAD units) which was between 14 – 37% more than other
286 amendments. Leaf chlorophyll was reduced by 14% with the addition of half and full rates of the compost.

287 FRAC and the control affected the least leaf concentration. Leaf chlorophyll concentration was not
 288 affected ($P>0.05$) by the interaction of fertility treatments and drought stress regimes at 7 WAI. However,
 289 drought regimes affected ($P=0.02$) leaf chlorophyll concentration at 7 WAP (Fig. 4B). D1 and D2 affected
 290 the highest leaf chlorophyll concentration (~ 35), about 17% more than D3. Leaf chlorophyll concentration
 291 at 10 WAP was not affected ($P>0.05$) by fertility treatments, drought stress regimes or their interactions.




292
 293 Fig. 4. SPAD leaf chlorophyll concentration affected by fertility treatments at 3 WAP (before drought
 294 imposition) (A); leaf chlorophyll concentration affected by drought stress regimes at 7 WAP (after drought
 295 imposition) (B). Error bars represent standard errors of the means. Different lower case letters on top of
 296 the bars mean significant differences between the treatment means.

297 3.3 Common bean yield parameters

298 Grain weight was affected by fertility treatments only ($P=0.05$) but not drought or its interaction with
 299 fertility treatments (Fig. 5A). FRG affected the highest grain weight (96 kg/ha). The addition of half and full
 300 rates of compost to FRG reduced grain weight by 84 and 56%, respectively. Weights affected by fertility
 301 treatments other than FRG were statistically similar.

302 There was a significant interaction between fertility treatments and imposed drought ($P=0.008$) on
 303 common bean dry biomass at harvest (Fig. 5B). FRG imposed with 40% FC drought stress at flowering
 304 affected the largest biomass (3055 kg/ha) while the control at 80% FC throughout the study affected the
 305 least. On average, FRG alone affected the highest biomass (2037 kg/ha) which was similar to FRG +


306 FRAC but between 48 to 70% higher than biomass affected by FRG + HRAC and other fertility
 307 treatments. The control affected 24% higher common bean biomass than compost application alone.
 308 Other yield parameters (number of pods, pod weight, number of seeds) were not affected by fertility
 309 treatments, drought imposition or their interactions ($P > 0.05$)



311 Fig. 5. Grain yield (kg/ha) affected by fertility treatments at harvest (A); biomass (kg/ha) by
 312 interaction between fertility treatments and drought imposition at harvest (B). Error bars represent
 313 standard errors of the means. Different lower case letters on top of the bars mean significant differences
 314 between the treatment means (A) and treatment interaction means (B). Upper case letters on top of the
 315 bars represent significant differences between corresponding fertility treatment means.


316 **3.4 Soil organic matter and nutrient statuses after harvest**

318 There was significant interaction between fertility treatments and drought imposition on organic carbon
 319 ($P=0.01$) and organic matter ($P=0.01$) at the end of the study (Fig. 6 A&B). Control x D2; FRAC x D1;
 320 FRG x D2; (FRG+HRAC) x D2 and (FRG+FRAC) x D1 affected up to 7% more organic matter than initial
 321 soil organic matter before treatment imposition. However, the application of the fertility treatments alone
 322 did not affect ($P > 0.05$) soil organic matter but in general soil organic carbon and matter followed the order
 323 D1 > D2 > D3 for drought imposition.



326

327



328


329 Fig. 6. Organic carbon (%) affected by interaction between fertility treatments and drought imposition at
 330 harvest (A); organic matter (%) affected by interaction between fertility treatments and drought imposition
 331 at harvest (B). Error bars represent standard errors of the means. Different lower case letters on top of
 332 the bars mean significant differences between the treatment interaction means. Short dash lines mark the
 333 initial organic carbon and organic matter percentages before treatment imposition.

334 At the end of harvest, all treatments had residual soil mineral N ($\text{NO}_3^- + \text{NH}_4^+$) levels lower than the initial
 335 soil mineral N concentration (121 mg/kg). The interaction between fertility treatments and drought
 336 imposition significantly affected ($P < 0.005$) residual soil mineral N (Fig. 7). FRG treatment with drought
 337 imposition at 16% FC (D3) retained the highest soil N concentration (113 mg/kg) which was similar to
 338 FRAC x D3; FRG x D1; FRG x D2; (FRG+HRAC) x D1,D2&D3 and (FRG+FRAC) x D1,D2&D3. The
 339 control at D3 retained the least amount of soil mineral N (23 mg/kg). On average, FRG, FRG + HRAC and
 340 FRG+FRAC retained the highest and similar concentrations of mineral N (~ 95 mg/kg) which was 25-53%
 341 more than concentrations retained by FRAC alone and the control. The fertility treatments, drought
 342 imposition and their interactions had no effect ($P > 0.05$) on residual phosphorus concentration and soil pH.

343

344

345



346

347

348

349

350


351

352 Fig.7. Residual mineral N (mg/kg) affected by interaction between fertility treatments and drought
353 imposition. Error bars represent standard errors of the means. Different lower case letters on top of the
354 bars mean significant differences between the treatment interaction means. Different uppercase letters
355 represent differences in corresponding fertility treatments (c). Short dash lines mark the initial mineral N
356 concentration before treatment imposition.

357

358 **3.5 Common bean nutrient uptake**

359 The fertility treatments, drought imposition and their interactions had no effect ($P>0.05$) on plant biomass
360 N and P uptake. However, fertility treatments alone ($P=0.002$) and drought imposition alone ($P=0.001$)
361 affected K uptake in common beans biomass (Fig. 8 A&B). FRG affected the highest K uptake (3%)
362 which was 5 to 18% more than FRAC, FRG+HRAC, FRG+FRAC but was statistically similar to them.
363 Drought imposition to 40% FC at flowering affected 2.7% common bean biomass K uptake, similar to
364 plants that had no moisture stress. However, K uptake was significantly reduced by more than 20% when
365 plants were water stressed till 16% FC at flowering.




366 A
 367 Fig. 8. Potassium uptake (%) in plant biomass as affected by soil fertility treatments (A). Potassium
 368 uptake in plant biomass as affected by drought imposition (B). Error bars represent standard errors of the
 369 means. Different lower case letters on top of the bars mean significant differences between the treatment
 370 means.

371 **3.6 Water use efficiency of common beans**

372 Water use efficiency was affected ($P=0.05$) by the soil fertility treatments. Common bean plants were
 373 most efficient with water use with the application of FRG (0.58 kg/ha/mm WUE) (Fig. 9A). Addition of half
 374 and full rates of compost to FRG reduced water use efficiency by 55 to 64%. The least efficient water use
 375 was affected by the control (0.018 kg/ha/mm) and FRG +HRAC (0.095 kg/ha/mm) (Fig. 9A).

376 There was a strong positive relationship ($R^2=73%$) between water use efficiency (WUE) and nitrogen use
 377 efficiency (NUE) (Fig. 9B)



378

379 Fig. 9. Water Use Efficiency (kg/ha/mm) of common beans affected by fertility treatments(A). Error bars
380 represent standard errors of the means. Different lower case letters on top of the bars mean significant
381 differences between the treatment means. Relationship between water use efficiency (WUE) and nitrogen
382 use efficiency (NUE)(B).

383 **4. DISCUSSION**

384 **4.1 Soil organic matter and water retention**

385 The increase in soil moisture by the full rate of glycine mix fertilizer(FRG) alone and its addition with half
386 (FRG + HRAC) and full rates (FRG + FRAC) of compost (Fig. 1) could not be attributed to organic matter
387 because none of these treatments on their own, affected soil organic matter (SOM) (Fig. 6). However, it is
388 possible that the short height and spreading architecture of common beans, and the relatively large leaf
389 number and area (Fig.2&3) affected by nutrient supply from FRG included treatments, shaded the soil
390 surface, reduced the reach of incident solar radiation, and reduced excessive evaporation. This
391 confirms previous reports that using live plants as soil cover increases soil moisture by allowing more water
392 to sink in and reducing evaporation [51]. Moreover, achieving improvement in SOM to consequently
393 impact soil water retention usually does not occur with one application because many research findings
394 which successfully achieved such, applied compost for two or more years [52,53, 54]. Generally,
395 maintaining soil moisture at 80% FC (no moisture stress) affected higher soil organic carbon and matter
396 percentages because optimal microbial activity occurs near field capacity moisture [51], while drought
397 stress reduces microbial activity and organic matter build-up [55].

398 **4.2 Common bean growth, yield, nutrient uptake and residual nitrogen**

399 Lower residual soil mineral N concentration below the initial level could be attributed to common bean N
400 uptake to meet its physiological needs [56,57]. Already a poor biological N fixer [56, 58, 59], the supply of
401 N from the treatments and drought imposition may have further compromised its nitrogen fixing abilities
402 [60, 61] leading to the unexpected decline in N. However, since no significant differences were observed
403 in N uptake by the plants and NUE affected by all treatments was extremely low, it is possible that aside
404 plant N uptake, immobilization may have also caused the decline leading to the differences in residual N
405 levels observed. Based on a previous compost study [62], the C:N ratio of our compost implied that, there
406 should have been a balance between N mineralization and immobilization. However, judging from the

407 non-corresponding increase in residual soil N above FRG, with the addition of half and full rates of
408 compost to FRG (Fig. 7), it is evident that N was immobilized more than mineralized with the addition of
409 compost. This is confirmed by the decline in crop growth rate (leaf number, area and chlorophyll
410 concentration) and yield (grain yield and biomass at harvest) parameters with the addition of the compost
411 alone or in combination with FRG relative to FRG alone (Fig. 2, 3, 4A and 5). According to a study[63],
412 the compost applied belongs to category 3 of organic amendments (because its total N was below 2.5%
413 but C: N ratio was below 25) which implies that it should be mixed with inorganic fertilizer for application,
414 just as practiced in our experiment. However, studies on composting municipal solid waste compost(as
415 used in our study) often report C:N ratios between 10 to 18 [64, 65, 66] when compost is completely
416 decomposed. The high carbon percentage (38%) in our compost, organic matter content above 65%
417 (calculated from C%) and C:N ratio of 23 suggest that the compost may not have been properly cured
418 and continued decomposition after application [64, 67, 68]. In such cases, continuous decomposition
419 leads to the loss of organic matter through microbial respiration and immobilization of N during the
420 decomposition process [66, 67, 68]. This could have contributed to why compost addition did not improve
421 SOM above initial levels (Fig. 6).

422 The generally low common bean yield in this study (highest yield was about 100 kg/ha compared to
423 average yield of 437 kg/ha in Sub-Saharan Africa [69]) was expected because of the sensitivity of the
424 flowering and pod initiation growth periods to drought events[35]. However, comparatively, the ready
425 supply of relatively high levels of P and K and a starter N from FRG (NPK 5:30:30) without high N
426 immobilization rates like compost included treatments, and the timing of its application, caused it to
427 increase common bean growth (leaf number, area, SPAD chlorophyll) and yield (grain yield and biomass)
428 parameters compared to other fertility treatments. Split application of FRG supplied a starter N during the
429 temporal N deficiency stage of seedling growth when cotyledon reserves were depleted, leading to fast
430 vegetative growth [70]. Split application of FRG at the pod development stage supplied the necessary
431 nutrients for dry matter partitioning into pods and grain yield [71]. Common bean yield increases of up to
432 3600 kg/ha with the application of 0- 280 kg/ha P₂O₅ and 0-200 kg/ha K₂O even in soils with inherently
433 high P and K levels have been reported [27]. Starter N application between 0-46 kg/ha N with

434 *Rhizobium* inoculation has also been found to increase common bean yield by 32%, though nodulation
435 and biological N fixation were compromised [72].

436 Higher K uptake by plants supplied with FRG included treatments was only an artifact of high K supply
437 from them. Higher K uptake affected by the constant supply of 80% FC moisture compared to the water
438 stressed plants (Fig. 8) confirms that K mobility and availability to the crops was increased by the
439 availability of water, since soil moisture is one of the key factors controlling K availability and uptake [73].
440 Since soil moisture levels below 100% FC do not cause significant K leaching [74], it can be assumed
441 that there was no or minimal K leaching in this study, hence low K uptake in water stressed plants could
442 not be attributed to K leaching. Other authors have also reported low K uptake in common beans under
443 severe moisture stress conditions [75].

444 **4.3 Water and nitrogen use efficiencies**

445 Consistent N supply from FRG and the high water retention affected by FRG, presented common bean
446 plants with better growth conditions (nutrients and water) to produce higher biomass and yield (Fig. 5).
447 These components translated into higher WUE for the same amounts of water supplied to all treatments.
448 Conditions that reduce soil evaporation have been reported to also increase the WUE of
449 crops [76]. Though NUE was generally extremely poor (Fig. 9B), the strong positive relationship between
450 NUE and WUE confirms that better supply and use of soil nitrogen by crops could allow them to efficiently
451 use water as well. On the reverse, there is also an intricate relationship between water and nitrogen
452 concentration in the soil that allows crops greater access to nitrogen with adequate water supply, through
453 transpiration driven mass flow of nutrients in the soil and uptake by roots [33]. Nitrogen affects stromal
454 and thylakoid proteins in leaves which in turn affects the photosynthetic capacity of plants [77]. Hence,
455 relatively higher availability of both water and N by FRG led to the higher WUE. A previous study [78] also
456 found 35 - 45% increases in common bean yield with the application of 80, 170 and 225 kg/ha N to wheat-
457 common bean rotations under drought conditions, compared to no nitrogen application. Nitrogen
458 immobilization affected by adding half and full rates of the compost to the fertilizer (FRG), and poor
459 nutrient supply from the control led to poor WUE affected by these treatments.

460 **5. CONCLUSION**

461 This study confirms that, drought at the reproductive and pod initiation stages of common beans, has
462 great impact and generally reduces its yield irrespective of agronomic practices implemented. However, a
463 good supply of nutrients at the right time may offset some of the yield decline in the event of drought.
464 Thus, the application of recommended rates of nitrogen, phosphorus and potassium at the vegetative and
465 reproductive stages of the crop increases growth and yield components and reduces the severity of short-
466 term droughts by improving the water use efficiency of the crop. However, research is still needed to
467 ascertain the exact amount of N fertilizer to apply to maximize biological nitrogen fixation by common
468 beans. Combined use of compost and fertilizer to improve common bean yield and mitigate drought
469 effects, should seriously consider the quality of the compost because compost would not complement
470 inorganic fertilizer to mitigate drought effects if it is not of required quality.

471 **ACKNOWLEDGEMENT**

472 The authors are grateful to the (CSIR)-Crops Research Institute, Ghana, and RDA- National Institute of
473 Agricultural Sciences, Korea, for providing an enabling environment, equipment and materials needed for
474 the implementation of this work and writing of this manuscript.

475 **STATEMENTS AND DECLARATIONS**

476 **Funding**

477 This project was self-funded

478 **AVAILABILITY OF DATA AND MATERIALS**

479 The dataset generated and/or analyzed during the study are available from the corresponding author
480 upon reasonable request.

481 **COMPETING INTERESTS**

482 Authors declare no conflicts of interests.

483 **REFERENCES**

- 484 1. Khatun M, Sarkar S, Era FM, Mominul, Islam AKM, Anwar MP, Fahad S, Datta R, Islam AKMA. Drought
485 stress in grain legumes: Effects, tolerance, mechanisms and management. *Agron.* 2021; 11, 2374.
486 <https://doi.org/10.3390/agronomy11122374>
- 487 2. Araujo SS, Beebe S, Crespi M, Delbreil B, Gonzalez EM, Gruber V, Lejeune-Henaut I, Link W,
488 Monteros MJ, Prats E. Abiotic stress responses in legumes: Strategies used to cope with environmental
489 challenges. *Crit Rev Plant Sci.* 2015; 34, 237-280. <https://doi.org/10.1080/07352689.2014.898450>

- 490 3. Porch, Beaver SJ, Debouck GD, Jackson AS, Kelly DJ, Dempewolf H. Use of wild relatives and closely
491 related species to adapt common bean to climate change. *Agron.*2013;3, 433-462.
492 <https://dx.doi.org/10.3390/agronomy3020433>
- 493 4. Polania J, Poschenreider C, Rao I, Beebe S. Estimation of phenotypic variability in symbiotic nitrogen
494 fixation ability of common bean under drought stress using N-15 natural abundance in grain. *Europe J*
495 *Agron.* 2016; 79, 66-73. <https://doi.org/10.1016/j.eja.2016.05.014>
- 496 5. Adonadag M-G, Ampadu B, Ampofo S, Adiali F. Climate change adaptation strategies towards
497 reducing vulnerability to drought in Northern Ghana. *Europe J Environ Earth Sci.*
498 2022.<https://doi.org/10.24018/ejgeo.2022.3.4.294>
- 499 6. Marinus W, Ronner E, van de Ven GWJ, Kanampiu F, Adjei-Nsiah S, Giller KE. What role for legumes
500 in sustainable intensification?- case studies in Western Kenya and Northern Ghana for PROIntensAfrica.
501 N2Africa, Putting nitrogen fixation to work for smallholder farmers in Africa. 2016.
502 www.edepot.wur.nl/397988
- 503 7. Beebe S, Chirwa R, Rubyogo JC, Clare M, Bodo R, Tumsa K, Mutari B, Nkalubo S, Chisale V,
504 Macharia D, Kamau E, Kweka S, Kilanjo M, Karanja D, Ugen M, Demissie DA, William M, Negash K,
505 Yilma B, Kabungo C, Bekele A. Enhancing common bean productivity and production in Sub-Saharan
506 Africa. In: Seven seasons of learnings. 2020.[https://tropicallegumeshub.com/wp-](https://tropicallegumeshub.com/wp-content/uploads/2020/06/Seven_seasons_Of_Learnings-06_Beebe_et_al.pdf)
507 [content/uploads/2020/06/Seven_seasons_Of_Learnings-06_Beebe_et_al.pdf](https://tropicallegumeshub.com/wp-content/uploads/2020/06/Seven_seasons_Of_Learnings-06_Beebe_et_al.pdf)
- 508 8. Smith MR, Veneklaas E, Polania J, Idupulapati MR, Beebe SE, Merchant A. Field drought conditions
509 impact yield but not nutritional quality of the seed in common bean (*Phaseolus vulgaris* L.) *Plos*
510 *One.*2019.<https://doi.org/10.1371/journal.pone.0217099>
- 511 9. Szilagyi L. Influence of drought on seed yield components in common bean. *Bulg J Plant Physio.* 2003;
512 43, 320-330.
513 [www.researchgate.net/publication/237551508_Influence_of_drought_on_seed_yield_components_in_co](http://www.researchgate.net/publication/237551508_Influence_of_drought_on_seed_yield_components_in_common_bean)
514 [mmon_bean](http://www.researchgate.net/publication/237551508_Influence_of_drought_on_seed_yield_components_in_common_bean)
- 515 10. Dipp CC, Marchese JA, Woyann LG, Bosse MA, Roman MH, Gobatto DR, Paludo F, Fedrigo K,
516 Kovali K, Finatto T. Drought stress tolerance in common bean: what about highly cultivated Brazilian
517 genotypes? *Euphytica* 2017; 213, 102. <https://doi.org/10.1007/s10681-017-1893-5>
- 518 11. Mwenye OJ, van RensburgL, van Biljon A, van der Merwe R. The role of proline and root traits on
519 selection for drought stress tolerance in soybeans: a review. *S Afr J Plant Soil* 2016; 33(4),1-12.
520 <https://doi.org/10.1080/02571862.2016.1148786>
- 521 12. Assefa T, Wu J, Beebe SE, Rao IM, Marcomin D, Claude RJ. Improving adaptation to drought stress
522 in small red common bean: phenotypic differences and predicted genotypic effects on grain yield, yield
523 components and harvest index. *Euphytica* 2015; 203(3), 477-489. [https://doi.org/10.1007/s10681-014-](https://doi.org/10.1007/s10681-014-1242-x)
524 [1242-x](https://doi.org/10.1007/s10681-014-1242-x)
- 525 13. Namugwanya M, Tenywa JS, Otabbong E, Mubiru DN, Masamba TA. Development of common bean
526 (*Phaseolus vulgaris* L.) production under low soil phosphorus and drought in sub-Saharan Africa: a
527 review. *J Sustain Dev.* 2014; 7(5), 128. <https://doi.org/10.5539/jsd.v7n5p128>
- 528 14. Villordo-Pineda E, Gonzalez-Chavira MM, Giraldo-Carbajo P, Acosta-Gallegos JA, Caballero-Perez J.
529 Identification of novel drought-tolerant-associated SNPs in common bean (*Phaseolus vulgaris*). *Front*
530 *Plant Sci.* 2015.<https://doi.org/10.3389/fpls.2015.00546>
- 531 15. Benlloch R, Lois ML. Sumoylation in plants: mechanistic insights and its role in drought stress. *J*
532 *Experiment Bot.* 2018.<https://doi.org/10.1093/jxb/ery233>

- 533 16. Herawati A, Mujiyo, Syamsiyah J, Baldan SK, Arifin I. IOP Conf. Ser.: Earth Environ Sci. 2021; 724
534 012014. <https://doi.org/10.1088/1755-1315/724/1/012014>
- 535 17. USDA. Effects on soil water holding capacity and soil water retention resulting from soil health
536 management practices implementation.
537 2018. [www.file:///C:/Users/Administrator/Downloads/AWC_Effects_on_Soil_Water_Holding_Capacity_and](http://www.file:///C:/Users/Administrator/Downloads/AWC_Effects_on_Soil_Water_Holding_Capacity_and_Retention.pdf)
538 [_Retention.pdf](http://www.file:///C:/Users/Administrator/Downloads/AWC_Effects_on_Soil_Water_Holding_Capacity_and_Retention.pdf)
- 539 18. Lal R. Soil organic matter and water retention. Agron J.2020;112 (1).
540 <https://doi.org/10.1002.agj2.20282>
- 541 19. Yang F, Zhang G-L, Yang J-L, Li D-C, ZhaoY-G, Liu F, Yang R-M, Yang F. Organic matter controls of
542 soil water retention in an alpine grassland and its significance for hydrological processes. J Hydro,
543 Elsevier. 2014; 519, Part D, 3086-3093. <https://doi.org/10.1016/j.hydro.2014.10.054>
- 544 20. Awuni GA, Reynolds DB, Goldsmith PD, Tamimie CA, Denwar NN. Agronomic and economic
545 assessment of input bundle of soybean in moderately acidic Savannah soils in Ghana. Agroeco Geosci &
546 Environ. 2020. <https://doi.org/10.1002/agg2.20085>
- 547 21. Khojely DM, Ibrahim SE, Sapey E, Han T. History, current status and prospects of soybean
548 production ad research in sub-Saharan Africa. The Crop J. 2018; 6, 226-235.
549 <https://doi.org/10.1016/j.cj.2018.03.006>
- 550 22. Goldsmith PD. The Faustian bargain of tropical soybean. Tropical Conserv Sci. 2017; 10, 1-4.
551 <https://doi.org/10.1177.1940082917723892>
- 552 23. Tamimie CA. Determinants of soybean adaptation and performance in northern Ghana (Master's
553 Thesis). Graduate College of the University of Illinois at Urbana-Champaign.
554 2017. [https://www.ideals.illinois.edu/bitstream/handle/2142/97450/TAMIMIE-THESIS-](https://www.ideals.illinois.edu/bitstream/handle/2142/97450/TAMIMIE-THESIS-2017.pdf?sequence=1&isAllowed=y)
555 [2017.pdf?sequence=1&isAllowed=y](https://www.ideals.illinois.edu/bitstream/handle/2142/97450/TAMIMIE-THESIS-2017.pdf?sequence=1&isAllowed=y)
- 556 24. Ngome AF. The contribution of nitrogen fixation by field-grown common beans (*Phaseolus vulgaris* L.)
557 to N balances in agricultural production systems of Kakamega District, Kenya. BIOTA East Africa.
558 2006. [www.researchgate.net/publication/320244278_he_Contribution_of_nitrogen_fixation_by_field-](http://www.researchgate.net/publication/320244278_the_Contribution_of_nitrogen_fixation_by_field-grown_common_beans_Phaseolus_vulgaris_L_to_N_balances_in_agricultural_production_systems_of_Kakamega_District_Kenya)
559 [grown_common_beans_Phaseolus_vulgaris_L_to_N_balances_in_agricultural_production_systems_of](http://www.researchgate.net/publication/320244278_the_Contribution_of_nitrogen_fixation_by_field-grown_common_beans_Phaseolus_vulgaris_L_to_N_balances_in_agricultural_production_systems_of_Kakamega_District_Kenya)
560 [Kakamega_District_Kenya](http://www.researchgate.net/publication/320244278_the_Contribution_of_nitrogen_fixation_by_field-grown_common_beans_Phaseolus_vulgaris_L_to_N_balances_in_agricultural_production_systems_of_Kakamega_District_Kenya)
- 561 25. Wondimu WG, Tana T. Yield response of common bean (*Phaseolus vulgaris* L.) varieties to combined
562 application of nitrogen and phosphorus fertilizers at Mechara, Eastern Ethiopia. J Plant Biol Soil Health.
563 2019; 4(2), 7.
564 [www.researchgate.net/publication/334680391_Yield_Response_of_Common_Bean_Phaseolus_vulgaris](http://www.researchgate.net/publication/334680391_Yield_Response_of_Common_Bean_Phaseolus_vulgaris_to_combined_Application_of_Nitrogen_and_Phosphorus_Fertilizers_at_Mechara_Eastern_Ethiopia)
565 [_to_combined_Application_of_Nitrogen_and_Phosphorus_Fertilizers_at_Mechara_Eastern_Ethiopia](http://www.researchgate.net/publication/334680391_Yield_Response_of_Common_Bean_Phaseolus_vulgaris_to_combined_Application_of_Nitrogen_and_Phosphorus_Fertilizers_at_Mechara_Eastern_Ethiopia)
- 566 26. Dejene T, Tama T, Urage E. Response of common bean (*Phaseolus vulgaris* L.) to application of lime
567 and phosphorus on acidic soil of Areka, Southern Ethiopia. J Nat Sci Res. 2016; ISSN 2225-0921.
568 [www.academia.edu/34852847/Response_of_Common_Bean_Phaseolus_vulgaris_L_to_Application_of](http://www.academia.edu/34852847/Response_of_Common_Bean_Phaseolus_vulgaris_L_to_Application_of_Lime_and_Phosphorus_on_Acidic_Soil_of_Areka_Southern_Ethiopia)
569 [Lime_and_Phosphorus_on_Acidic_Soil_of_Areka_Southern_Ethiopia](http://www.academia.edu/34852847/Response_of_Common_Bean_Phaseolus_vulgaris_L_to_Application_of_Lime_and_Phosphorus_on_Acidic_Soil_of_Areka_Southern_Ethiopia)
- 570 27. Carvalho MCS, Nascente AS, Ferreira GB, Mutadiua CAP, Denardin JE. Phosphorus and potassium
571 fertilization increase common bean grain yield in Mozambique. Revista Brasileira de Engenharia
572 Agricolae Ambiental. 2018; 22(5), 308-314. <http://dx.doi.org/10.1590/1807-1929/agriambi.v22n5p308-314>
- 573 28. He X, Qiu H, Xie K, Wang Y, Hu J, Li F, An J. Effects of water saving and nitrogen reduction on the
574 yield, quality, water and nitrogen use efficiency of *Isatis indigotica* in Hexi Oasis. Sci Rep. 2022; 12,
575 550. <https://doi.org/10.1038/s41598-021-04585-x>

- 576 29. Liu Q, Xu H, Mu X, Zhao G, Gao P, Sun W. Effects of different fertilization regimes on crop yield and
577 soil water use efficiency of millet and soybean. Sustainability.2020; 12.
578 4125. <https://doi.org/10.3390/su12104125>
- 579 30. Sharma B, Molden D, Cook S. Water use efficiency in agriculture: measurement, current situation and
580 trends. In: Managing water and fertilizer for sustainable agricultural intensification.
581 2012. www.publications.iwmi.org/pdf/H046807.pdf
- 582 31. Wang B, Liu W, Dang T. Effect of phosphorus on crop water and nitrogen use efficiency under
583 different precipitation year in dryland. Proceedings of International Symposium on water resources and
584 environmental protection.2011; 2111-2114. <https://doi.org/10.1109/ISWREP.2011.5893679>
- 585 32. Ritchie JT. Efficient water use in crop production: Discussion on the generality between biomass
586 production and evapotranspiration. In: Taylor, H.M., Jordan, W., Sinclair, T.R. (eds). Limitations to
587 efficient water use in crop production. American Society of Agronomy, Madison, Wisconsin.1983; 29-
588 44. <https://doi.org/10.2134/1983.limitationstoeficientwateruse.c2>
- 589 33. Plett DC, Ranathunge K, Melino VJ, Kuya N, Uga Y, Kronzucker HJ. He intersection of nitrogen
590 nutrition and water use in plants: new path toward improved crop productivity. J Experiment Bot. 2020;
591 71(15), 4452-4468. <https://doi.org/10.1093/jxb/eraa049>
- 592 34. Voltr V, Mensik L, Hlisnikovsky L, Hruska M, Pokorny E, Pospisilova L. The soil organic matter in
593 connection with soil properties and soil inputs. Agron.2021;11.779.
594 <https://doi.org/10.3390/agronomy11040779>
- 595 35. Ntukamazina N, Onwonga RN, Sommer R, Makankusi CM, Mburu J, Rubyogo JC, Moral MT. Effect of
596 excessive and minimal soil moisture stress on agronomic performance of bush and climbing bean
597 (*Phaseolus vulgaris* L.). Cogent Food and Agric. 2017;3(1).
598 <https://doi.org/10.1080/23311932.2017.1373414>
- 599 36. Manjeru P, Madanzi T, Makedredza B, Ncilzah A, Sithole M. Effects of water stress at different growth
600 stages on yield and yield components of common bean (*Phaseolus vulgaris*). African Crop Sci Conf
601 Proceeds. 2007; 8, 299-303. <https://doi.org/10.13140/RG.2.1.2500.5924>
- 602 37. Smith MR, Dinglasan E, Veneklaas E, Polania J, Rao IM, Beebe SE, Merchant A. Effect of drought
603 and low P on yield and nutritional content in common bean. Front Plant Sci. 2022;
604 <https://doi.org/10.3389/fpls.2022.814325>
- 605 38. Rivera M, Polania J, Ricaurte J, Borrero G, Beebe S, Rao I. Soil compaction induced changes in
606 morpho-physiological characteristics of common beans. J Soil Sci and Plant Nutri. 2019; 19, 217-227.
607 <https://doi.org/10.1007/s42729-019-0007-y>
- 608 39. Blake GR, Hartge KH . Bulk density In: Klute, A Ed, Methods of soil analysis, part 1- Physical and
609 Mineralogical methods, 2nd edition, Agronomy Monograph 9, American Society of Agronomy- Soil
610 Science Society of America, Madison. 1986; 363-382. <https://doi.org/10.2136/sssabookser512edc13>
- 611 40. Lawrence WJK. Soil sterilization. Unwin Brothers Limited. 1955.
- 612 41. Gardener WH . Water content. In: Klute, A Methods of soil analysis, part 1 Physical and mineralogical
613 methods Agronomy Monograph 9 2nd Edition. SSSA Book Series. 1986.
614 <https://doi.org/10.2136/sssabookser5.1.2ed.c21>
- 615 42. Thomas GW . Soil pH and soil acidity In: Sparks, DL Ed, Methods of soil analysis: part 3 Chemical
616 Methods, Book Series No5. SSSA and ASA, Madison, WI. 1996; 475-489.
617 <https://doi.org/10.2136/sssabookser5.3.c16>

- 618 43. Walkley A, Black I A. An examination of the Degtjareff method for determining soil organic matter and
619 a proposed modification of the chromic acid titration. *Soil Sci.* 1934; 37, 29-38.
- 620 44. Sherrod LA, Dunn G, Peterson GA, Kolberg RL . Inorganic carbon analysis by modified pressure-
621 calcimeter method. *Soil Sci Soc Am J.* 2002; 66: 299 – 305. <https://doi.org/10.2136/sssaj2002.2990>
- 622 45. Bray RH, Kurtz LT. Determination of total organic and available forms of phosphorus in soils. *Soil Sci.*
623 1945; 59, 39-45.
- 624 46. Gee, GW, Bauder, JW . Particle size analysis In: Klute, A Ed *Methods of soil analysis, part 1 Physical*
625 *and mineralogical methods Agronomy Monograph 9 2nd Edition.* ASA/SSSA, Madison, WI. 1986 383-
626 411. <https://doi.org/10.2136/sssabookser5.1.2ed.c15>
- 627 47. Bremmer JM. Determination of nitrogen in soil by Kjeldahl method. *J Agri Sci.* 1960; 55, 11-33.
- 628 48. Burton JD, Riley JP. Determination of soluble phosphate, total phosphorus in sea water and of total
629 phosphorus in marine muds. *Microchimica Acta.* 1956. 44; 1350-1365.
630 <https://doi.org/10.1007/BF01223539>
- 631 49. Thomas GW. Exchangeable cations. *Methods of soil analysis, part 2, chemical and microbiological*
632 *properties.* Second Edition. Page AL (editor). Agron, ASA, SSSA, Madison, WI. 1982; 159-165
- 633 50. EU Nitrogen Expert Panel. Nitrogen use efficiency (NUE)- an indicator for the utilization of nitrogen in
634 agriculture and food systems. Wageningen University, Netherlands.2015. [www.eunep.com/wp-](http://www.eunep.com/wp-content/uploads/2017/03/Report-NUE-Indicator-Nitrogen-Expert-Panel-18-12-2015.pdf)
635 [content/uploads/2017/03/Report-NUE-Indicator-Nitrogen-Expert-Panel-18-12-2015.pdf](http://www.eunep.com/wp-content/uploads/2017/03/Report-NUE-Indicator-Nitrogen-Expert-Panel-18-12-2015.pdf)
- 636 51. FAO. The importance of soil organic matter : Key to drought-resistant soil and sustained food
637 production. Eds. Bot, A. and Benites, J. 2005; www.fao.org/3/a0100e/a0100e00.htm
- 638 52. Adugna G. A review on the impact of compost on soil properties, water use and crop productivity.
639 *Agric Sci Res J.* 2018; 4(3): 93-104. <https://doi.org/10.14662/ARJASR2016010> .
- 640 53. Rivero C, Chirenje T, Ma LQ, Martinez G. Influence of compost on soil organic matter quality under
641 tropical conditions. *Geoderma*, 2004; 123 (3-4), 355-361. <https://doi.org/10.1016/j.geoderma.2004.03.002>
- 642 54. McGiffen M, Lebron I, Nguoajo M, Hutchinson CM. Soil organic amendments change low organic
643 matter agroecosystems. *Acta Horticulturae.* 2004; <https://doi.org/10.17660/ActaHortic.2004.638.32>
- 644 55. Bogati K, Walczak M. The impact of drought stress on soil microbial community, enzyme activities and
645 plants. *Agron.*2022; 12, 189. <https://doi.org/10.3390/agronomy12010189>
- 646 56. Leal FT, Filla VA, Bettol JVT, Sandrini, F de O FT, Mingotte FLC, Lemos LB. Use efficiency and
647 responsivity to nitrogen of common bean cultivars. *Agric Sci.* 2019; 43. [https://doi.org/10.1590/1413-](https://doi.org/10.1590/1413-7054201943004919)
648 [7054201943004919](https://doi.org/10.1590/1413-7054201943004919)
- 649 57. Shumi D, Alemayehu D, Afeta T, Debelo B. Response of common bean (*Phaseolus vulgaris* L.)
650 varieties to rates of blended NPS fertilizer in Adola District, Southern Ethiopia. *J Plant Biol Soil Health.*
651 2018; 5(1), 11. [www.avensonline.org/wp-content/uploaods/JPBSSH-2331-8996-05-0022.pdf](http://www.avensonline.org/wp-content/uploads/JPBSSH-2331-8996-05-0022.pdf)
- 652 58. Peoples MB, Hauggaard-Nielsen H, Jensen ES. The potential environmental benefits and risks
653 derived from legumes in rotations. In: *Nitrogen fixation in crop production.* Eds. Emerich, D.W., Krishnah,
654 H.B. Madison, USA; ASA,CSSA,SSSA, 2009; 349-385
- 655 59. Hardardson G. Methods for enhancing symbiotic nitrogen fixation. *Plant Soil.* 2004; 152, 1-17.
656 <https://doi.org/10.1007/BF00016329>
- 657 60. Rodino AP, Riveiro M, De Ron AM. Implications of the symbiotic nitrogen fixation in common bean
658 under seasonal water stress. *Agron.* 2021; 11, 70. <https://doi.org/10.3390/agronomy11010070>

659 61. Reinprecht Y, Schram L, Marsolais F, Smith TH, Hill B, Pauls KP. Effects of nitrogen application on
660 nitrogen fixation in common bean production. *Front Plant Sci Sec: Plant Breeding*. 2020;
661 <https://doi.org/10.3389/fpls.2020.01172>

662 62. Brust GE. Management strategies for organic vegetable fertility. In: *Safety and practice for organic*
663 *food*. Science Direct. 2019; ISBN 978-0-12-812060-6, 397-408. <https://doi.org/10.1016/C2016-0-02314-8>

664 63. Palm CA, Gachengo CN, Delve RJ, Cardish RJ, Giller K. Organic inputs for soil fertility management
665 in tropical agroecosystems. *Agric Ecosys Environ*, Elsevier. 2001; 83, 27-42.
666 <https://doi.org/10.3390/agronomy8070120>

667 64. Ajaweed AN, Hassan FM, Hyder NH. Evaluation of physio-chemical characteristics of bio-fertilizer
668 produced from organic solid waste using compost bins. *Sustainability*, 2022; 14, 4738.
669 <https://doi.org/10.3390/su14084738>

670 65. Khater E. Some physical and chemical properties of compost. *Int J Waste Res*. 2015; 5(1).
671 <https://doi.org/10.4172/2252-5211.1000172>

672 66. Mangan F, Barker A, Bodine S, Borten P. Compost use and soil fertility. University of Massachusets
673 Extension. 2013. www.ag.umass.edu/vegetable/fact-sheets/compost-use-soil-fertility

674 67. Sullivan D, Bary AI, Miller RO, Brewer LJ. Interpreting compost analyses. Oregon State University
675 Extension Service. 2018.
676 www.catalog.extension.oregonstate.edu/sites/catalog/files/project/pdf/em9217.pdf

677 68. Kreiling A. How does uncured compost harm plants. The Solana Center Composter.
678 2012. www.solanacompost.wordpress.com/2012/02/18/how-does-uncured-compost-harm-plants/

679 69. Beebe SE, Rao IM, Blair MW, Acosta-Gallegos JA. *Frontiers in Physiology*. 2013.
680 <https://doi.org/10.3389/fphys.2013.000035>

681 70. Nurudeen AR, Larbi A, Kotu BK, Tetteh FM, Hoeschle-Zeledon I. Does nitrogen matter for legumes?
682 Starter nitrogen effects on biological and economic benefits of cowpea (*Vigna unguiculata* L.) in Guinea
683 and Sudan Savannah of West Africa. *Agron*. 2018; 8(7): 1-12. <https://doi.org/10.3390/agronomy8070120>

684 71. Kakiuchi J, Kamiji Y. Relationship between phosphorus accumulation and dry matter partitioning in
685 soybeans. *Plant Prod Sci*. 2015; 18:3, 344 – 355. <https://doi.org/10.1626/pps.18.344>

687 72. Habete A, Buraka T. Effect of Rhizobium inoculation and nitrogen fertilization on nodulation and yield
688 response of common bean (*Phaseolus vulgaris* L.) at Boloso Sore, Southern Ethiopia. *J Biol, Agric*
689 *Health*. 2016; 6(13). www.core.ac.uk/download/pdf/234662059.pdf

690 73. Hopkins M. 4 Key factors affecting potassium uptake. University of Minnesota Extension Soil
691 Scientists. 2016. www.croplife.com/crop-inputs/fertilizer/4-key-factors-affecting-potassium-uptake

692 74. Mendes W da C, Junior JA, da Cunha PCR, da Silva A R, Evangelista AWP, Casaroli D. Potassium
693 leaching in different soils as a function of irrigation depths. *Soil, water & Plant Manage*. 2016; 20(11),
694 972-977. <https://doi.org/10.1590/1807-1929/agriambi.v20n11p972-977>

695 75. Dogan N, Akinci S. Effects of water stress on the uptake of nutrients by bean seedlings (*Phaseolus*
696 *vulgaris* L.). *Fresenius Environ Bulletin*. 2011; 20(8), 2163-2173.
697 www.researchgate.net/publication/287464271_Effects_of_water_stress_on_the_uptake_of_nutrients_by_bean_seedlings_phaseolus_vulgaris_L

699 76. Hatfield JL, Dold C. Water-use efficiency: advances and challenges in a changing climate. *Front Plant*
700 *Sci, Sec: Plant Physiology*. 2019. <https://doi.org/10.3389/fpls.2019.00103>

701 77. Waraich EA, Ahmad R, Ashraf MY, Saifullah, Ahmad M. Improving agricultural water use efficiency
702 by nutrient management in crop plants. *Acta Agriculturae Scandinavica, Section B- Soil & amp, Plant Sci.*
703 2011; 61(4), 291-304. <https://doi.org/10.1080/09064710.2010.491954>

704 78. Dianatmanesh M, Kazemeini SA, Bahrani MJ, Shakeri E, Alima M, Amjad SF, Mansoor N, Poczai P,
705 Lalarukh I, Abbas MHH, Abdelhafez AA, Hamed MH. Yield and yield components of common bean as
706 influenced by wheat residue and nitrogen rates under water deficit conditions. *Environ Tech & Inno.* 2022;
707 28, 102549. <https://doi.org/10.1016/j.eti.2022.102549>

708

709