

# Occurrence of Heavy Metal and Antibiotic Resistant Bacteria in Soils from Selected Mechanic Workshops in Ibadan Metropolis, Nigeria

### ABSTRACT

**Aim:** This study investigated the occurrence of antibiotic and heavy metal co - resistance in bacteria indigenous to mechanic workshops.

**Study design:** The study is an experimental study

**Place:** Samples were collected at mechanic workshop

**Methodology:** Soil samples were collected from three mechanic workshops and tested for the presence of heavy metals. Hydrocarbon-utilizing bacteria was also isolated from the soils using Bushnell Haas medium supplemented with 1% engine oil or petrol. Isolates obtained were subjected to antibiotic resistance test and those that showed extensive drug resistance to the tested antibiotics were further tested for heavy metal resistance. Selected isolates were then identified using 16S rRNA gene sequence.

**Results:** The soil samples contained excessive amounts of Lead, Iron, Copper and Zinc, while the concentrations of Nickel, Cobalt, Cadmium and Chromium were within WHO permissible limit. A total of 10 hydrocarbon-degrading isolates were obtained from the soil samples, seven of which were gram negative and three were gram positive. Three of the isolates showed extensive drug resistance to 14 of the tested antibiotics. three isolates were then subjected to heavy metal resistance test and all of them showed resistance to the tested heavy metals. They were identified and given ascension numbers as *Stenotrophomonas maltophilia* MW392903, *Achromobacterxylooxidans* MW392904 and *Pseudomonas aeruginosa* MW392905.

**Conclusion:** It was observed that heavy metal resistance and antibiotic resistance can be selected for simultaneously as organisms adapt ways to cope with man's activities in the environment, and while traits like hydrocarbon utilization and heavy metal resistance make the organisms promising in bioremediation, the inadvertent possession of antibiotic resistance by these environmental isolates poses a challenge to the health sector.

**Keywords:** Heavy metals resistance, Antibiotic resistance, *Stenotrophomona smaltophilia*, *Achromobacterxylooxidans*, *Pseudomonas aeruginosa*

## 1. INTRODUCTION

Heavy metals are chemical elements found in nature and components of the earth's crust that have a specific gravity at least five times that of water<sup>1</sup>. Arsenic (5.7), cadmium (8.65), iron (7.9), lead (11.34), and mercury (13.546) are some well-known metallic metals with a specific gravity greater than or equal to 5<sup>2</sup>. They are regularly recycled from one earth compartment to the next due to natural processes such as weathering, erosion, and biological activity since they are non-degradable in the environment. Heavy metals have been known to accumulate to harmful or toxic amounts as a result of substantial industrialization and urbanization, as well as the rise of indiscriminate dumping of untreated industrial wastewater and municipal sewage into the environment<sup>5</sup>.

Heavy metal levels in soil have been linked to poor plant development, lower water and nutrient uptake, and a variety of enzymatic disruptions<sup>6</sup>. Heavy metal contamination puts food safety in jeopardy. Because of the high level of heavy metal contamination in China, the food safety conference in 2009 showed that one sixth of the cultivated land was contaminated by heavy metals, and an area of more than 20 million hectares was threatened with being barred from agricultural practices<sup>7,8</sup>. Heavy metal uptake and accumulation have also been found in edible vegetables, grains, and seafood, according to several studies<sup>9</sup>. Heavy metal accumulation in human tissues and organs has resulted in cardiovascular, kidney, and neurological system disorders, as well as bone diseases<sup>10,11</sup>. To address the problem of heavy metal pollution in the environment, various strategies have been developed. Chemical precipitation and solvent extraction have been used as physicochemical methods<sup>12</sup>. These approaches, on the other hand, have shown to be difficult to apply on big surfaces, expensive, and ecologically unfriendly, as they may necessitate the use of harmful reagents in the restoration process<sup>12,13</sup>.

Although large levels of heavy metals have a deleterious influence on microbial populations in the impacted environment, some bacteria may tolerate or even thrive in the presence of specific metals<sup>14</sup>. Bacterial tolerance to heavy metals is defined as the bacteria's ability to cope with metal toxicity through intrinsic properties, whereas bacteria resistance is defined as the bacteria's ability to survive in higher concentrations of toxic metals through detoxifying mechanisms activated by the presence of the specific metals<sup>4</sup>. Heavy metal resistant bacteria are a superior option for heavy metal decontamination, and they have already been used successfully in the developed world<sup>10,15</sup>. However, there are few research in Nigeria looking into the possible use of metal-resistant bacteria for bioremediation<sup>16</sup>.

The rise in the number of bacteria resistant to antimicrobial agents puts the control of infectious diseases at jeopardy. This is due to the fact that infections produced by resistant germs frequently fail to react to standard treatment, resulting in extended sickness and an increased risk of mortality. Antibiotic resistance is a type of drug resistance in which a microorganism can survive antibiotic exposure. Antibiotic resistance is primarily caused by bacterial genetic mutation<sup>17</sup>. Antimicrobial drugs are used inappropriately and irrationally, creating favorable conditions for resistant microbes to arise, spread, and persist. Regardless matter the severity of the need for the antibiotic, the longer it is exposed to the environment, the greater the danger of resistance developing. As antibiotic resistance grows increasingly frequent, there is a larger need for alternative treatments. Despite the search for new antibiotic medicines, the number of newly approved antibiotics has continued to decline<sup>18,19</sup>. As a result, antibiotic resistance is a major issue.

In a heavy metal contaminated ecosystem, heavy metal resistance and antibiotic resistance can be selected for at the same time, and this co-selection has also been recorded in agriculture, animal husbandry, waste water treatment systems, and sediments<sup>10</sup>. Organisms possessing this dual feature could be useful in bioremediation of metal-polluted ecosystems, as well as in overcoming the inhibitory effect heavy metals have on the biodegradation of organic contaminants. Furthermore, in a contaminated environment, such dual resistance organisms might be able to compete effectively with antibiotic-producing flora.

## **2. MATERIALS AND METHODS**

### **2.1 Soil Samples Collection**

Soil samples were collected from three different mechanic workshops at sites that showed heavy oil activity. A total of three samples were collected from each workshop. Sample 1 was collected at Sango, 7° 25' 47.1" N, 3° 53' 31.2" E, Sample 2 at Ologuneru situated between 7° 25' 04.0" N, 3° 51' 23.1" E and Sample 3 at Orogun situated between 7° 27' 08.0" N, 3° 54' 44.1" E. Each of these areas are located within Ibadan metropolis, Oyo State, Nigeria. The soil samples were collected in clean polythene bags and transported to the laboratory. The control sample was collected from the Botanical Garden, University of Ibadan, where there are no anthropogenic activities such as car repairs, commercial activities, and drainage influence.

## 2.2 Equipment

Beakers (Pyrex), Conical flasks (Pyrex), Autoclave, Petri-dishes, Forceps, Cotton wool, Spirit lamps, McCartney bottles, Mortar and Pestle, Filter Paper, Measuring Cylinders, Refrigerator, Spatula, Test tubes, Weighing Balance SUN-5202, Incubator, UV Spectrophotometer V-730, Petrol, Engine Oil, Inoculating Loop, Antibiotic discs, Cock borer, Hand gloves, Nose masks, Compound Microscope, Slides, Standard Erlenmeyer flask, 0.5mm Sieve, Bulk Scientific 210/211 VGP Atomic absorption spectrophotometer.

## 2.3 Chemicals/ Reagents/ media used

Crystal Violet, Lugol's Iodine, Immersion Oil, Safranin, Hydrogen peroxide, Nitric Acid, Chloric Acid, Ethanol, Polyacrylamide Gel, Zinc chloride, Lead sulphate, Copper sulphate, Iron sulphate, De-ionized distilled water, Bushnell-Haas medium, Nutrient Agar, Bacteriological Agar, Nutrient Broth, Mueller-Hinton Agar.

## 2.4 Analysis of Heavy Metals in Soil

Soil samples were homogenized and gently crushed multiple times with a mortar and pestle before being passed through a 0.5mm mesh for analysis. 15mL of freshly made aqua regia (HNO<sub>3</sub>: HCl, 2:1) was added to a homogenized sample (0.5g) in a standard Erlenmeyer flask. The contents of the beaker were heated on a digestion block while it was covered. In a dessicator, the mixture was cooled before being filtered through a Whatman No. 42 filter paper into a 25mL standard volumetric flask. The filtrate was diluted to 25mL with de-ionized distilled water<sup>20</sup>. Blank solutions were also prepared using aqua regia and de-ionized water. The digested sample solutions were analysed using the atomic absorption spectrometer {Buck Scientific 210/211 VGP Atomic absorption spectrophotometer (AAS)}. Standard solutions of the various heavy metals were analysed. All soil samples were analysed in triplicate to minimize error.

## 2.5 Enrichment and Isolation of Hydrocarbon Degrading Bacteria

One gram of soil sample was taken from each of the soil samples and serial dilution was carried out to the tenth dilution factor. Bushnell-Haas medium (BH)<sup>21</sup> was used for the isolation of hydrocarbon-degrading bacteria from the soil. The BH medium consist of the following composition in g/L; 0.2 MgSO<sub>4</sub>, 0.02 CaCl<sub>2</sub>, 1.0 K<sub>2</sub>HPO<sub>4</sub>, 1.0 KH<sub>2</sub>PO<sub>4</sub>, 1.0 NH<sub>4</sub>NO<sub>3</sub>, 0.05 FeCl<sub>3</sub>, pH 7.0. BH was supplemented with 1% (v/v) engine oil and/or petrol as the sole carbon source. Three milliliter of dilution factors 10<sup>-3</sup>, 10<sup>-7</sup> and 10<sup>-9</sup> were added to 250 mL Erlenmeyer flasks containing 100 mL sterile BH medium supplemented with engine oil or petrol and the flasks were incubated at 30 °C for 7 days on rotary shaker (150 rpm). After incubation, 0.1 mL of culture was plated on solid BH medium supplemented with 1% (v/v) engine oil and/or petrol as the sole carbon source using the spread plate method and incubated for 48 hours. Colonies that grew were sub-cultured till pure cultures were obtained, which were then maintained on Nutrient Agar prepared according Manufacturer's instruction.

## 2.6 Characterization of Isolated Bacteria

Isolates were characterized using conventional morphological methods.

### 2.6.1 Gram's Staining

Twenty four-hour old culture was prepared. A smear was made by placing a drop of water on the slide and then transferring the microbial inoculum to the drop of water with a sterile cooled loop, it was then mixed and spread by means of a circular motion of the inoculating loop. The smear was air dried and heat fixed and was gently flooded with crystal violet for 1 minute and washed under running water. The smear was gently treated with three drops of Lugol's iodine and left for one minute and gently washed with running water again. The smear was decolorized with 95% concentration ethyl alcohol reagent and was

gently washed under running water. Counterstaining with safranin was carried out after which the slide was dried using a filter paper and examined under the microscope using the x100 objective lens with oil immersion.

## **2.7 Antibiotic Susceptibility Testing of Isolated Bacteria**

Single disc diffusion method<sup>22</sup> was used to examine isolates susceptibility to antimicrobial agents. Mueller-Hinton Agar (MHA) was used for this test and microbial sample preparations were carried out using 18-24 hour old culture which was adjusted to 0.5 Mc Farland Standard corresponding to  $1.5 \times 10^8$  CFU/ML. A sterile swab stick was used to spread the culture standard on the whole surface of the MHA plate after which the antibiotic discs were placed on the MHA plate with the aid of a sterile forceps. Incubation was subsequently done for 24 hours. The antibiotic sensitivity discs utilized were; streptomycin (30µg), septrin (15µg), erythromycin (15µg), zinnacef (10µg), amoxicillin (30µg), rocephin (10µg), ciprofloxacin (30µg), pefloxacin (10µg), gentamicin (30µg), ofloxacin (10µg), chloramphenicol (10µg), sparfloxacin (30µg), augmentin (10µg), ampiclox (10µg). The diameter of the inhibitory zones around the respective antibiotic discs was then measured with the aid of a meter rule<sup>22</sup>.

## **2.8 Heavy Metal Resistance (HMR) test on Selected Isolates**

Mueller-Hinton Agar (MHA) was the culture medium used and 18-24 hour old culture was streaked on the plate with the aid of a sterile inoculating loop. A cork borer was then used to bore wells in the MHA streaked plate. The metals  $Pb^{2+}$ ,  $Cu^{2+}$ ,  $Fe^{2+}$ , and  $Zn^{2+}$  were used as  $PbCl_2$ ,  $CuSO_4$ ,  $FeCl_2$  and  $ZnCl_2$  salts, respectively. Stock solutions of the metals were prepared in six varying concentrations (1g/ml, 0.1g/ml, 0.01g/ml, 0.001g/ml, 0.0001g/ml and 0.00001g/ml) and 1mL each of the solution were placed in the bored holes. The control experiment was carried out by inoculating the pure isolates on basal media without the heavy metals. The results were read after 24hours<sup>23</sup>.

## **2.9 Molecular Identification**

The 16S ribosomal RNA (rRNA) gene was analyzed to identify the selected bacteria strains. After the cells were cultured overnight in Nutrient Broth, the genomic DNA was extracted using ZR Bacterial DNA Miniprep. The 16S rRNA gene fragment of strains was then amplified by PCR amplification using forward primer >16S-27F(5'-AGAGTTTGATCMTGGCTAG-3') and reverse primer >16S-1492R (5'-CGGTTACCTTGTTACGATT-3'). The 16S rRNA PCR product was sequenced by 3130XL genetic analyser for Applied Biosystems. The Isolate sequences gotten after the polymerase chain reaction were opened with Bio-Edit and the sequences obtained were compared with data available in GenBank database using the Mega 6.0 Blast network service of the National Centre for Biotechnology Information (NCBI)<sup>24</sup>. A phylogenetic tree was constructed in MEGA 6.0 software by carrying out the multiple alignments using Jukes-Cantor method. Bootstrap values were analyzed based on 1000 replications.

## **3. RESULTS**

### **3.1 Analysis of Heavy Metals present in the Soil Samples**

The soil samples were analyzed for a total of 8 heavy metals namely Iron (Fe), Copper (Cu), Zinc (Zn), Lead (Pb), Cadmium (Cd), Chromium (Cr), Nickel (Ni) and Cobalt (Co). Iron and Cadmium had the highest and lowest concentrations respectively for each of the soil samples. The soil sample obtained from the mechanic workshop in Orogun (Sample 3) had the highest Fe concentration while that of Sango (Sample 1) had the least Fe concentration as seen in Table 1.

### **3.2 Characterization of Isolated Bacteria**

#### **3.2.1 Gram Staining Result and Pigments of Isolated Bacteria**

Ten (10) bacteria were isolated from this study of which 7 were gram negative and 3 were gram positive while out of the ten (10) isolates, nine (9) were cream in colour and one (1) was green in colour.

### 3.3 Antibiotic Susceptibility test of Isolated Bacteria

Most of the isolates showed multiple drug resistance. However, three (3) of them 27<sup>B</sup>A<sup>2</sup>, 23<sup>B</sup>A<sup>2</sup> and 17<sup>B</sup>A<sup>1</sup> showed extensive drug resistance as seen in Table 2 and 3.

### 3.4 Heavy metal resistance (HMR) test on selected isolates

The result showed that the organisms were resistant to the heavy metals. At higher concentrations of 1g/ml and 0.1g/ml, there were visible precipitates of the heavy metals around the cultures but none at lower concentrations as seen in Plate 1 to 3.






Plate 1: Plate HMR-1 showing the visible precipitates of Lead (all in g/ml) around the culture



0.00001


**Plate 2: Plate HMR-2 showing the visible precipitates of Iron (all in g/ml) around the culture**



**Plate 3: Plate HMR-3 showing the visible precipitates of Copper (all in g/ml) around the culture**

### 3.5 Molecular Identification of Bacteria with desired traits

Isolates coded  $17^B A^1$ ,  $23^B A^2$  and  $27^B A^2$  were identified as *Stenotrophomonas maltophilia*, *Achromobacter xylosoxidans* and *Pseudomonas aeruginosa*, respectively after subjecting their isolated DNA to 16S ribosomal RNA gene amplification and subsequent sequencing of the amplified genes. *Stenotrophomonas maltophilia*, *Achromobacter xylosoxidans* and *Pseudomonas aeruginosa* has a 98.67%, 97.99% and 97.17% identity respectively using BLAST 2.10.0N+. The 16S ribosomal RNA gene of each of the isolates was submitted to GenBank and each was assigned the following accession numbers: *Stenotrophomonas maltophilia*: MW392903, *Achromobacter xylosoxidans*: MW392904 and *Pseudomonas aeruginosa*: MW392905. The sizes of the amplicons were about 1,428bp for isolate  $17^B A^1$ , 1,442bp for isolate  $23^B A^2$  and 1,413bp for isolate  $27^B A^2$  and the DNA ladder used was Hyper ladder 1. The gel image obtained after running the amplified 16S rRNA gene of the three isolates with the molecular marker is shown in Fig 4.1.




**Fig.1: Gel image for PCR of the three isolates with molecular marker/ladder.**


Source: *Laboratory Analysis, 2020*

KEY: M= Molecular marker/ladder  
1= *Stenotrophomonas maltophilia*,  
2= *Achromobacter xylosoxidans*  
3= *Pseudomonas aeruginosa*


### **3.6 Phylogenetic Analysis of Bacteria with desired traits**



**Fig.2. Neighbor-joining phylogenetic tree based on 16S rRNA gene sequences showing *Stenotrophomonas maltophilia* found in the GenBank Database. The bar indicates 0.005 substitutions per nucleotide position and values at nod represent percentage of 1000 bootstrap replicates.**



**Fig. 3. Neighbor-joining phylogenetic tree based on 16S rRNA gene sequences showing *Achromobacter xylosoxidans* found in the GenBank Database. The bar indicates 0.1 substitutions per nucleotide position and values at nod represent percentage of 1000 bootstrap replicates.**



**Fig. 4. Neighbor-joining phylogenetic tree based on 16S rRNA gene sequences showing *Pseudomonas aeruginosa* found in the GenBank Database. The bar indicates 0.005 substitutions per nucleotide position and values at nod represent percentage of 1000 bootstrap replicates.**

**Table 1: Concentrations of the various heavy metals found in the soil samples**

Sample No	g/kg Fe	g/kg Cu	g/kg Zn	g/kg Co	g/kg Cr	g/kg Cd	g/kg Pb	g/kg Ni
1	20.51±1.50	0.27±0.01	0.51±0.01	0.01±0.00	0.06±0.00	0.000	0.20±0.00	0.03±0.00
2	21.34±0.50	0.05±0.00 <sub>a</sub>	0.34±0.03 <sub>a</sub>	0.01±0.00	0.04±0.00	0.000	0.03±0.00	0.01±0.00
3	32.27±0.70 <sub>a,b</sub>	0.11±0.00 <sub>a,b</sub>	1.42±0.03 <sub>a,b</sub>	0.02±0.00	0.05±0.00	0.000	0.22±0.00	0.03±0.00

Values are presented as mean ± standard deviation. <sup>a</sup>value is significant when compared with sample 1 <sup>b</sup>value is significant when compared with sample 2

**Table 2: Antibiotic susceptibility test of Isolated Bacteria**

Name of Antibiotics	33 <sup>B</sup> T	29 <sup>B</sup> A <sup>1</sup>	17 <sup>B</sup> A <sup>2</sup>	33 <sup>A</sup>	23 <sup>B</sup> A <sup>1</sup>	27 <sup>B</sup> A <sup>2</sup>	29 <sup>B</sup> A <sup>2</sup>	17 <sup>A</sup>	23 <sup>B</sup> A <sup>2</sup>	17 <sup>B</sup> A <sup>1</sup>
<b>CN</b>	-ve	-ve	-ve	16	-ve	<b>-ve</b>	-ve	-ve	<b>-ve</b>	<b>-ve</b>
<b>PEF</b>	15	15	15	17	15	<b>-ve</b>	13	16	<b>14</b>	<b>12</b>
<b>OFX</b>	11	11	13	17	12	<b>-ve</b>	-ve	20	<b>12</b>	<b>11</b>
<b>S</b>	-ve	-ve	-ve	16	-ve	<b>-ve</b>	17	15	<b>-ve</b>	<b>-ve</b>
<b>SXT</b>	11	-ve	10	15	10	<b>-ve</b>	17	15	<b>-ve</b>	<b>-ve</b>
<b>CH</b>	12	-ve	-ve	13	-ve	<b>-ve</b>	-ve	-ve	<b>-ve</b>	<b>-ve</b>
<b>SP</b>	-ve	-ve	-ve	17	-ve	<b>-ve</b>	17	-ve	<b>-ve</b>	<b>-ve</b>
<b>CPX</b>	24	22	21	17	22	<b>-ve</b>	22	20	<b>-ve</b>	<b>-ve</b>
<b>AM</b>	-ve	-ve	-ve	17	-ve	<b>-ve</b>	-ve	-ve	<b>-ve</b>	<b>-ve</b>
<b>AU</b>	-ve	-ve	-ve	13	-ve	<b>-ve</b>	-ve	-ve	<b>-ve</b>	<b>-ve</b>

**AU-** augmentin **CN-** gentamycin **PEF-** perfloxacin **OFX-** tarivid **S-** streptomycin **SXT-** septrin  
**CH-** chloramphenicol **SP-** sparfloracin **CPX-** ciprofloxacin  
The diameters of susceptibility are measured in millimeter (mm). -ve means Negative

**Table 3: Antibiotic susceptibility test of Isolated Bacteria**

Name of Antibiotics	33 <sup>B</sup> T	29 <sup>B</sup> A <sup>1</sup>	17 <sup>B</sup> A <sup>2</sup>	33 <sup>A</sup>	23 <sup>B</sup> A <sup>1</sup>	27 <sup>B</sup> A <sup>2</sup>	29 <sup>B</sup> A <sup>2</sup>	17 <sup>A</sup>	23 <sup>B</sup> A <sup>2</sup>	17 <sup>B</sup> A <sup>1</sup>
<b>CN</b>	15	-ve	-ve	15	-ve	<b>-ve</b>	-ve	-ve	<b>-ve</b>	<b>-ve</b>
<b>PEF</b>	16	-ve	-ve	17	16	<b>-ve</b>	12	-ve	<b>15</b>	<b>-ve</b>
<b>E</b>	-ve	-ve	-ve	-ve	-ve	<b>-ve</b>	-ve	-ve	<b>-ve</b>	<b>-ve</b>
<b>S</b>	15	-ve	-ve	17	21	<b>-ve</b>	-ve	15	<b>-ve</b>	<b>-ve</b>
<b>SXT</b>	-ve	-ve	-ve	12	-ve	<b>-ve</b>	-ve	18	<b>-ve</b>	<b>-ve</b>
<b>Z</b>	-ve	-ve	-ve	-ve	-ve	<b>-ve</b>	-ve	-ve	<b>-ve</b>	<b>-ve</b>
<b>R</b>	-ve	-ve	-ve	20	-ve	<b>-ve</b>	-ve	-ve	<b>-ve</b>	<b>-ve</b>
<b>CPX</b>	20	16	22	18	20	<b>-ve</b>	21	20	<b>-ve</b>	<b>-ve</b>
<b>AM</b>	-ve	-ve	-ve	-ve	-ve	<b>-ve</b>	-ve	-ve	<b>-ve</b>	<b>-ve</b>
<b>APX</b>	-ve	-ve	-ve	-ve	-ve	<b>-ve</b>	-ve	-ve	<b>-ve</b>	<b>-ve</b>

*R- rocephin E- erythromycin CPX- ciprofloxacin S- streptomycin SXT- septrin PEF- perfloxacin CN- gentamycin APX- ampiclox Z- zinacef AM- amoxicillin*

*The diameters of susceptibility are measured in millimeter (mm). -ve means Negative*

#### 4. DISCUSSION

Using one-way Anova, the concentrations of heavy metals present in the three soil samples were compared relative to one another. The mean values of Fe (Iron), Cu (Copper) and Zn (Zinc) were significant when compared to one another while that of Co (Cobalt), Cr (Chromium), Cd (Cadmium), Pb (Lead) and Ni (Nickel) were insignificant.

Iron (Fe) had the greatest mean concentration of all the metals studied, ranging from 20.5 to 32.3 grams per kilogram. Several studies have found high quantities of iron in soils when compared to other metals, confirming that natural soils contain considerable amounts of iron, though not as much as soils where anthropogenic activities may have played a role<sup>25,26,27,28</sup>. The mean iron value found in this study is higher than that found by other researchers.<sup>28,29,30,31,32</sup> The soil samples from the mechanic workshop in Orogun had the highest quantity of iron. The high values are likely due to the rusting of ancient car bodywork, as well as welding and panel beating activities on the job sites. In humans, acute Fe exposure can cause vomiting, cardiac depression, and metabolic acidosis<sup>26</sup>.

Zinc contents range from 0.51 to 1.42 g/kg in soil samples. The range of results found in this study is higher than prior studies' findings<sup>28,29,33,34</sup>. The highest zinc concentrations were found in Orogun. The high zinc content of this sample can be attributed to the site's antiquity as well as its position along the roadside. The wear and tear of car bodywork with galvanized steel surfaces could be to blame for high zinc levels in roadside soil<sup>35,36</sup>. Zinc is also utilized in vehicle brake linings because of its heat conducting qualities, and it can be released via mechanical abrasion, engine oil combustion, and vehicle tyres<sup>37,38</sup>. The acceptable limit of zinc in soil, according to the DPR standard, is 0.14g/kg, and the maximum zinc content in this study is 1.42g/kg. This result is more than the maximum amount of zinc that can be found in the soil. According to the WHO, the acceptable limit is 0.050 g/kg.

Chromium (Cr) concentrations ranged from 0.04 to 0.06 g/kg, and were found to be highest in Sango. Values obtained are slightly higher than those reported by some authors<sup>34,35,36</sup>. According to one of them, the presence of chromium in roadside soil is linked to the usage of chromic plating on various automobile

parts to prevent corrosion. Chromium is carcinogenic, and people exposed to chromium-containing dust develop cancer of the respiratory organs<sup>36</sup>. The acceptable limit of Chromium in soil is 0.1g/kg, according to WHO and DPR standards, and the maximum chromium concentration in this study is 0.06g/kg. This number is lower than the chromium allowed limit in soil.

Lead (Pb) concentrations in all the sample sites were in the range of 0.03 to 0.22 g/kg. This range of values for lead is higher in some studies and also lower than those of some other studies<sup>32,38,39</sup>. The concentration of lead in the soil is likely to have been derived from vehicle exhaust fumes containing some lead-rich aerosols<sup>36</sup>. Tetraethyl lead is used as an anti-knocking compound in gasoline, according to studies, and it is released during automotive emissions and fossil fuel combustion<sup>40</sup>. According to one study, petrol engine exhaust emissions account for roughly 80% of all lead in the air<sup>38</sup>. Furthermore, lead levels in the roadside might be linked to the deterioration of car brake linings and tyres<sup>41</sup>. Lead poisoning poses a major threat to one's health<sup>41</sup>.

Nickel (Ni) values in soils from all sites varied from 0.01 to 0.03 grams per kilogram. This study's highest nickel value is higher than values published by several authors<sup>28,33,42</sup>. Airborne particles produced by brakes and tyre wear can include significant amounts of nickel, making this a likely source of nickel in the soils studied. The diesel used in automobiles<sup>35</sup> could also be a source of anthropogenic nickel input in the studied locations. Sango and Orogun had the highest Ni levels (0.03 g/kg). Exposure to high levels of nickel from plants grown in nickel-rich soil increases the risk of lung, nose, laryngeal, and prostate cancers, as well as respiratory failures, birth defects, and heart problems<sup>43</sup>. According to WHO standards, the maximum nickel concentration in soil is 0.035g/kg, while the maximum nickel concentration in this study is 0.03g/kg. This value is somewhat lower than the maximum amount of nickel that can be found in soil.

Copper (Cu) concentrations ranged from 0.05 to 0.27 (g/kg). This value is higher than the value reported by a previous researcher<sup>31</sup>. Copper, lead, and antimony are all present in high concentrations in used oils that leach into the earth as leachates. Copper can also be obtained via metal bearing wear. Although, copper is required for plant growth, but only in minute levels. An excess of copper ions causes headaches, nausea, vomiting, and diarrhoea, and excessive quantities induce anemia, gastrointestinal disorders, and in extreme cases, liver and kidney failure. <sup>44</sup>. The acceptable limit of copper in soil is 0.036g/kg, according to WHO and DPR standards, and the maximum copper concentration in this study is 0.27g/kg. This value is higher than the permissible limit for copper in the soil.

Cobalt (Co) was equally detected in all the soil samples. The sample concentration ranges from 0.01 to 0.02 g/Kg. The highest concentration value was obtained at Orogun. The values are higher than the DPR permissible limit of Cobalt in soil.

In mechanic workshops, there is a constant change in the soil micro-organisms as a result of deliberate spillage of used engine oil and other petrochemical products. This alters the biomass and ecology of the soil such that microbial communities undergo a dynamic change in specie composition and grasses can no longer grow on the soil. The colour and texture of the soil are also affected<sup>22</sup>. Some studies have shown that, oil-polluted soils are dominated by Gram negative bacteria as was seen also in this research project<sup>46,47</sup>. The result of the present study revealed that soils of mechanic workshops may harbor hydrocarbon degraders that came about as a result of exposure to hydrocarbons due to indiscriminate disposal of the spent engine oil collected from the crankcase of motor vehicles, motor bikes and other machinery.

The distribution pattern of the hydrocarbon utilizing isolates gotten from this study showed that gram negative rods had the highest occurrence at 70%. This is close to a study which isolate occurrence was 80% <sup>48</sup>. In that same study, gram positive cocci also had the least (10%) which correlates to the value gotten in this research where gram positive cocci had the lowest percentage.

The antibiotic susceptibility profile of the test bacterial cultures revealed that all the bacterial isolates were also multi drug resistant. Heavy metal resistance and antibiotic resistance can be selected for simultaneously in a heavy metal contaminated ecosystem. This co-selection has also been reported in agriculture, animal husbandry, waste water treatment system and sediments<sup>49</sup>. According to some authors resistance genes to both antibiotics and heavy metals could be closely located on the same

plasmid in bacteria and are thus more likely to be transferred together in the environment<sup>50,51</sup>. None of the bacterial isolates in this study have plasmids, thus it could be that the microorganisms while developing other mechanisms to cope with excess heavy metals such as efflux channels to pump out excess heavy metals may have adapted the same process for resisting antibiotics<sup>52</sup>.

At 0.00001g/ml, 0.0001g/ml, 0.001g/ml, and 0.01g/ml, of Iron, zinc, copper and lead, all the bacteria isolated from the hydrocarbon polluted soil samples showed resistance to the heavy metals tested. However, at higher concentrations of 1g/ml and 0.1g/ml, there were visible precipitates of all the heavy metals around the culture. It has been observed that certain microorganisms are able to utilize heavy metals as electron donors during metabolism and are also able to precipitate heavy metals by reducing them to their insoluble forms such as chlorides or sulphates forms that are less bioactive<sup>52</sup>. At concentrations of 0.01g/ml, 0.001g/ml, 0.0001g/ml and 0.00001g/ml, the microorganisms could be able to utilize the heavy metals, as sources of electrons during metabolism. However, at higher concentrations of 1g/ml and 0.1g/ml, visible precipitates were seen around cultures which most likely means that the microorganisms were precipitating the heavy metals by reducing them to their insoluble forms. The exit of the metals from the cells could be via efflux pumps<sup>53</sup>.

The isolates exhibiting extensive drug resistance were identified as *Stenotrophomonas maltophilia*, *Achromobacter xylosoxidans* and *Pseudomonas aeruginosa* on the basis of their cultural, morphological and molecular characteristics. In polluted soil samples, similar findings of heavy metal resistance bacteria have been reported<sup>54,55</sup>. From a lagoon area, a group of researchers discovered *Pseudomonas* and *Streptococcus* sp. that were resistant to Cr and Pb<sup>54</sup>. From soil samples gathered from a mining site, another group discovered *Pseudomonas* sp. resistant to Pb<sup>55</sup>.

The selected heavy metal bacterial isolates in this investigation were shown to be resistant to a variety of antibiotics. *Stenotrophomonas maltophilia* had 92.9 percent resistance to the 14 antibiotics tested, *Achromobacter xylosoxidans* had 85.7 percent, and *Pseudomonas aeruginosa* had 100 percent resistance. *Pseudomonas aeruginosa* has also been found to have the highest resistance pattern<sup>56</sup>. Heavy metal and antibiotic resistance in bacteria have been linked in a number of studies<sup>56,57</sup>. Five heavy metal resistant bacteria strains, *Micrococcus* sp., *Nocardia* sp., *Acinetobacter junni*, *Actinomyces turicensis*, and *Pseudomonas aeruginosa*, were found to be resistant to all 18 antibiotics tested in a study by some researchers, which is similar to the findings of this study, in which *Pseudomonas aeruginosa* was resistant to all 14 antibiotics tested<sup>35</sup>. Metal and antibiotic resistance in bacteria most likely helps them adapt faster in the presence of high amounts of heavy metals in their environment by spreading resistant components. The organisms with antibiotic and heavy-metal resistance isolated and identified in this work could be useful in bioremediation of metal-polluted environments, as well as helping to overcome heavy metal inhibition of organic pollutant biodegradation. Furthermore, in a contaminated environment, such dual resistance organisms might be able to compete effectively with antibiotic-producing flora.

The environment is subject to anthropogenic pressures. Places like mechanic villages exert a significant amount of such pressure on the environment. Microbial communities are highly sensitive to such environmental changes and the excessive heavy metals as well as hydrocarbon pollution can impose selection pressures on soil microbes. Microbes that survive usually evolve a system that could either be based on biochemical or genetically encoded mechanisms or both in order to be metabolically active and grow. While these could help select for organisms that are useful for bioremediation, the downside is the inadvertent conferment of traits like antibiotic resistance which is a major source of concern to the health sector.

#### **COMPETING INTERESTS DISCLAIMER:**

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of

knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

## REFERENCES

- <sup>1</sup>Ali, S., Sardar, K., Hameed, S., Afzal, S., Fatima, S., Shakoor, B. M., Bharwana, S. A. and Tauqeer, H. M. Heavy metals contamination and what are the impacts on living organisms. *Greener Journal of Environmental Management and Public Safety* (2013)**2**: 172-179.
- <sup>2</sup>Oluwatosin, G. A., Adeoyolanu, O. D., Ojo, A. O., Are, K. S., Dauda, T. O. and Aduramigba-Modupe, V. O. Heavy metal uptake and accumulation by edible leafy vegetable (*Amaranthushybridus*L.) grown on urban valley bottom soils in southwestern Nigeria. *Soil and Sediment Contamination: (2010)* **19**: 1–20
- <sup>3</sup>Orisakwe, O. E., Asomuqua, R., Afonne, O. J., Anisi, S. N., Obi, E. and Dioka, C. E. Impact of effluents from a car battery manufacturing plant in Nigeria on water, soil, and food qualities. *Arch. Environ. Health* (2004) **59**: 31-36.
- <sup>4</sup> International Union of Pure and Applied Sciences (IUPAC)—Heavy metalsII- a meaningless term? *Pure Applied Chemistry* (2002)**74**: 793-807.
- <sup>5</sup> Adewole, M. B. and Uchegbu, L. U. Properties of soils and plants uptake within the vicinity of selected automobile workshops in Ile-Ife Southwestern, Nigeria. *Ethiopian Journal of Environ. Studies Manage.* (2010) **3**: 31-35.
- <sup>6</sup> Nordberg, G. F. and Nordberg, M. Biological monitoring of cadmium. In: Clarkson TW, Friberg L, Nordberg GF, Sager PR, eds. Biological monitoring of toxic metals. New York: Plenum Press. pp. (2001) 151-168.
- <sup>7</sup> Adie, G. U. and Osibanjo, O. Assessment of soil pollution by slag from an automobile battery manufacturing plant in Nigeria. *African Journal of Environmental Science and Technology* (2009)**3**: 239-250.
- <sup>8</sup>Ekundayo E. O. and Killham K. Lead solubilization and accumulation by two strains of *Pseudomonas* obtained from a contaminated Alfisol's effluent in Southwestern Nigeria. *Environmental Monitoring and Assessment* (2001)**71**: 243-254.
- <sup>9</sup>Olukoya D. K., Smith S. I. and M. O. Ilori Isolation and characterization of heavy metals resistant bacteria from Lagos Lagoon. *Folia Microbiol(Praha)*. (1997) **42**: 441-444.
- <sup>10</sup>Galadima, A., Okoronkwo, M. U., Mustapha, D. G., and Leke, L. Petrol in Nigeria: a fuel or a killer? —Is shift to hydroisomerisation not overdue? *Elixir International Journal* (2012)**43**: 6893-6897.
- <sup>11</sup> Nies, D. H., Microbial heavy-metal resistance. *Applied Microbiology and Biotechnology* (1999) **51**: 730-750.
- <sup>12</sup> Agarwal, S.K. Studies on the effect of the auto exhaust emission on the *Mitragynapatriflora*. Ajmeer, India: MDS University; Master Thesis.(1999)
- <sup>13</sup>Galadima, A., Muhammad, N. U. and Garba, Z. N. Spectroscopic investigation of heavy metals in waste water from University students' halls of residence. *International Journal of Chemistry*. (2010) **20**: 239-244.

- <sup>14</sup>. Yao, Z. T., Li, J. H., Xie, H. H., and Yu, C. H. Review on remediation technologies of soil contaminated by heavy metals. *Proc. Environ. Sci.* (2012) **16**: 722–729
- <sup>16</sup>. Kumar, A., Bisht, B. S. and Joshi, V. D. Bioremediation potential of three acclimated bacteria with reference to heavy metal removal from waste. *International Journal of Environmental Sciences* (2011)**2**: 896-908.
- <sup>17</sup>. R.Laxminarayan. Battling resistance to antibiotics and pesticides: an economic approach. Washington, DC: *Resources for the Future*.2003.
- <sup>18</sup>. J.A.Dromigny, J.D.Perrier-Gros-Claude. Antimicrobial resistance of *Salmonella enterica* serotype Typhi in Dakar, Senegal. *Clinical Infectious Disease*. 2003; **37**: 465–466.
- <sup>19</sup>. E.Zuccato,S.Castiglioni, R.Fanelli. Identification of the pharmaceuticals for human use contaminating the Italian aquatic environment. *Journal of Hazardous Materials*. 2005; **122** (3), 205-209.
- <sup>20</sup>. A.J,Adebayo, J.T.Jayoye, I.I.Oguntimehin and L.Lajide. Delineation of Heavy Metals in Soils from Auto-Mechanic Workshops within Okitipupa, Ondo State, Nigeria. *International Research Journal of Public and Environmental Health* 2017: 136-147
- <sup>21</sup>. S.D. Lima, A.F. Oliveira, R. Golin, V.C. Lopes, D.S. Caixeta, Z.M. Lima and E.B. Morais. Isolation and characterization of hydrocarbon degrading bacteria from gas station leaking contaminated ground water in Brazil. *Brazilian journal of biology* (2020), **2**: 354-361
- <sup>22</sup>. National Committee for Clinical Laboratory Standards (NCCLS). Methods for Dilution of Antimicrobial Susceptibility Tests for Bacteria that grow Aerobically 2004
- <sup>23</sup>. D.Kratochvil and B.Volesky. Advances in the biosorption of heavy metals. *Trends in Biotechnology*,2012 **16**:291-300.
- <sup>24</sup>.L.C.Trindade,E.Marques,D.B.Lopes,M.A.S.V.Ferreira. Development of a molecular method for detection and identification of *Xanthomonas campestris* sp. *viticola*. *Summa Phytopathologica*, 2007**33**(1):16-23
- <sup>25</sup>. Ademoroti C.M.A. Environmental Chemistry and Toxicology. Foludex Press Ltd Ibadan, (1996) pp.150-188
- <sup>26</sup>. Aluko OO, Oluwande PA.Characterization of leachates from a municipal solid waste landfill site in Ibadan, Nigeria. *J. Environ. Health Res.*, (2003) **2**: 83-84
- <sup>27</sup>. Oguntimehin I andpinmoroti K. Profile of Heavy Metals from Automobile Workshops in Akure Nigeria. *Journal of Environmental Science and Technology*. (2008) **1**:19-26
- <sup>28</sup>. Osakwe SA. Heavy metal contamination and physicochemical characteristics of soils from automobile workshops in Abraka, Delta State, Nigeria.*International Journal of Natural Science Research*(2014) **2**(4):48-58.
- <sup>29</sup>. Osakwe SA. Chemical speciation and mobility of some heavy metals in soils around automobile waste dumpsites in Northern part of Niger Delta, South Central Nigeria.*Journal of Applied Science and Environmental Management*.(2010) **14**(4):13-130
- <sup>30</sup>. Osakwe SA. Assessment of the effects of wood processing industries in selected parts of Delta State, Nigeria on the soils and vegetation in their vicinities. *IOSR Journal of Applied Chemistry*(2012)**3**:22-30.

- <sup>31</sup>. Nwachukwu MA, Feng H and Alinnor J. Assessment of Heavy Metals Pollution in Soils and their Implications within and Around Mechanic Villages. *International Journal of Environmental Science and Technology*. (2011)7(2):347- 358.
- <sup>32</sup>. Nwachukwu MA, Ntesat B and Mbaneme FC. Assessment of direct soil pollution in automobile junk market. *Journal of Environmental Chemistry and Ecotoxicology*. (2013)
- <sup>33</sup>. Idugboe SO, Tawari-Fufeyin P Midonu AA. Soil pollution in two auto mechanic villages in Benin city, Nigeria. *IOSR Journal of Environmental Science and Toxicology* (2014)8(1): 9-14
- <sup>34</sup>. Iwegbue CMA, Armioro FO and Onyenibe S. Study of heavy metal speciation in sediments impacted, with crude oil in the Niger Delta, Nigeria. *Italian Journal of Chemical Society* (2006) 97(11-12): 1143 – 1155.
- <sup>35</sup>. Zakir HM, Sultana N and Akter M. Heavy Metal contamination in Roadside soils and Grasses: A case study from Ahaka City, Bangladesh. *Journal of Chemistry, Biology and Physical Science*. (2014)4(2):1661-1673.
- <sup>36</sup>. El – Gamal IM. Distribution pattern of some heavy metals in soil and plants along el-Moukattam highway, Egypt. *International Chemistry and Environmental Health Management*. (2000) pp.518-524.
- <sup>37</sup>. Matthews –Amune OC, Kakulu S. Investigation of heavy metal levels in road-side agricultural soil and plant samples in Adogo, Nigeria, *Academic Journal of Environmental Science*. (2013) 2:31-35
- <sup>38</sup>. Pam AA, Sha'Ato R, Offem OJ. Contributions of Automobile Mechanic Sites to Heavy Metals in Soil: A Case Study of North Bank Mechanic Village Makurdi, Benue State, Central Nigeria. *Journal of Chemistry, Biology and Physical Science*. (2013a) 3(3):2337-2347.
- <sup>39</sup>. Pam AA, Sha'Ato R, Offem OJ. Evaluation of heavy metals in soils around automechanic workshop clusters in Gboko and Makurdi, Central Nigeria. *Journal of Environmental Chemistry and Ecotoxicology*, (2013b) (11):298-306
- <sup>40</sup>. Onder S, Dursun S, Gezgin S Demirbas A. Determination of heavy metal pollution in grass and soil of city centre green areas Konya, Turkey, *Polish Journal of Environmental Studies* (2007) 16 (1):145-154
- <sup>41</sup>. Zhang F, Yan X, Zeng C, Zhang M, Shrestha S, Devkota LP, Yao T. Influence of traffic activity on heavy metal concentrations of roadside farmland soil in mountainous areas. *International Journal of Environmental Research and Public Health*, (2012) 9(5):1715-1731
- <sup>42</sup>. Najib NWAZ, Mohammed SA, Ismail SA, Ahmad WA. Assessment of Heavy Metal in soil due to Human Activities in Kangar, Perlis, Malaysia 2012
- <sup>43</sup>. Lenntech BV. Heavy metals Available <http://www.Lenntech.com.pt/processes/pesadol-heavy-metals.html> 2009
- <sup>44</sup>. USEPA, Integrated Risk Information System (IRIS). National Center for Environmental Assessment, Office of Research and Development, Washington, DC. 1999
- <sup>45</sup>. Megharaj S.J., Singleton I., McClure N.C. and Naidu R. Influence of petroleum hydrocarbon contamination on microalgae and microbial activities in a long term contaminated soil. *Archives on Environmental Contamination and Toxicology*, (2000) 38: 439-445.
- <sup>46</sup>. MacNaughton S.J., Stephen J.R., Venosa A.O., Davis G.A., Chang Y.J. and White D.C.. Microbial population changes during bioremediation of an experimental oil spill. *Journal of Applied Environmental Microbiology*, (1999) 65; 3566-3574.

- <sup>47.</sup> Kaplan C.W. and Kitts C.L. Bacterial succession in a petroleum land treatment unit. *Journal of Applied Environmental Microbiology*, (2004) 70; 1777-1786.
- <sup>48.</sup> Yuni Sri Rahayu, Yulani, Guntur Trimulyono. Isolation and Identification of Hydrocarbon degradation bacteria and phosphate solubilizing bacteria in oil contaminated soil in Bojonegoro East Java, Indonesia. *Indonesian Journal of Science and Technology*. 2019: 134-147
- <sup>49.</sup> Kim, D.J., Lee, D.I., Keller, J. Effects of temperature and free Ammonia On nitrification and nitrite accumulation. In landfill leachate and analysis of its nitrifying Bacterial community by FISH. *Bioresource Technology*, (2006) 97:459-468.
- <sup>50.</sup> Rajbanshi, A. Study on heavy metal resistance bacteria in guesswork sewage treatment plant. *Our Nature*. (2008)6: 52-57.
- <sup>51.</sup> Nies, D.H. Efflux-mediated heavy metal resistance in prokaryotes. *FEMS Microbiology Rev*, (2003) 27: 313-339
- <sup>52.</sup> Valls M, Lorenzo V. Exploiting the genetic and biochemical capacities of bacteria for the remediation of heavy metal pollution. *FEMS Microbiology Reviews* 2002: 26, 327-338
- <sup>53.</sup> Trevors J.T, Oddie K.M, Belliveau B.H. Metal Resistance in Bacteria. *FEMS Microbiology Reviews*. 2005: 32 (39-54)
- <sup>54.</sup> Olukoya D. K., Smith S. I. and M. O. Ilori. Isolation and characterization of heavy metals resistant bacteria from Lagos Lagoon. *Folia Microbiology (Praha)*. (1997)42: 441-444.
- <sup>55.</sup> Nwaugo V. O., Onyeagba, R. A., Akubugwo, E. I. and Ugbogu, O. Soil bacterial flora and enzymatic activities in zinc and lead contaminated soil. *Biokemstri*(2008) 20: 77-84.
- <sup>56.</sup> Mgbemena, C. I., Nnokwe, J. C., Adjero L.A. and Onyemekara N. N. Resistance of bacteria isolated from Otamiri River to heavy metals and some selected antibiotics. *Current Research Journal of Biological Sciences* (2012) 4: 551-556.
- <sup>57.</sup> Oyetibo, G. O., Ilori, M. O., Adebusoye, S. A., Obayori, O. S. and Amund O. O. Bacteria with dual resistance to elevated concentrations of heavy metals and antibiotics in Nigeria in contaminated systems. *Environmental Monitoring Assessment*(2010) 168: 305-314.