

A New Exponential Type Ratio Estimator for the Population Mean in Systematic Sampling

Abstract

In this article we proposed an exponential ratio type estimator for the estimation of finite population mean under systematic sampling. The mean square error of the suggested estimator is computed up to the first degree of approximation and we find suggested estimator is efficient as compared with existing estimators. Furthermore this result is supported by numerical examples as well.

Keywords: Exponential ratio type estimator, Systematic sampling, mean square error (MSE), efficiency.

Introduction:

In the literature of survey sampling, a simple technique of utilizing the known information of the population parameters of the auxiliary variables is through ratio, product, and regression method of estimations using different probability sampling designs such as simple random sampling, stratified random sampling, cluster sampling, systematic sampling, and double sampling. In the present paper we will use knowledge of the auxiliary variables under the framework of systematic sampling. Due to its simplicity, systematic sampling is the most commonly used probability design in survey of finite populations; see W. G. Madow and L. H. Madow [23]. Apart from its simplicity, systematic sampling provides estimators which are more efficient than simple random sampling or stratified random Sampling for certain types of population.

Hajek [9], Cochran [24] and Gautschi [25]. later on the problem of estimating the population mean using information on auxiliary variable has also been discussed by various authors including Quenouille [15], Hansen et al. [12], Swain [1], Banarasi et al. [2], Kadilar et, al. [3], Robson [4], Singh et, al. [5], Singh et, al. [6], Singh at, al. [7], Singh et al. [8], Kushwaha et, al.[10], Khan at, al. [11], Khan et, al [13], Singh [14], Shukla [16], Koyuncu et, al. [17], R. Singh et al. [18], R. Singh et al. [19], Bahl et, al. [20], Srivastava et, al. [21], Tailor et al. [22], and Ozel Kadilar, et, al. [26].

Consider a finite population $U = U_1, U_2, U_3 \dots \dots U_N$ of size N units. A sample of size n is taken at random from the first k units and every k^{th} subsequent unit then, $N = nk$ where n and

k are positive integers thus, there will be k samples each of size n and observe the study variate y and auxiliary variate x for each and every unit selected in the sample.

Let (y_{ij}, x_{ij}) for $i = 1, 2, \dots, k$, $j = 1, 2, \dots, n$ indicate the value of j^{th} unit in the i^{th} sample. Then, the systematic sample means are defined as follows:

$\bar{y}_{st} = t_0 = 1/n \sum_{j=1}^n y_{ij}$, and $\bar{x}_{st} = t_0 = 1/n \sum_{j=1}^n x_{ij}$, unbiased estimators of the population means

$\bar{Y} = 1/n \sum_{j=1}^n y_{ij}$, and $\bar{X} = 1/n \sum_{j=1}^n x_{ij}$, of y on x

To obtain estimators upto first order of approximation, using the following errors terms:

$$e_0 = \bar{y}_{sys} - \bar{Y}/\bar{Y},$$

$$e_1 = \bar{x}_{sys} - \bar{X}/\bar{X},$$

$$e_2 = \bar{z}_{sys} - \bar{Z}/\bar{Z},$$

such that $E(e_i) = 0$ for $i = 0, 1$ and 2

and

$$\rho_{yx} = \frac{s_{yx}}{s_y s_x}$$

$$\rho_{yz} = \frac{s_{yz}}{s_y s_z}$$

$$\rho_{xz} = \frac{s_{xz}}{s_x s_z}$$

$$k = \frac{\rho_{yx} C_y}{C_x}$$

$$k^* = \frac{\rho_{yz} C_y}{C_z}$$

$$\rho_y^* = \{1 + (n-1)\rho_y\}$$

$$\rho_x^* = \{1 + (n-1)\rho_x\}$$

$$\rho_z^* = \{1 + (n-1)\rho_z\}$$

$$\rho^{**} = \frac{\rho_y^*}{\rho_x^*}$$

$$\rho_2^{**} = \frac{\rho_y^*}{\rho_z^*}$$

$$\rho_{1}^{**} = \frac{\rho_y^*}{\rho_z^*}$$

Where

ρ_y, ρ_x, ρ_z are intra class correlation among the pair of units for the variables y,x and z.

2. Estimators in Literature:

In this part, we consider some estimators of the finite population mean in the sampling literature [27-29]. The variance and MSE's of all the estimators computed here are obtained upto first order of approximation.

The variance of the unbiased estimator for population mean is

$$var(t_0) = \lambda \bar{Y}^2 \varphi_0 \quad (1.1)$$

Swain [14] and Shukla [16] suggested the classical ratio and product estimators for finite population mean by are given by

$$t_1 = \bar{y}_{sy} \left(\frac{\bar{X}}{\bar{x}_{sy}} \right) \quad (1.2)$$

$$t_2 = \bar{y}_{sy} \exp \left(\frac{\bar{z}_{sy}}{\bar{Z}} \right) \quad (1.3)$$

The mean square errors upto first orders of approximation are given as follows:

$$MSE(t_1) = \lambda \bar{Y}^2 [\varphi_0 + \varphi_2 (1 - 2k\sqrt{\rho^{**}})] \quad (1.4)$$

$$MSE(t_2) = \lambda \bar{Y}^2 [\varphi_0 + \varphi_3 (1 + 2k\sqrt{\rho_2^{**}})] \quad (1.5)$$

Singh et al. [20] suggested the following ratio and product type exponential estimators as:

$$t_3 = \bar{y}_{sy} \exp \left(\frac{\bar{X} - \bar{x}_{sy}}{\bar{X} + \bar{x}_{sy}} \right) \quad (1.6)$$

$$t_4 = \bar{y}_{sy} \exp \left(\frac{\bar{x}_{sy} - \bar{X}}{\bar{x}_{sy} + \bar{X}} \right) \quad (1.7)$$

The mean square errors upto first orders of approximation are given as follows:

$$MSE(t_3) = \lambda \bar{Y}^2 \left[\varphi_0 + \frac{\varphi_2}{4} (1 - 4k\sqrt{\rho^{**}}) \right] \quad (1.8)$$

$$MSE(t_4) = \lambda \bar{Y}^2 \left[\varphi_0 + \frac{\varphi_2}{4} (1 + 4k\sqrt{\rho^{**}}) \right] \quad (1.9)$$

Tailor et al. [25] define the following ratio-cum product estimator as:

$$t_5 = \bar{y}_{sy} \left(\frac{\bar{X}}{\bar{x}_{sy}} \right) \left(\frac{\bar{z}_{sy}}{\bar{Z}} \right) \quad (2.0)$$

The MSE upto first order of approximation, is given by

$$\text{MSE}(t_5) = \lambda \bar{Y}^2 [\varphi_0 + \varphi_2(1 - 2k\sqrt{\rho^{**}}) + \varphi_3(1 - 2k^{**}\sqrt{\rho_1^{**}}) + 2\varphi_4\sqrt{\rho_y^*\rho_z^*}] \quad (2.1)$$

Where,

$$\varphi_0 = \rho_y^* C_y^2$$

$$\varphi_1 = 2C_x^2 \sqrt{\rho_y^* \rho_x^*}$$

$$\varphi_2 = \rho_x^* C_x^2$$

$$\varphi_3 = \rho_z^* C_z^2$$

$$\varphi_4 = k^* C_z^2$$

Proposed estimator

In this section, motivated Kadilar (2016) we proposed estimator for population mean under systematic sampling as given by :

$$t_{RK} = \bar{y}_{sy} \left(\frac{\bar{x}_{sy}}{\bar{X}} \right)^\alpha \exp \left(\frac{\bar{X} - \bar{x}_{sy}}{\bar{X} + \bar{x}_{sy}} \right) \quad (2.2)$$

To obtain estimators upto first order of approximation, using the following errors terms:

$$E(e_0^2) = \lambda \rho_y^* C_y^2,$$

$$E(e_1^2) = \rho_x^* C_x^2$$

$$E(e_0 e_1) = \lambda C_x^2 \sqrt{\rho_y^* \rho_x^*}$$

Where, $\lambda = \left(\frac{N-1}{nN} \right)$

Expressing (2.2) in terms of e's

$$t_{RK} = \bar{Y}(1 + \varepsilon_0)(1 + \varepsilon_1)^\alpha \exp \left(\frac{\bar{X} - \bar{X}(1 + \varepsilon_1)}{\bar{X} + \bar{X}(1 + \varepsilon_1)} \right)$$

$$t_{RK} = \bar{Y}(1 + \varepsilon_0)(1 + \varepsilon_1)^\alpha \exp \left[\frac{\varepsilon_1}{2} \left(1 + \frac{\varepsilon_1}{2} \right)^{-1} \right] \quad (2.3)$$

$$t_{RK} = \bar{Y}(1 + \varepsilon_0)(1 + \varepsilon_1)^\alpha \exp \left[\frac{\varepsilon_1}{2} \left(1 + \frac{\varepsilon_1}{2} + \frac{\varepsilon_1^2}{4} + \dots \right) \right]$$

$$t_{RK} = \bar{Y}(1 + \varepsilon_0) \left(1 + \alpha \varepsilon_1 + \frac{\alpha(\alpha - 1)}{2} \varepsilon_1^2 + \dots \right) \exp \left[\frac{\varepsilon_1}{2} \left(1 + \frac{\varepsilon_1}{2} + \frac{\varepsilon_1^2}{4} + \dots \right) \right] \quad (2.4)$$

From (2.4)

$$t_{RK} - \bar{Y} \cong \bar{Y} \left[\alpha \varepsilon_1 + \frac{\alpha(\alpha - 1)}{2} \varepsilon_1^2 + \frac{\varepsilon_1}{2} + \frac{\alpha \varepsilon_1^2}{2} + \frac{3\varepsilon_1^2}{8} + \varepsilon_0 + \alpha \varepsilon_1 \varepsilon_0 + \frac{\varepsilon_0 \varepsilon_1}{2} \right] \quad (2.5)$$

Squaring (2.5) on both sides and then taking expectation, the MSE of the estimator \bar{y}_{RK}

$$MSE(t_{RK}) = \lambda \bar{Y}^2 \left[\varphi_0 + \frac{K\varphi_1}{2} + \alpha K\varphi_1 + (\alpha^2 + \alpha) \frac{\varphi_2}{2} + \frac{\varphi_2}{8} \right] \quad (2.6)$$

Obtain the optimum α to minimize $MSE(\bar{y}_{RK})$. Differentiating $MSE(\bar{y}_{RK})$ w.r.t α and equating the derivative to zero. Optimum value of α is given by:

$$\alpha = -\frac{(K\varphi_1 + \varphi_2)}{2\varphi_2}$$

Using the value of α_{opt} in (2.6), we get the minimum value of \bar{y}_{RK}

$$MSE_{min}(t_{RK}) = \lambda \bar{Y}^2 \varphi_0 [1 - \rho_{yx}^2] \quad (2.7)$$

It follows from (2.7) that the proposed estimator \bar{y}_{RK} at its optimum condition is equal efficient as that of the usual linear regression estimator.

Efficiency Comparisons

In this section, the MSE of traditional estimators t_0, t_1, t_2, t_3, t_4 and t_5 are compared with the MSE of proposed estimator \bar{y}_{RK} .

From (1.1) - (2.0) and (2.1)

$$\begin{aligned} [var(t_0) - MSE_{min}(\bar{y}_{RK})] &> 0 \\ [\lambda \bar{Y}^2 \varphi_0 \rho_{yx}^2] &> 0 \end{aligned} \quad (2.8)$$

$$\begin{aligned} [MSE(t_1) - MSE_{min}(\bar{y}_{RK})] &> 0 \\ \lambda \bar{Y}^2 [\varphi_0(1 - 2k\sqrt{\rho^{**}})] - [\lambda \bar{Y}^2 \varphi_0 \rho_{yx}^2] &> 0 \end{aligned} \quad (2.9)$$

$$\begin{aligned} [MSE(t_2) - MSE_{min}(\bar{y}_{RK})] &> 0 \\ \lambda \bar{Y}^2 [\varphi_0(1 + 2k\sqrt{\rho^{**}})] - [\lambda \bar{Y}^2 \varphi_0 \rho_{yx}^2] &> 0 \end{aligned} \quad (3.0)$$

$$\begin{aligned} [MSE(t_3) - MSE_{min}(\bar{y}_{RK})] &> 0 \\ \lambda \bar{Y}^2 \left[\frac{\varphi_0}{4} (1 - 4k\sqrt{\rho^{**}}) \right] - [\lambda \bar{Y}^2 \varphi_0 \rho_{yx}^2] &> 0 \end{aligned} \quad (3.1)$$

$$\begin{aligned} [MSE(t_4) - MSE_{min}(\bar{y}_{RK})] &> 0 \\ \lambda \bar{Y}^2 \left[\frac{\varphi_0}{4} (1 + 4k\sqrt{\rho^{**}}) \right] - [\lambda \bar{Y}^2 \varphi_0 \rho_{yx}^2] &> 0 \end{aligned} \quad (3.2)$$

$$[MSE(t_5) - MSE_{min}(\bar{y}_{RK})] > 0$$

$$\lambda \bar{Y}^2 [\varphi_0(1 - 2k\sqrt{\rho^{**}}) + \varphi_3(1 - 2k^{**}\sqrt{\rho_1^{**}}) + 2\varphi_4\sqrt{\rho_y^* \rho_z^*}] - [\lambda \bar{Y}^2 \varphi_0 \rho_{yx}^2] > 0 \quad (3.3)$$

Empirical Study

To examine the merits of the proposed estimator over the other existing estimators at optimum conditions, we have considered natural population data sets from the literature. The sources of population are given as follows.

(Source: Tailor et al. [20]). Consider Population

$$N = 15, n = 3, \bar{X} = 44.47, \bar{Y} = 80, \bar{Z} = 48.40, C_y = 0.56, C_x = 0.28, C_z = 0.43$$

$$S_y^2 = 2000, S_x^2 = 149.55, S_z^2 = 427.83, S_{yx} = 538.57, S_{yz} = -902.86, S_{xz} = -241.06,$$

$$\rho_{yx} = 0.9848, \rho_{yz} = -0.9760, \rho_{xz} = -0.9530, \rho_y = 0.6652, \rho_x = 0.707, \rho_z = 0.5487.$$

To find (PREs) of the estimator, we use the following formula

$$PRE(t_\alpha, t_0) = MSE(t_0)/MSE(t_\alpha) \times 100, \text{ For } \alpha = 0, 1, 2, 3, 4, 5 \text{ and } RK$$

Table 1 The percent relative efficiency of different estimators with respect to t_0 .

Population		
Estimator	$MSE(t_\alpha)$	$PRE(t_\alpha, t_0)$
t_0	1455.08	100.00
t_1	373.32	389.62
t_2	768.06	189.45
t_3	820.09	177.43
t_4	1044.42	139.32
t_5	187.08	777.79
t_{RK}	43.88	3316.04

Conclusion

A new exponential type estimator is proposed under systematic sampling and the properties of the suggested estimator are obtained up to first order of approximation. It has been seen that the suggested estimator performed better than the existing estimator both theoretically and empirically.

References

- [1] A. K. P. C. Swain, "The use of systematic sampling ratio estimate," *Journal of the Indian Statistical Association*, vol. 2, pp. 160–164, 1964.

- [2] Banarasi, S. N. S. Kushwaha, and K. S. Kushwaha, "A class of ratio, product and difference (R.P.D.) estimators in systematic sampling," *Microelectronics Reliability*, vol. 33, no. 4, pp. 455–457, 1993.
- [3] C. Kadilar, and H. Cingi, "An improvement in estimating the population mean by using the correlation coefficient," *Hacettepe Journal of Mathematics and Statistics*, vol. 35, no. 1, pp. 103–109, 2006.
- [4] D. S. Robson, "Application of multivariate polykeys to the theory of unbiased ratio-type estimation," *The Journal of the American Statistical Association*, vol. 59, pp. 1225–1226, 1957.
- [5] H. P. Singh and R. Singh, "Almost unbiased ratio and product type estimators in systematic sampling," *Questiio*, vol. 22, no. 3, pp. 403–416, 1998.
- [6] H. P. Singh and R. S. Solanki, "An efficient class of estimators for the population mean using auxiliary information in systematic sampling," *Journal of Statistical Theory and Practice*, vol. 6, no. 2, pp. 274–285, 2012.
- [7] H. P. Singh and N. K. Jatwa, "A class of exponential type estimators in systematic sampling," *Economic Quality Control*, vol. 27, no. 2, pp. 195–208, 2012.
- [8] H. P. Singh, R. Tailor, and N. K. Jatwa, "Modified ratio and product estimators for population mean in systematic sampling," *Journal of Modern Applied Statistical Methods*, vol. 10, no. 2, pp. 424–435, 2011.
- [9] J. Hajek, "Optimum strategy and other problems in probability sampling," *Casopis pro Pestovani Matematiky*, vol. 84, pp. 387–423, 1959.
- [10] K. S. Kushwaha and H. P. Singh, "Class of almost unbiased ratio and product estimators in systematic sampling," *Journal of the Indian Society of Agricultural Statistics*, vol. 41, no. 2, pp. 193–205, vi, 1989.
- [11] M. Khan and H. Abdullah, "A note on a differencetype estimator for population mean under twophase sampling design," *SpringerPlus*, vol. 5, article 723, 7 pages, 2016.
- [12] M. H. Hansen, W. N. Hurwitz, and M. Gurney, "Problems and methods of the sample survey of business," *Journal of the American Statistical Association*, vol. 41, no. 234, pp. 173–189, 1946.
- [13] M. Khan and R. Singh, "Estimation of population mean in chain ratio-type estimator under systematic sampling," *Journal of Probability and Statistics*, vol. 2015, Article ID248374, 2 pages, 2015.
- [14] M. P. Singh, "Ratio-cum-product method of estimation," *Metrika*, vol. 12, pp. 34–42, 1967.

- [15] M. H. Quenouille, "Notes on bias in estimation," *Biometrika*, vol. 43, pp. 353–360, 1956.
- [16] N.D. Shukla, "Systematic sampling and product method of estimation," in *Proceeding of the all India Seminar on Demography and Statistics*, Varanasi, India, 1971.
- [17] N. Koyuncu and C. Kadilar, "Family of estimators of population mean using two auxiliary variables in stratified random sampling," *Communications in Statistics: Theory and Methods*, vol. 38, no. 13–15, pp. 2398–2417, 2009.
- [18] R. Singh, S. Malik, M. K. Chaudhary, H. Verma, and A. A. Adewara, "A general family of ratio-type estimators in systematic sampling," *Journal of Reliability and Statistical Studies*, vol. 5, no.1, pp. 73–82, 2012.
- [19] R. Singh, S. Malik, and V. K. Singh, "An improved estimator in systematic sampling," *Journal of Scientific Research*, vol. 56, pp. 177–182, 2012.
- [20] S. Bahl and R. K. Tuteja, "Ratio and product type exponential estimators," *Journal of Information & Optimization Sciences*, vol. 12, no. 1, pp. 159–164, 1991.
- [21] S. K. Srivastava and H. S. Jhaji, "A class of estimators of the population mean using multi-auxiliary information," *Calcutta Statistical Association Bulletin*, vol. 32, no. 125-126, pp. 47–56, 1983.
- [22] T. Tailor, N. Jatwa, and H. P. Singh, "A ratio-cum-product estimator of finite population mean in systematic sampling," *Statistics in Transition*, vol. 14, no. 3, pp. 391–398, 2013.
- [23] W. G. Madow and L. H. Madow, "On the theory of systematic sampling. I," *Annals of Mathematical Statistics*, vol. 15, pp. 1–24, 1944.
- [24] W. G. Cochran, "Relative accuracy of systematic and stratified random samples for a certain class of populations," *The Annals of Mathematical Statistics*, vol. 17, pp. 164–177, 1946.
- [25] W. Gautschi, "Some remarks on systematic sampling," *Annals of Mathematical Statistics*, vol. 28, pp. 385–394, 1957.
- [26] Ozel Kadilar, G. A new exponential type estimator for the population mean in simple random sampling. *Journal of Modern Applied Statistical Methods*, 15(2), 15, 207-214 (2016).
27. Zaman, T. (2021). An efficient exponential estimator of the mean under stratified random sampling. *Mathematical Population Studies*, 28(2), 104-121.

28. Zaman, T., & Kadilar, C. (2021). Exponential ratio and product type estimators of the mean in stratified two-phase sampling. *AIMS Mathematics*, 6(5), 4265-4279.

29. Zaman, T., & Kadilar, C. (2021). New class of exponential estimators for finite population mean in two-phase sampling. *Communications in Statistics-Theory and Methods*, 50(4), 874-889.

UNDER PEER REVIEW