

Original Research Article

Susceptibility of the dengue vector *Aedes aegypti* to various insecticides in Lahore, Pakistan

ABSTRACT

Objective

Present studies were carried out to examine the susceptibility of larvae and adults of *Aedes aegypti* (Ae. aegypti) from laboratory and field populations of Lahore city against four major groups of commonly used insecticides; **neonicotenoids** **neonicotinoids** (Imidacloprid 5% SC), phenyl-pyrazoles* (fipronil 2.5% EC), pyrethroids (Deltamethrin 2.5% SC) and organophosphates (DDVP 50% EC) world health organization standard procedures were observed.

Methods

For laboratory strain, adults and larvae were collected from the Insectary of National Institute of Malaria Research and Training (NIMRT), Lahore, Pakistan. Insecticides of four major groups. Larval bioassays were conducted according to the World Health Organization procedure with minor modifications. Larval mortality was finalized after 48 hours for Deltamethrine, Imidacloprid, DDVP and 72 hours for Fipronil.

Results

Larval and adult toxicity results were different for wild and laboratory strains exposed to different groups of insecticides. Regression analysis showed non-significant trends in **mortality**. Imidacloprid proved to be the most toxic to wild larval strains of

Comment [T1]: You need to clarify this sentence in the result or discussion part. Where is your explanation?

Ae. aegypti collected from different localities of Lahore while Fipronil was also active for wild larval samples.

Conclusion

Deltamethrin showed the least activity against both adults and larval strains. The susceptibility of the field strains was lower than laboratory strains; the resistant resistance ratio varies from insecticide to insecticide.

Key words Keywords *Aedes aegypti*, neonicotenoids neonicotinoids, phenylpyrazoles, pyrethroids, organophosphates

Background

Dengue fever is a mosquito-borne mosquito-borne disease widespread throughout the world, there is no safe affordable and effective vaccine available. For the purpose to control the disease vector control is an effective method (Kuno 1997, Mohanty and Prakash 2008). However, Dengue Fever is less widespread in the temperate than in tropic countries. The life cycle of pathogens is delayed by the cold weather in temperate climate temperate climate, and due to the shorter life span mosquitoes usually die before pathogens become infective (Busvine 1980) Climate is mostly favorable for the breeding of mosquitoes in Pakistan (Schafer and Lundstrom 2001) *Ae. aegypti* (Linnaeus) and *Ae. albopictus* (Skuse) are the important vectors of dengue fever in tropical and subtropical regions of the world (Lee and Yap 1999). Major diseases like dengue fever, dengue hemorrhagic fever (DHF) are spread by *Ae. aegypti*. Neurotoxic insecticides are either neuro inhibitory or neuro excitatory. The behavioral effects of neuro excitation on organism organisms are tremors, hyperactivity and rigid paralysis, in the case of neuro inhibition, lower immobility and flaccid paralysis occur. The mortality is caused by energy depletion and neuromuscular fatigue (Michael 2003). In Pakistan, a major strategy to prevent and control vector-borne vector-borne diseases like

Comment [T2]: You can merge your result part and conclusion together.

dengue fever is ~~the~~ use of insecticides. During ~~the~~ dengue outbreak in 2011, insecticides used were reportedly not effective. Although some previous studies indicated that there is a high level of resistance to some commonly used insecticides, ~~but~~ the level of resistance was not studied for ~~the~~ *Aedes* population ~~at-in~~ Lahore. According to World Health Organization (WHO) standards and control program, there is a continuous need to monitor insecticide resistance. The present study was conducted to determine the activity of different groups of insecticides against laboratory and field strains of larvae and adults of *Ae. aegypti*.

Material and Methods

Mosquito collection sites

For wild strain, larvae were collected from those sites in Lahore where insecticides were used at high frequency and quantity. For laboratory strain, adults and larvae were collected from the Insectary of National Institute of Malaria Research and Training (NIMRT), Lahore, Pakistan.

Mosquitoes

For wild strain of *Ae. aegypti*, first star larvae were collected from the field using plastic trays and were kept under laboratory conditions of controlled temperature ($25^{\circ}\text{C} \pm 2$) and relative humidity respectively ($70\% \pm 5$) with a constant photoperiod (light: dark, 14h:10h). These larvae were fed on a mixture of biscuits, dried yeast and ~~fat free~~ fat-free milk (1:1:1). After 3-5 days these wild larvae hatched into adults, and were used for adult bioassays. for, larvae bioassays laboratory strains were used. Adults ~~taken~~ took from insectary from larvae collected in 2009 from indoor and outdoor populations of mosquitoes in Lahore, Pakistan.

Insecticides

Insecticides of four major groups: ~~neonicotinoids~~ ~~neonicotinoids~~ (imidacloprid 5% SC), phenylpyrazoles (fipronil 2.5% EC), pyrethroids (deltamethrin 2.5% SC) and organophosphates dichlorvos (DDVP 50% EC) were used during this study. Stock solutions of four insecticides, DDVP, deltamethrin, fipronil and imidacloprid were prepared in distilled water.

Larval bioassay

Larval bioassays were conducted according to the World Health Organization procedure (WHO, 1981) with minor modifications. For the larval bioassay, fourth instars larvae (batches of 20) were introduced in 100 ml of water in ~~a~~ plastic cups containing different concentrations of ~~the above mentioned~~ ~~above-mentioned~~ insecticides. Into each cup, 0.02g of ~~the powdered~~ ~~powdered~~ mixture was added to feed ~~the~~ larvae. Five different concentrations for each insecticide ranging from 0.001 to 0.5µl were used to determine lethal concentrations. Bioassays were done in three replicates for each concentration ~~in order to~~ ~~to~~ get valid results. In ~~the~~ control test, distilled water was used. Plastic cups were inspected after 24h and the numbers of dead larvae were recorded. Larval mortality was finalized after 48 hours for deltamethrin, imidacloprid, DDVP and 72 hours for fipronil. In bioassay moribund larvae ~~was~~ ~~were~~ considered dead.

Adult bioassay

Adult bioassays were conducted according to the World Health Organization procedure (WHO 1981) with some modifications. ~~Three~~ ~~From three~~ to five days old adults (batches of 25) were exposed to Whatman # 1 filter paper treated with 0.05% Imidacloprid, 0.01% Fipronil, 0.5% Deltamethrin and 0.5% Dichlorvos (DDVP) in a glass test tube under the same laboratory conditions as for larvae. In control, no treatment was given to filter paper. After 24 h mortality was observed.

Data analysis

Data from bioassays were expressed in mean \pm S.E.M by using Minitab statistical software (Version 13.20). LC₅₀ with their 95% confidence intervals was estimated by using the EPA Probit analysis program (version 1.5). Results were statistically significant when $P \leq 0.05$. Duncan's multiple range tests ~~was~~ were applied to compare the concentrations of insecticides with a significant difference at the 5 % level using New Co stat. Log concentration-~~probability~~ probability regression lines were drawn for the insecticides used in larval and adult bioassays. The resistance status was determined according to WHO criteria.

Results

From table 1, it is evident that larval mortality for fipronil ranges from 18-99% for laboratory strain of *Ae.aegypti* at 0.005- 0.003 μ l and ~~for~~ field strain ranges from 15-97% at 0.008-0.048 μ l showing its high activity as compared to other insecticides. From all tested insecticides against *Ae .aegypti* larvae, deltamethrin proved to be ~~resistance-resistant~~ as its percentage mortalities ranges from 15-85% at 0.2-0.65 μ l in the laboratory strain and for field strain it decreases to 10-80% at 0.3-0.7 μ l. Bioassays results indicate that the percentage mortality of *Ae.aegypti* exposed to larvicides ~~were~~ was lower in the field populations than in the laboratory strain.

As far as LC₅₀ is concerned, deltamethrin shows the highest LC₅₀ values 0.4 and 0.52 μ l for laboratory and field strain depicting its low effectiveness in the control of *Ae.aegypti*. The LC₅₀ values were recorded in the lab and field strain of *Ae. aegypti* after treatment with fipronil were 0.012 and 0.022 μ l respectively. While imidacloprid and DDVP showed intermediate activities (table 1).

The diagnostic doses of imidacloprid, fipronil, deltamethrin and DDVP are 0.05 μ l, 0.03 μ l, 0.7 μ l and 0.5 μ l respectively (table1). Hundred adult females derived from wild larvae after one generation were exposed to each group of insecticides. The lowest percentage mortality rate (75%) was observed in the deltamethrin treated group, representing the development of resistance in the field population. The highest percentage mortality rate (90%) was observed in the fipronil treated group. According

to WHO criteria of adult susceptibility test *Ae.aegypti* population showed high tolerance to imidacloprid, fipronil and DDVP but high resistance to deltamethrin.

Discussion

The present study was performed to determine the susceptibility of *Ae.aegypti* to different groups of insecticides. In this study, it was observed that laboratory strain was more susceptible than field strain. From the laboratory results of mortality after exposure to different groups of insecticides, fipronil was the most active followed by imidacloprid. The LC₅₀ values of imidacloprid and fipronil were markedly low. New classes of insecticides, such as imidacloprid and fipronil were registered in the last decade and are not commonly used for mosquitoes control in Pakistan, therefore *Ae.aegypti* were more susceptible to these insecticides.

From our results, the outstanding performance of fipronil was in agreement with Pridgeon et al. (2008) who reported relative potency of 19 insecticides against female *Ae.aegypti* (L). The insecticides used for bioassay against *Ae. aegypti* larvae and adult adults have a different mode-modes of actions-action for killing. In order to investigate the efficacy of each insecticide, six concentrations were applied to estimate 0-100% mortality. After treatment, these female-female mosquitoes were transferred to plastic cups provided with 10% sucrose solution. Mortality was recorded after 24 hours. Among 19 insecticides used fipronil was considered highly effective against all tested *Ae. aegypti*. Similarly, Sandra (2010) also conducted the a study in which toxic baits were prepared by using active ingredients of five insecticidal groups (macrocyclic lactones, neonicotinoids, phenylpyroles, pyrethroids and pyrroles) in sucrose solution. These toxic baits were used against *Anopheles quadrimaculatus*, *Ae.taeniorhynchus* and *Culex quinquefasciatus*. In *Ae. taeniorhynchus*, imidacloprid was considered to be the most toxic insecticide in the present study.

Previous studies and the present study reveals that insecticides used previously for control of dengue vectors caused the development of resistance. *Ae.aegypti* was highly

resistant to the pyrethroid group of insecticides. ~~Cross-resistance~~Cross-resistance can be developed in mosquitoes between pyrethroid and non-pyrethroid insecticides. This phenomenon was well explained in the study of Sathantriphop et al. (2006). In that study, the potency of insecticides (pyrethroids, organochlorines, carbamates and organophosphates) ~~were-was~~ recorded for 1 hour against *Cx. quinquefasciatus* and *Ae. aegypti* females. When *Ae. aegypti* females were tested against insecticides, it was found that resistant to permethrin, and highly tolerant to deltamethrin but high susceptibility was observed ~~to~~to organophosphate (fenitrothion).

Our study emphasized the susceptibility of *Ae. aegypti* to commonly used (DDVP and Deltamethrin) insecticides and newly introduced fipronil and imidacloprid in Lahore city. Our results reveal an increase in the level of tolerance to operational usage of larvicides and adulticides after dengue outbreaks in Lahore. The main problem was the development of resistance in vectors which can be managed by monitoring susceptibility status in vector control programs. Resistance developed in mosquitoes to these insecticides especially deltamethrin (pyrethroids) ~~was-resulted~~resulted from household use. Insecticidal products such as liquid, mat, coil, and cream formulations have ingredients of pyrethroids. These products play an important role in the development of resistance in *Ae. aegypti*. Frequent use of insecticides with the same modes of action can accelerate the development of resistance.

The use of low concentrations of pyrethroids and organophosphates for ~~mosquitoes~~ mosquito control was effective and safe (Chareonviriyaphap et al. 1999, Somboon et al. 2003). Thus insecticides play an imperative role in vector control. ~~Application-The~~ application of larvicides and adulticides are principal methods for the control of ~~vector~~ vector-borne diseases (Reiter and Gubler 1997). Resistance in ~~the mosquitoes'~~ mosquitoes population was due to incomplete and infrequent coverage in examining and reporting. The extensive application of pyrethroids for mosquito and agriculture pest control caused an indirect contribution ~~for-to the~~ development of resistance to these classes of insecticides. Therefore the results of the present study suggest the need for

Comment [T3]: How this phenomenon was well explained?. I still do not see it in the next sentences. Pyrethroid resistance can be explained by two major mechanisms: increased detoxification and changes in target site insensitivity.

mosquitoes control ~~professional~~ professionals to search for strategies to prevent or delay the development of ~~insecticides~~ insecticide resistance in Lahore city, Pakistan.

References

1. **Ashrafi, S. H., M. Anwarullah, R. A. Khan, and S. A. Ali. 1966.** Mass rearing of mosquitoes, houseflies and cockroaches in PCSIR Laboratories for insecticide testing. *Sci. Ind.* 4: 312-316.
2. **Busvine, J. R. 1980.** Recommended methods for detection and measurement of resistance of agricultural pests to pesticides, p. 44-46. FAO, Rome.
3. **Chareonviriyaphap, T., B. Aum-aung, and S. Ratanatham. 1999.** Current insecticide resistance patterns in mosquito vectors in Thailand. *Southeast Asian. J. Trop. Med. Public. Health.*30(1): 184-194.
4. Kamgang, B., Marcombe, B., Chandre, F., Nchoutpouen, E., Nwane, P., Etang, J., Corbel, V. and Paupy, V. 2011. Insecticide susceptibility of *Aedes aegypti* and *Aedes albopictus* in Central Africa. *Parasit. Vectors.* 4: 79-87.
5. Kuno, G. 1997. Factors influencing the transmission of dengue viruses. In: Dengue and dengue hemorrhagic fever. CAB International, London, United Kingdom. Pp: 61-68.
6. Lee, C. Y. and Yap, H. H. 1999. Overview on urban pests: A Malaysian perspective. In: Urban Pests Control: A Malaysian Perspective Vector Control Research Unit. University Sains Malaysia. Pp. 1-7.
7. Michael, E. S. 2003. Neurological effects of insecticides. In: Encyclopedia of Pest Management(eds. by Marcel Dekker, I.) Purdue University, West Lafayette, Indiana, U.S. Pp: 50-80.
8. Mohanty, S. S. and Prakash, S. 2008. Laboratory and field evaluation of the fungus *Chrysosporium lobatum* against larvae of the mosquito *Culex quinquefasciatus*. *Parasitol. Res.*102: 881-886.

Comment [DKct4]: Authors need to write references in accordance with the journal's new reference style, effective from 09 Oct, 2012

9. Paul, A., Harrington, L. C., and Scott, J. G. 2006. Evaluation of novel insecticides for control of dengue vector *Aedes aegypti* (Diptera: Culicidae). *J. Med. Entomol.* 43: 55-60.
10. Reiter, P. and Gubler, D. J. 1997. Surveillance and control of urban dengue vectors. In: *Dengue and Dengue Haemorrhagic Fever*. CAB International, [New York New York](#). Pp. 425-462.
11. Rozendaal, A. J. 1999. Mosquitoes and other biting ~~diptera~~ [Diptera](#). In: *Vector control methods use by individuals and communities*. A.I.T.B. S. publishers and distributors, Dehli. Pp: 12-17.
12. Sandra, A. A. 2011. Susceptibility of adult mosquitoes to insecticides in aqueous sucrose baits. *J. Vector Ecol.* 36(1): 59-67.
13. Sathantriphop, S., Paeporn, P. and Supaphathom, K. 2006. Detection of insecticides resistance status in *Culex quinquefasciatus* and *Aedes aegypti* to four major groups of insecticides. *Trop. Biomed.* 23(1): 97-101.
14. Schafer, M. and Lundstrom, J. O. 2001. Comparison of mosquito (Diptera: Culicidae) fauna characteristics of forested wetlands in Sweden. *Ann. Entomol. Soc. Am.* 94(4): 576-582.
15. Somboon, P., Prapanthadara, L. A. and Suwonkerd, W. 2003. Insecticide susceptibility tests of *An. minimus*, *Aedes aegypti*, *Aedes albopictus*, and *Culex quinquefasciatus* in northern Thailand. *Southeast Asian. J. Trop. Med. Public Health.* : 87-93.

Table 1: Susceptibility of *Ae .aegypti* larvae (lab and field strains) against various insecticides

Formatted: Centered

Insecticides	Mosquito strains	Effective concentrations mg/l	Larval mortality (%) ^a	Statistical parameters		Resistance Ratio (RR ₅₀) ^b
				LC ₅₀ /μl	Slope	
Imidacloprid 5% SC	Lab strain	0.03-0.13	16-95	0.060	4.00	2.1
	Field strain	0.05-0.20	18-90	0.127	3.72	
Fipronil 2.5% EC	Lab strain	0.005-0.03	18-99	0.012	2.82	1.8
	Field strain	0.008-0.04	15-97	0.022	3.00	
Deltamethrin 2.5% SC	Lab strain	0.2-0.65	15-85	0.40	3.66	1.3
	Field strain	0.3-0.70	10-80	0.520	6.06	
DDVP 50% EC	Lab strain	0.05-0.19	17-92	0.104	3.91	1.8
	Field strain	0.07-0.35	19-88	0.187	2.90	

a: Three replicates , 20 larvae each; control mortalities ranged from 0.0% -3.0%

b: RR₅₀: = LC₅₀ of field strain / LC₅₀ of the lab strain

Table 2: Susceptibility of *Ae .aegypti* adults against various insecticides

Insecticides	Diagnostic dosages/ mg/μl	No. of mosquitoes exposed	Mortality	Susceptibility status
Imidacloprid 5% SC	0.05	100	85	Tolerant
Fipronil 2.5% EC	0.03	100	90	Tolerant
Deltamethrin 2.5% SC	0.7	100	75	Resistant
DDVP 50% EC	0.5	100	80	Tolerant

While susceptibility of *Ae. aegypti* adults against diagnostic dosages of various insecticides, ~~several number of~~ mosquitoes exposed and mortality are shown in Figure 1.




Figure 1: Susceptibility of *Ae. aegypti* adults against diagnostic dosages of various insecticides, number of mosquitoes exposed and mortality

UNDER PEER REVIEW

Formatted: Numbered + Level: 1 +
Numbering Style: 1, 2, 3, ... + Start at: 1 +
Alignment: Left + Aligned at: 0" + Indent at:
0,3"

UNDER PEER REVIEW