

Original Research Article

The Prevalence and Distribution of Enteric Bacteria in Meat and Meat Products

Abstract

Food-borne diseases are the most serious international health issue, causing economic losses and lives. The enteric bacteria is the most difficult bacterial contamination of raw and processed beef products worldwide. It is also the most prevalent type of food poisoning. **Aim:** The current study sought to determine the presence and distribution of enteric bacteria associated with various types of meat (red and white) and their fresh and frozen products. **Methods:** 36 meat samples were collected from local markets in Nineveh Governorate. The contamination of various meat samples with enteric bacteria was assessed. The bacteria were subsequently isolated and identified using culture, microscopy, and biochemical techniques. Vitek-2 device was used to verify the diagnosis. **Results:** All of the meats tested were contaminated with enteric bacteria to varied degrees. Fresh chicken meat had the highest rate of enteric bacterial infection, with a logarithm of 1.12×10^8 CFU/g. The frozen beef samples had the least contamination, with a logarithm of 7.4×10^4 CFU/g. The results revealed that 57 isolates from the intestine family bacteria included 13 species: *C. freundii*, *C. koseri*, *E. coli*, *Enterobacter* spp., *K. oxytoca*, *K. pneumoniae*, *P. mirabilis*, *P. vulgaris*, *P. stuartii*, *S. paratyphi A*, *S. typhi*, *Shigella* spp., and *Y. enterocolitica*. *E. coli* was found in the most meat varieties analyzed, accounting for 19% of the total. **Conclusions:** Because it is evident that meat can be contaminated with a wide variety of hazardous bacteria, basic hygiene procedures help to decrease the amount of contaminated microbes.

Keywords: Enteric bacteria, Meat contamination, *E. coli*, Culture media

Introduction

Food-borne illnesses are the most serious international health issue, resulting in economic losses and lives. According to the World Health Organization (WHO), 600 million individuals worldwide are affected by foodborne infections each year. Every year, around 420,000 people die as a result of antibiotic-resistant microorganisms [1]. Meat is one of the most perishable foods and a high-protein source. Meat, on the other hand, poses a high risk of food poisoning since it contains all of the ingredients that promote bacterial growth. As a result, meat, like other perishable goods, is preserved through canning and freezing [2, 3].

The category of enteric bacteria is the most common contamination of meat and its numerous derivatives globally. Not only that, but this bacteria is linked to all occurrences of food poisoning, including *E. coli*, *Proteus* spp, *Salmonella* spp, and *Shigella* spp [4]. Its pathogenicity is primarily determined by its ability to release and secrete numerous toxins, as well as its ability to

Comment [PM1]: Must be improve

adhere to surfaces, form biofilms, and produce various virulence enzymes, particularly hemolytic and proteolytic enzymes [5]. The manner of preparing meat, the tools used in dealing with it, the personal cleanliness of the employees, and the method of storing and exhibiting it make it susceptible to contamination [6]. Bacterial food poisoning instances are classified into two kinds. The first is a case of food poisoning induced by the presence of bacteria. The other is poisoning from bacteria toxins generated or released in food. Poisoning-related disease symptoms include abdominal discomfort, diarrhea, vomiting, nausea, fever, breathing difficulties, and death in severe situations [7]. The severity and impact of this form of disease are determined by the interplay of the pathogen transmitted by food, the host, the food, and the environment [8].

Given the health and economic importance of bacterial contamination of meat sources, as well as the pathogenicity importance of members of the enterobacteriaceae, the current study was designed to assess the prevalence and distribution of different types of contaminated enteric bacteria in various meat and its products.

Materials and Methods

36 meat samples were acquired from various sources (fresh beef, fresh lamb, fresh chicken, fresh fish, frozen red meat, frozen pastrami, frozen chicken liver, frozen chicken breast, frozen burger). Meat samples were collected in sterile boxes from local markets in Nineveh Governorate and sent immediately to the laboratory in less than two hours to undertake the initial isolation stages for enteric bacteria.

Isolation and estimation of contamination of meat with enteric bacteria:

Formatted: Font: Not Bold, Italic

10g of meat samples are placed in 90ml of peptone water media (LAB/England). A mixer (Stainless Steel/Chania) crushed it. Using normal saline, 1ml of solution was diluted into multiple decimal dilutions. 1ml of the final dilutions was inoculated on MacConkey agar (LAB/England). The number of colonies was determined by CFU/g. The isolated bacteria were classified based on their capacity to ferment lactose. The microorganisms were cleansed and prepared for further testing [9].

Phenotypic diagnosis:

Formatted: Font: Not Bold, Italic

The diagnosis was made based on the phenotypic characteristics of bacterial colonies in terms of form, color, texture, and fermentation type on MacConkey media. Gram stain was used for microscopic inspection [10].

Biochemical tests:

Formatted: Font: Not Bold, Italic

All isolates were subjected to the following biochemical [11] tests: Oxidase, indol, methyl red (MR)/Voges-Proskauer (VP), citrate utilization, triglyceride iron test (TSI), and Urease test. The Vitek-2 system was used for confirmatory diagnostic testing.

Results and Discussion

The results revealed a clear variance in the extent of contamination with different meat sources, as indicated in Table (1) and Figure (1, 2). The manner of preparing meats, the tools used in dealing with them, the personal cleanliness of the employees, and the method of keeping and presenting them may be the source of contamination of different meats with different enteric bacteria. As a result, it becomes an ideal environment for bacterial reproduction [12].

According to the study, fresh chicken meat had the highest percentage of enteric bacterial infection, with a logarithm of 1.12×10^8 CFU/g. The high pollution rate might be attributed to the frequent use of the chicken feather removal device and the failure to wash it after extended periods of time, as well as the failure to change the water for long durations of time. Fresh fish had the second highest contamination rate of 1.16×10^7 CFU/g, which could be attributed to the placing of fish in dirty water ponds, the frequent use of these ponds, and the lack of water exchange for extended periods of time. These ponds are thought to be an ideal environment for the growth of microorganisms.

Table 1: Average number of bacterial colonies (CFU/g) in different meat samples

Sample source	Average no. of colonies CFU/g
Fresh beef meat	7.2×10^5
Fresh sheep meat	6.8×10^5
Fresh chicken meat	1.12×10^8
Fresh fish	1.16×10^7
Frozen beef	7.4×10^4
Frozen chicken liver	4.7×10^5
Frozen chicken meat	1.31×10^5
Frozen Iraqi beef burger	5.8×10^4
Frozen beef pastrami	5.1×10^4

Formatted: Font: 12 pt

Formatted: Space Before: 6 pt, After: 6 pt

Formatted: Space Before: 6 pt, After: 6 pt

Formatted: Space Before: 6 pt, After: 6 pt

Formatted: Space Before: 6 pt, After: 6 pt

Formatted: Space Before: 6 pt, After: 6 pt

Formatted: Space Before: 6 pt, After: 6 pt

Formatted: Space Before: 6 pt, After: 6 pt

Formatted: Space Before: 6 pt, After: 6 pt

Formatted: Space Before: 6 pt, After: 6 pt

Formatted: Space Before: 6 pt, After: 6 pt




Figure 1: Contamination percentages in different meat sources log₁₀(CFU)/g of enterobacteria.

Comment [PM2]: error bar?

Formatted: Font: 12 pt

Formatted: Font: 12 pt




Figure 2: Models representing the extent of contamination with enteric bacteria on McConkeyagar media in different meat samples. A: Showing a sample of fresh beef from (the third dilution). B:Result of culturing a frozen chicken liver sample from (the third dilution).

Formatted: Font: 12 pt

Prevalence and distribution of enteric bacteria:-

Formatted: Font: Not Bold, Italic

Enterobacteriaceae were isolated by using MacConkey agar media divided based on their ability to ferment lactose (pink colonies) and non-lactose-fermenting bacteria (transparent colonies) [10]. Enterobacteriaceae isolates of meat sources identified based on the agronomic, microscopic, and biochemical features mentioned in Table (2), as well as confirmation of the isolates' diagnosis by using Vitek-2. Isolated bacteria were dispersed among 13 distinct species, which included: *C. freundii*, *C. koseri*, *E.coli*, *Enterobacter* spp., *K. oxytoca*, *K. pneumoniae*, *P. mirabilis*, *P. vulgaris*, *P. stuartii*, *S.paratyphi A*, *S. typhi*, spp. *Shigella* and *Y. enterocolitica*. All

bacterial types were isolated from fresh chicken, fresh beef, fresh lamb, fresh fish, frozen red meat, frozen pastrami meat, frozen chicken breast, frozen chicken liver, and frozen burger.

Table 2: Biochemical tests of bacteria isolated from different meat sources.

Bacteria Sp	Oxidase	Urease	TSI Test			IMViC Test			
			H ₂ S	Gas	S/B	CIT	VP	MR	IND
<i>C. koseri</i>	-	-	-	+	A/A	+	-	+	+
<i>E.coli</i>	-	-	-	+	A/A	-	-	+	+
<i>P. vulgaris</i>	-	+	+	-	K/A	-	-	+	+
<i>P. mirabilis</i>	-	+	+	+	K/A	+	-	+	-
<i>Enterobacter</i>	-	+	-	+	A/A	+	+	-	-
<i>P. stuartii</i>	-	-	-	+	K/A	+	-	+	+
<i>Shigella</i>	-	-	-	-	K/A	-	-	+	-
<i>C. freundii</i>	-	-	+	+	A/A	+	-	+	-
<i>S.paratyphi A</i>	-	-	-	+	K/A	-	-	+	-
<i>S. typhi</i>	-	-	+	-	K/A	-	-	+	-
<i>Y. enterocolitica</i>	-	+	-	-	K/A	-	-	+	-
<i>K. pneumoniae</i>	-	+	-	+	A/A	+	+	-	-
<i>K. oxytoca</i>	-	+	-	+	A/A	+	+	-	+

TSI: Triple Sugar Iron, S/B: butt/slant, Gas: gas production, A: acid, K: alkaline, IND: Indole, MR: Methyl red, VP: Voges-Proskauer, CIT: Citrate.

Because most isolated bacteria are harmful, they cause diseases such as intestinal infections, urinary tract infections, pneumonia, meningitis, and sepsis. The majority of them also produce intestinal toxins, which induce food poisoning [13]. Clinically, the majority of gut bacteria genera are categorized as opportunistic infections. Some of them are classed as basic pathogens, such as *Salmonella*, *Shigella*, and *Yersinia*, as well as being one of the most common food-borne pathogens, particularly *E.coli* bacteria [4]. These are the species that naturally inhabit the human intestine and spread easily by hand contamination, water, and food. Treatment and control are difficult, especially in low-income nations [14].

The results revealed a difference in the presence of the isolated enteric bacterial species, which implies variance in the size and kind of contamination. Some meats had only two species (frozen chicken liver), whereas others contained seven types (fresh beef). The remaining samples contained 6-3 enteric bacteria, according to the Table (3). *E. coli* was the most common from different isolate sources (6 sources), reflecting the extensive dissemination. This was followed by *C. koseri*, *Enterobacterspp* (5 sources), and *Proteus* (4 sources). In terms of the number and

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

Formatted: Space Before: 3 pt, After: 3 pt

percentage of isolates, it was discovered that *E. coli* had the highest isolates 19%, followed by *C. koseri* 16%, and *Proteus mirabilis* 12.3%. *Enterobacter* spp. and *Shigella* spp. each accounted for 10.5% of the total, Figure (3).

Table 3: Types of enteric bacteria isolated from different meat sources.

Source of sample	No. of isolates	Bacteria (no.)
Fresh chicken	6	<i>Citrobacterkoseri</i> (2), <i>E. coli</i> (1)
		<i>Enterobacterspp</i> (1), <i>Proteus mirabilis</i> (1)
		<i>Proteus vulgaris</i> (1)
Fresh beef meat	13	<i>Citrobacterkoseri</i> (2), <i>E. coli</i> (2)
		<i>Enterobacterspp</i> (1), <i>Proteus mirabilis</i> (4)
		<i>Proteus vulgaris</i> (1), <i>Providenciastuartii</i> (1)
		<i>Shigellaspp</i> (2)
Fresh sheep meat	7	<i>E. coli</i> (4), <i>Enterobacterspp</i> (1)
		<i>Proteus mirabilis</i> (2)
Fresh fish	4	<i>Citrobacterfreundii</i> (1), <i>E. coli</i> (1)
		<i>S.paratyphiA</i> (1), <i>Shigellaspp</i> (1)
Frozen red meat	10	<i>Citrobacterkoseri</i> (1), <i>E. coli</i> (2)
		<i>Enterobacterspp</i> (2), <i>Salmonellatyphi</i> (1)
		<i>Shigellaspp</i> (2), <i>Yersinia enterocolitica</i> (2)
Frozen beef pastrami	3	<i>E.coli</i> (1), <i>Proteus mirabilis</i> (1)
		<i>S.paratyphiA</i> (1)
Frozen chicken liver	3	<i>Citrobacterkoseri</i> (2), <i>Klebsiellapneumoniae</i> (1)
Frozen breast chicken	8	<i>Citrobacterfreundii</i> (2),
		<i>Citrobacterkoseri</i> (2), <i>Klebsiellaoxytoca</i> (1), <i>Proteus vulgaris</i> (1) <i>S.paratyphi A</i> (2)
Frozen beef burger	3	<i>Enterobacterspp</i> (1), <i>Shigellaspp</i> (1)
		<i>Klebsiella pneumoniae</i> (1)

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt




Figure 3: The proportions of isolating types of enterobacteria from the studied meat sources.

Members of the enteric family play a significant role in the transmission of foodborne illnesses. Foodborne infections continue to be a real and major health and economic hazard around the world [1]. According to the report of WHO in 2021, 600 million people worldwide suffer from foodborne infections each year, and 420,000 people die as a result of foodborne diseases, particularly antibiotic-resistant bacteria. The spread is frequently caused by fecal contamination of food sources, which contain several bacterial species [15].

Conclusions

Because it is obvious that meat can be infected with a wide range of dangerous bacteria, good hygiene procedures help to limit the quantity of contaminated germs. Furthermore, a commitment to take precautions when cutting and preserving meat, as well as applying sound manufacturing processes, is required to keep meat contamination to an acceptable level. Future research should concentrate on the precise characterization of the population structure of foodborne pathogens in order to better understand epidemiology, pathogenicity, and antibiotic resistance profiles.

Comment [PM3]: Can be improve

References

1. Lee, H. and Yoon, Y. (2021) 'Etiological agents implicated in foodborne illness worldwide', *Food Science of Animal Resources*, 41(1), pp. 1–7. doi: 10.5851/KOSFA.2020.E75.

2. Ebner, C., Morgan, A., & Manuel, C. (2021). Packaging of perishable food products. In *Food Safety and Quality-Based Shelf Life of Perishable Foods* (pp. 105-134). Springer, Cham.
3. Mustefa, W. S. (2021). Microbiological evaluation of meat sold in butcheries shop of Cheleleka town in ancharworeda, West Hareрге, Oromia, Ethiopia Western Ethiopia. *Journal of Food Science and Nutrition Therapy*, 7(1), 033-039.
4. Li, P. (2022) 'Foodborne Pathogens of Enterobacteriaceae, Their Detection and Control', in Jiang, H. (Ed.). Rijeka: IntechOpen, p. Ch. 10. doi: 10.5772/intechopen.102086.
5. Victorien, D., Phénix, A., Eugénie, A., Esther, D., Christina, D., Jerrold, A. et al (2020). Enterobacteria responsible for urinary infections: a review about pathogenicity, virulence factors and epidemiology.
6. Tribudi, Y. A., Lestari, O. A., Alfius, M., & Hidayatullah, A. (2020). Antibiotic Residues and Microbial Contamination in Animal-Derived Foodstuffs in Pontianak City, Indonesia. *Jurnal Ilmudan Teknologi Hasil Ternak (JITEK)*, 15(3), 152-161.
7. Aljamali, N. M. (2021). Review on food poisoning (types, causes, symptoms, diagnosis, treatment). *Global Academic Journal of Pharmacy and Drug Research*, 3(4), 54-61.
8. Gourama, H. (2020). Foodborne pathogens. In *Food safety engineering* (pp. 25-49). Springer, Cham.
9. Olu-Taiwo, M., Obeng, P. and Forson, A. O. (2021) 'Bacteriological Analysis of Raw Beef Retailed in Selected Open Markets in Accra, Ghana', *Journal of Food Quality*, 2021. doi: 10.1155/2021/6666683.
10. Procop, G. W., Church, D. L., Hall, G. S., Janda, W. M., Koneman, E. W., Schreckenberger, P. C., & Woods, G. L. (2018). Koneman's color atlas and textbook of diagnostic microbiology, pp. 29-60.
11. Atlas, R. and Snyder, J. (2015) 'Reagents, stains, and media: bacteriology', *Manual of clinical microbiology*, pp. 316-349.
12. Bersisa, A., Tulu, D., & Negera, C. (2019). Investigation of bacteriological quality of meat from abattoir and butcher shops in Bishoftu, Central Ethiopia. *International Journal of Microbiology*, 2019.
13. González, L. V. (2021) 'Human Pathogenic Enterobacteriales', *Reference Module in Biomedical Sciences*. doi: 10.1016/B978-0-12-818731-9.00160-9.
14. Lynch III, J. P., Clark, N. M. and Zhanel, G. G. (2021) 'Escalating antimicrobial resistance among Enterobacteriaceae: focus on carbapenemases', *Expert Opinion on Pharmacotherapy*, 22(11), pp. 1455-1474.
15. Yulistiani, R. and Praseptiangga, D. (2019) 'Contamination level and prevalence of foodborne pathogen Enterobacteriaceae in broiler and backyard chicken meats sold at traditional markets in Surabaya, Indonesia', *Malaysian Applied Biology*, 48(3), pp. 95-103.