

Public Awareness and prevalence of Depression Symptoms among Patients Attending Primary Health Care Centers in Qatar

Abstract

Background

Public awareness is a key element in the management of depression, public awareness campaigns have contributed to the improvement of public knowledge attitudes, and to reduction of stigma towards depression.

Aim

The aim is to identify public awareness towards depression and to estimate prevalence of depression among participants

Methods

Descriptive cross-sectional study, this study was conducted in (West Bay Health Center, Al-Rayan Health Center and Al-Daayen Health Center) affiliated to Primary Health Care Corporation in Qatar, it included all selected patients attending three randomly chosen health centers, with total response rate representing 82 % with total sample size 316 participant conducted during from January to March 2021, self-administered validated questionnaire including basic demographic data, awareness about symptoms of depression, risk factors, attitudes, treatment and help seeking behaviors also completing the PHQ-9 screening for depression questionnaire.

Results

The general awareness about depression among participants lies within the range of 48% to 95.7%, participants attitudes towards depression revealing that good attitudes regard willing to work with depressed people (88.9%) and willing to make friends with them (82%) while 39.4% agree to marry from them, participants preferences regard whom to seek advice regard depression revealing that in order psychologist (89.1%) then psychiatrist (87.8%) while family physicians (66.7%), PHQ-9 score among participants revealing average prevalence of depression reaching 19.7% at the cut point score of 10 with males reaching 17% and females 20.7%.

Conclusions

Generally, in Qatar the level of public awareness of depression among participants was acceptable and they had positive attitudes towards depressed people also there is some statistical significant awareness results related to females in comparison to males. However, further efforts must be done to build public health educational programs related to depression in order to further raise its awareness in the community.

Key words

Public, awareness, Prevalence, Depression, PHCC, Qatar

Introduction

The WHO global estimates suggest mental disorders affect more than 1 in 4 people in the course of their lives, and about 1 in 10 adults at any one time.(1). A study of Qatari nationals attending Primary Health Care Centers (2), revealed that the almost one-quarter of all Qatari adults who attended the primary health care setting presented with at least one type of mental disorder. In 2010 showing disease burden (3) revealing that: mental health disorders account for 15% of the global disease burden, Depression is considered the main cause of disability globally, Mental disorders represent four of the top ten causes of disability globally .Worldwide, mental health issues are well recognized as the most bothersome of all classes of disorders because of their high rate of prevalence, with early age of onset, its chronicity and associated functional impairment (4). In Qatar, a study of public perceptions of mental health issues conducted in Primary Health Care Centers (5) revealed that the level of mental disease knowledge in Qatar was low and that misconceptions exist around the causes of mental illnesses and also the capabilities of people with mental health problems. Stigma can deny individuals with a mental health issues from learning, employment, and fulfilling relationships.(6) In Qatar, some people have experienced negative impacts associated with mental illness, including reduced marriage concepts and employment opportunities, as well as social isolation. In most cases, primary health care physicians will provide the first point of contact for service users and their families. Integrating mental health services into primary health care systems can be an important solution to addressing human resource shortages to deliver mental health interventions (7) that is why primary health care workers will facilitate the early identification of mental problems and the treatment of mild common mental disorders such as depression and anxiety. Public awareness is a key element in the management of depression (8). Public awareness campaign has contributed to the improvement of public knowledge and attitudes, and to reduction of stigma towards depression (9). Almost every person, during their lifetime, will have direct contact with a mental disorder patient (10). Thus, knowledge about mental disorders is necessary as it directly affects a person's approach and behavior towards those patients (11). In this study, we will assess the degree of public awareness regarding major depression, its associated symptoms, contributing factors, believes and attitudes towards depressed patients, as well as available treatment options.

Methods

Study design: Descriptive cross-sectional study.

Study setting: This study was conducted in (West Bay Health Center, Alrayan Health Center and Aldayen Health Center) affiliated to Primary Health Care Corporation in Qatar

Study subjects: It included all selected patients attending three randomly chosen health centers. With total response rate representing 82 % with total sample size 316 participant conducted during from January to March 2021.

Data collection methods: Self-administered validated questionnaire including basic demographic data, awareness about symptoms of depression, risk factors, attitudes, treatment and help seeking behaviors also completing the PHQ-9 screening for depression questionnaire. For illiterate people there is help from an interviewer was done. The questionnaire is available upon request (msalem@phcc.gov.qa).

Data analysis: data were entered into a personal computer and were analyzed using SPSS version 15. Frequency analysis responses was carried out and the results were tabulated and compared using the chi-square goodness of fit test .Tests of significance (i.e. Fischer exact test) were applied .P-values less than 0.05 were considered as statistically significant.

Ethical considerations: This research project is approved from Institutional Review Board (IRB) in Primary Health Care Corporation, Qatar

Results

Table (1) shows the sociodemographic characters of participant where females represent 64.9% ,Qatari represents 27.7% ,married in 69.9% ,well educated in 95% of participant ,and governmental work represent 62.2%.

Table (2) shows the general awareness about depression among participants which reveals that participant are aware about tested items within the range of 48% to 95.7% and females agree with statistical significance regard treatment of depression by will power.

Table (3) shows the awareness regard symptoms of depression revealing that participant are aware about tested symptoms within a range of 71% to 93.5% with no statistical significant difference between males and females.

Table (4) shows the awareness regard risk factors about depression revealing that participants are aware about tested items within a range of 54.1% to 85.6% with no statistical significance differences between males and females.

Table (5) shows the awareness regard treatment of depression revealing that participants are aware of tested treatment options in a range of 49.5% to 98.7% and females agree with statistical significance regard choosing vitamins and analgesics as options of treatment.

Table (6) shows participants attitudes towards depression revealing that good attitudes regard willing to work with them (88.9%) and willing to make friends with them (82%) while 39.4% agree to marry from them and 10.6% consider them mentally retarded also males agree with statistical significance regard consideration of mental retardation among depressed people.

Table (7) shows awareness regard whom to seek advice regard depression revealing that in order psychologist (89.1%) then psychiatrist (87.8%) while family physicians (66.7%) and females agree in statistical significance regard psychiatrist seeking and self-help.

Figure (1) shows PHQ-9 score among males and females revealing nearby score in moderate depression score, predominant female in moderately severe category and predominant male in severe category. It also shows PHQ-9 score among participants revealing average prevalence of depression reaching 19.7% at the cut point score of 10 with males reaching 17% and females 20.7%

Figure (2) shows overall PHQ-9 score among participants with total prevalence of 19.7% at cut point of 10 (moderate, moderately severe and severe depression)

Discussion

The sociodemographic characters of participant where that females represent 64.9%, Qatari represents 27.7%, and married in 69.9%, well educated in 95% of participant, and governmental work represent 62.2%.

This study indicated acceptable public awareness about major depression, its symptoms, risk factors, available treatments, and attitudes. With female respondents having higher degree of awareness regard depression, this matched study done in Jordan revealing significant higher degree of awareness among females regard depression compared to male respondents in Jordanian study (12). Women and younger people have more substantial knowledge about key aspects of depression and its treatments in Australian study (13).

Regard awareness towards treatment options despite good awareness score still females believe in vitamins and analgesics as modality of treatment which may be due to beliefs that drugs for depression may have side effects or addiction properties (14) another belief in our study reveal that females agree with statistical significance regard treatment of depression by will power. That is why we have to change these beliefs about treatment of depression by different methods through campaigns or communication with public through social media emphasizing need for support highlighting role of support clinics in Primary Health Care Corporation.

Participant attitudes regard low rate of agreement regard marriage from depressed in addition to believing that they may have mental retardation (significant towards male respondents), these negative attitudes may underestimate the depressive disorders in the community and clinical setting (15) this in concordance with Turkish study (16) revealing female respondents had less stigmatizing attitudes.

Participants agree upon considering family and friends as first choice for support in case of depressive symptoms followed by professionals (psychologists and psychiatrists) with low rank in comparison regard family physician and pharmacists, this is in contradictory with the Australian study which reveal choice of general practitioners as the first choice (13), this difference actually may be due to cultural issues favoring family support in Arab communities. But we have to raise awareness regard mental health services in the community and role of family physicians as key leaders in managing mental health issues with appropriate knowledge and appropriate referral systems in place.

In this study, PHQ-9 score among participants revealing average prevalence of depression reaching 19.7% at the cut point score of 10 with males reaching 17% and females 20.7%, This matches study done in Qatar (17) revealing that the most common ICD-10 disorders were specific generalized anxiety disorders (20.4) %, and major depression (19.1%), with a higher prevalence in women, this agreement about the prevalence despite use different tools and study population confirm that the prevalence is actually around 19% which warrants further efforts to tackle the problem and its consequences ,another important observation is that the prevalence is constant from 2014 to 2022 despite changing community and challenges such as covid-19 epidemics .

Conclusions

Generally, in Qatar the level of public awareness of depression among participants was acceptable and they had positive attitudes towards depressed people also there is some statistical significant awareness results related to females in comparison to males. However, further efforts must be done to build public health educational programs related to depression in order to further raise its awareness in the community.

Limitations

The limitations of the study could be related to small sample size in comparison to other studies ,there is some participants who cannot read and write so they need help in completing the questionnaire ,this study include Qatari and non-Qatari people as well which may affect the results in the form of culture differences ,the female sampled population were more than males which may give result differences ,finally the dependence upon PHQ-9 questionnaire as key in depression diagnosis may be better done after face to face consultations despite referral to mental health services was done for whom score revealed moderate or severe depression.

Recommendations

We still have room of improvement dealing with mental health problems specially depression and its awareness by focusing on repeated audits regard community awareness and also correcting beliefs regard issues related to stigmatization and modalities of treatment, also modify help seeking behaviors favoring family physicians consultation and early recognition of depressive symptoms before affecting persons quality of life .

Table (1) Sociodemographic data among sampled population

Demographic data	N	%
Age (in years)		
18-30	79	25.30%
31-40	107	34.30%
41-50	76	24.40%
>50	50	16.00%
Total	312	
Gender		
Female	198	64.90%
Male	107	35.10%
Total	305	
Nationality		
Qatari	86	27.70%
Non-Qatari	225	72.30%
Total	311	
Social status		
Single	76	26.00%
Married	204	69.90%
Divorce or widow	12	4.10%
Total	292	
Education level		
literate	6	1.90%
Primary	12	3.80%
Secondary	70	22.30%
Bachelor	149	47.50%
Postgraduate	77	24.50%
Total	314	
Occupation		
Governmental	178	62.20%
Non-governmental	58	20.30%
Not employed/housewife/retired	50	17.50%
Total	286	
Income		
Enough	193	68.40%
Not enough	89	31.60%
Total	282	

Table (2) General awareness about depression among sampled population

General Awareness	N (%)			p-value
	Total	Male	Female	
Depression is a treatable disease	286 (95.7)	99 (95.2)	187 (95.9)	0.772*
Depression can be treatable by will power	258 (88.7)	81 (81.0)	177 (92.7)	0.003
Depression may affect any age	257 (88.3)	86 (86.0)	171 (89.5)	0.373
Causes of depression are well known scientifically	177 (62.1)	58 (58.6)	119 (64.0)	0.372
Depression run in families	137 (48.2)	52 (53.1)	85 (45.7)	0.238

*Fisher's exact test

Table (3) Awareness about symptoms of depression among sampled population

Symptoms	n (%)			p-value
	Total	Male	Female	
Loss of interest	256 (88.0)	90 (89.1)	166 (87.4)	0.664
negative feelings	267 (92.7)	92 (92.0)	175 (93.1)	0.736
sadness	274 (93.5)	92 (91.1)	182 (94.8)	0.221
change in behaviours	268 (92.4)	99 (96.1)	169 (90.4)	0.077
drug abuse	198 (71.0)	71 (72.4)	127 (70.2)	0.688
suicide	225 (80.1)	79 (81.4)	146 (79.3)	0.676

Table (4) Awareness about risk factors of depression among sampled population

Risk Factors	n (%)			p-value
	Total	Male	Female	
Unemployment	250 (85.6)	88 (87.1)	162 (84.8)	0.592
poverty	231 (81.1)	80 (81.6)	151 (80.7)	0.856
drug abuse	222 (80.7)	74 (78.7)	148 (81.8)	0.544
divorce	218 (78.1)	74 (77.1)	144 (78.7)	0.758
heavy work	208 (73.0)	70 (69.3)	138 (75.0)	0.301
severe disease	235 (84.8)	84 (87.5)	151 (83.4)	0.368
marriage	146 (54.1)	44 (48.4)	102 (57.0)	0.179
others	152 (76.0)	50 (78.1)	102 (75.0)	0.629

Table (5) Awareness of treatment of depression among sampled population

Treatments	n (%)			p-value
	Total	Male	Female	
family support	295 (98.7)	102 (98.1)	193 (99.0)	0.612*
exercise	282 (95.6)	94 (93.1)	188 (96.9)	0.143*
antidepressants	233 (81.2)	82 (82.8)	151 (80.3)	0.605
vitamins	173 (62.2)	50 (52.6)	123 (67.2)	0.017
herbs	135 (49.5)	41 (45.6)	94 (51.4)	0.367
analgesics	109 (40.7)	28 (30.8)	81 (45.8)	0.018

*Fisher's exact test

Table (6) Public attitude towards depression

Public Attitude	n (%)			p-value
	Total	Male	Female	
I am willing to work with them	264 (88.9)	93 (88.6)	171 (89.1)	0.898
I am willing to make friends with them	237 (82.0)	80 (80.0)	157 (83.1)	0.518
I am willing to marry one with depression	108 (39.4)	43 (43.9)	65 (36.9)	0.259
they are mentally retarded	30 (10.6)	16 (16.0)	14 (7.7)	0.029

Table (7) Awareness about whom to seek for help among sampled population

Person to seek help from	n (%)			p-value
	Total	Male	Female	
family and friends	267 (94.3)	92 (93.9)	175 (94.6)	0.804
psychiatrists	231 (87.8)	76 (81.7)	155 (91.2)	0.025
self help	207 (80.5)	61 (68.5)	146 (86.9)	<0.001
psychologist	228 (89.1)	78 (86.7)	150 (90.4)	0.366
family physician	162 (66.7)	54 (63.5)	108 (68.4)	0.447
pharmacist	70 (29.8)	21 (25.9)	49 (31.8)	0.348

Figure (1) : PHQ – 9 score distribution between females and males





Figure (2) PHQ-9 score among sampled population



References

1. The world health report 2001: Mental Disorders affect one in four people, treatment available but not being used: 28th September 2001.
2. Ghuloum S, Bener A, Abou-Saleh MT. Prevalence of mental disorders in adult population attending primary health care setting in Qatari population. *J Pak Med Assoc.* 2011 Mar;61(3):216-21. PMID: 21465930.
3. Murray, C. et al, 2012, Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990—2010: a systematic analysis for the Global Burden of Disease Study 2010, *The Lancet*, Volume 380, Issue 9859, 15/22/29, December 2012.
4. Bener, A. et al, 2012: Measuring burden of diseases in a rapidly developing economy: state of Qatar, *Global Journal of Health Sciences*, Volume 5, Issue 2, pp 134 – 144, 21 December 2012.
5. Ghuloum S, Bener A & Burghut FT, 2011, Ethnic Differences in the Knowledge, Attitude and Beliefs towards Mental Illness in Traditional Fast Developing Country, *Psychiatria Danubina* Vol 23. No 2. 157-164
6. Sartorius N. Stigma: what can psychiatrists do about it? *Lancet.* 1998 Sep 26;352(9133):1058-9. doi: 10.1016/S0140-6736(98)08008-8. PMID: 9759771.
7. Funk M, Saraceno B, Drew N, Faydi E. Integrating mental health into primary healthcare. *Ment Health Fam Med.* 2008 Mar;5(1):5-8. PMID: 22477840; PMCID: PMC2777555.
8. Gargiulo A, Ebmeier KP. Awareness key in managing depression in later life. *Practitioner.* 2008;252(1705):32,34,37 (passim).
9. Jones AR, Cook TM, Wang J. Rural-urban differences in stigma against depression and agreement with health professionals about treatment. *J Affect Disord.* 2011;134 (1–3):145–50.
10. Kessler RC, McGonagle KA, Zhao S, Nelson CB, Hughes M, Eshleman S, et al. Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States. Results from the National Comorbidity Survey. *Arch Gen Psychiatry.* 1994;51(1):8–19.
11. Jorm AF. Mental health literacy. Public knowledge and beliefs about mental disorders. *Br J Psychiatry.* 2000;177: 396–401.
12. Al-Azzam S, Alzoubi KH, Ayoub NM, Khader YS, Karasneh RA, Al-Husein BA, Al-Horani MA. An audit on public awareness of depression symptoms in Jordan. *Int J Occup Med Environ Health.* 2013 Aug;26(4):545-54. doi: 10.2478/s13382-013-0128-9. Epub 2014 Jan 25. PMID: 24052153.
13. Highet NJ, Hickie IB, Davenport TA. Monitoring awareness of and attitudes to depression in Australia. *Med J Aust.* 2002;176 Suppl:S63–8.

14. Jorm AF, Nakane Y, Christensen H, Yoshioka K, Griffiths KM, Wata Y. Public beliefs about treatment and outcome of mental disorders: A comparison of Australia and Japan. *BMC Med.* 2005;3:12.
15. Postuvan V, Becaj J, Marusic A. Nurses' attitudes towards depression: a study in Slovenia. *Psychiatr Danub* 2007;19(1-2):61-7.
16. Savrun BM, Arıkan K, Uysal O, Cetin G, Poyraz BC, Aksoy C, et al. Gender effect on attitudes towards the mentally ill: A survey of Turkish university students. *Isr J Psychiatry Relat Sci.* 2007;44(1):57-61.
17. Suhaila Ghuloum, Abdulbari Bener, Elnour E. Dafeeah, Tariq Al-Yazidi, Ahmed ElAmin Mustapha and Ahmed E. Zakareia. Lifetime Prevalence of Common Mental Disorders in Qatar: Using WHO Composite International Diagnostic Interview (WHO-CIDI) *International Journal of Clinical Psychiatry and Mental Health*, 2014, Vol. 2, No. 1 9