

Case study

Bacteriological Profile and their drug sensitivity profile in Diabetic Foot Ulcer in a Tertiary Care Center

Comment [FG1]: Change to "Identification and Antimicrobial Resistant Pattern of Bacteria Isolated from Diabetic Foot Ulcer Cases in Tertiary Care Center, Pune"

Abstract-

Aim: The present study was an attempt to evaluate the different microorganisms infecting the Diabetic foot ulcers (DFU). The aim of the current study was to compare the different bacteria infecting Diabetic Foot Ulcers. And to know the antibiotic susceptibility patterns among the isolates.

Study design- This is a Prospective observational study of patients treated at Tertiary health Care Centre, Pune.

Type of study- Prospective and observational hospital-based study

Period of study- From February 2021 to January 2022

Sample size- Tissue culture samples were collected from 100 patients.

Conclusion: This study demonstrated that among the isolates from the DFUs, multidrug-resistant bacteria predominated. Determining the antibiotics for the empirical therapy of diabetic ulcers will be made easier with knowledge of the pattern of antibiotics resistance among the isolates. Thus, the likelihood of subsequent development of antibiotic resistance as well as the indiscriminate use of antibiotics can be reduced.

Keywords: Diabetic Foot ulcers, Bacteriological profile, Drug sensitivity profile

Comment [FG2]: Poorly written. Please rewrite according to guideline. The Key findings in Results were not mentioned.

INTRODUCTION

Diabetic foot ulcer is a dangerous and common complication of diabetes mellitus (DM) that significantly increases the cost of treatment.[1]

Diabetic foot ulcers were found in 4.54% of patients newly diagnosed with type 2 diabetes mellitus in India; of these, 46.1% had neuropathic, 19.7% had ischemic, and 34.2% had neuroischemic foot ulcers.[2] Infections, which account for 40% to 80% of instances of DFU morbidity and mortality, are most frequent complication in DFU.[3]

Poor microvascular circulation prevents phagocytic cells from reaching the infected location, which impairs the effectiveness of antibiotics in the infected tissue.[4]

Infection causes the development of microthrombi, which aggravate ischemia, necrosis, and progressive gangrene necessitating limb amputation.

DFUs are chronic in nature and patients with DFUs usually require several episodes of hospitalization. Patients are often exposed to several antibiotics which increase their risk of developing multidrug-resistant infection.[5]

Mostly, the diabetic foot infections (DFIs) are mixed bacterial infections, and the proper antibiotic selection is necessary for the treatment of these infections., based on the culture and the antimicrobial susceptibility testing results.

AIM

The aim of the current study was to assess the various bacteria infecting the DFU and to know the antibiotic susceptibility patterns to these isolates.

METHODS

Inclusion criteria-

All the diabetic patients who attended the outpatient department of the study center with foot ulcer or infection.

Exclusion criteria-

Other foot ulcers and foot infection in persons without diabetes

The Institutional Ethical Committee's approval was obtained prior to conducting the study.

A clinical history was obtained in relation to the Patients with diabetes(PWD's) demographics, the length of the diabetes and foot condition, the type of diabetes treatment previously received, and the existence of any systemic disorders.

PWD underwent clinical evaluation as well, and the foot ulcers were rated using Wagner's grade (Wagner and Meggitt, 1970):

- 0 - No ulceration in a high-risk foot
- 1 - Superficial ulcer of skin or subcutaneous tissue
- 2 - Ulcers extend into tendon, bone, or capsule
- 3 - Deep ulcer with osteomyelitis or abscess

Comment [FG3]: Before you mention the effect of foot ulcers, you need to mention what are the common bacteria that infect the lesion if not in India, in other Asian countries or other part of the world, not to forget to mention what was the current Antibiotic treatment used to treat this condition.

Comment [FG7]: Wrong font and size, Sentences are hanging please rewrite.

Comment [FG4]: Do you mean to identify

Comment [FG5]: Does your study have Human ethic clearance or approval? Please indicate the clearance number

Comment [FG6]: What is the sampling size, how many samples were taken from how many people? What are their demographic profiles?

4 - Gangrene of toes or forefoot (localized gangrene)

5 - Extensive gangrene requiring a major amputation.

Based on the presence of neuropathy, ischemia, and infection, ulcer foot type was determined. Investigations such as monofilament nerve conduction velocity testing, biothesiometry, and Doppler-based ankle brachial index estimation were carried

out for this in addition to the clinical history and examination of PWD.

After the debridement, tissue samples were taken. Before obtaining a tissue sample, no antimicrobial or antiseptic agent was used to the wound. Empirical broad spectrum antibiotic coverage was started for every patient with DFU according to institutional protocol.

A deep tissue specimen (including fat, fascia, muscles, and bone) was also taken from the wound. The samples were put into sterile transport containers and delivered to the microbiology laboratory for aerobic microbial culture as soon as possible.

Anaerobic and fungal cultures were not performed for this study.

Most of the bacterial isolates were identified using VITEK 2 Compact system, and a few isolates were identified manually.[6]A direct Gram-stained smear of the specimen was examined.

The specimens were inoculated onto blood agar, chocolate agar, Mac Conkey's agar, and thioglycollate medium. The inoculated plates were incubated at a temperature of 37°C overnight, and the plates were examined for growth on the following day.

The organisms were identified on the basis of their Gram-staining properties, and further analysis was done in VITEK® 2 Compact system (BioMérieux).

Antibiotic susceptibility testing

A bacterial suspension was matched with McFarland standard of 0.5ml in 2.5 ml of a 0.45% sodium chloride solution with a VITEK® 2 DensiChek instrument (BioMérieux) with the incubation temperature kept at 35.5°C.

The isolates were subjected to a colorimetric measurement using a fresh optical reading head every 15 minutes for a maximum incubation time of 10 hours[7]

Comment [FG12]: What do you mean by manually? Phenotypic? Genotypic? Biochemistry?

Comment [FG8]: Should start at a new subheading on tissue sampling

Comment [FG13]: Model; year

Comment [FG9]: What is the name of the antibiotics used

Comment [FG10]: What is the size or quantity of specimen taken eg in cm/mm or in gram etc.

Comment [FG11]: As soon as possible is subjective. Be specific on the duration eg. Within 2 hours, within 24 hours etc.

VITEK® 2 database version 4.01 was used to analyze the data for organism identification in kinetic mode after 2 h of incubation. The interpretations provided were then considered for the analysis.

RESULTS

In the present study, out of 100 PWD, 38 patients were below 50 years and 62 patients were above 50 years.

In our study, 81 patients were males and 19 were females.

In the present study, Grade I DFU was seen in 6 (6%) PWD, Grade II in 21 (21%), Grade III in 48 (48%), Grade IV in 21 (21%), and Grade V in 4 (4%).

In this study, 62 (62%) PWD cases had neuropathic conditions, 18 (18%) had neuropathic cases combined with sepsis, 11 (11%) had neuroischemic conditions, and 9 (9%) had neuroischemia plus sepsis.

In this study, there were 29 (29%) polymicrobial cases, 64 (64%) monomicrobial cases, and in 7 (7%) cases, the culture was sterile.

Gram-positive bacterial growths were present in 41(41%) cases, whereas Gram-negative growth was seen in 59 (59%) cases.

The most common single bacterial growth was that of *S. aureus* (27%), followed by *E. coli* (20%), and *Enterococcus* spp. (15%).53% of the Gram-negative bacilli

were Extended spectrum beta lactamase (ESBL) producers, 41% were Methicillin resistant staphylococcus aureus (MRSA), and 19% were Vancomycin resistant enterococci(VRE).

Comment [FG14]: How many antibiotic were tested? how many class of antibiotic? Which guidelines were followed? CLSI?

Bacterial sensitivity pattern is obtained as follow:

Antibiotic	Sensitivity pattern (%)		
	<i>Staphylococcus aureus</i>	<i>Enterococcus</i>	<i>Enterobacteriaceae</i>
Ampicillin			11
Amoxicillin-clavulanic acid			64
Piperacillin-tazobactam			73
Cefalacin			24
Ceftriaxone			56
Cefoxitin			26
Cefixime			11
Ertapenem			76
Ofloxacin			42
Ticarcillin-clavulanic acid			14
Ceftazidime			64
Cefoperazone-sulbactam			
Cefepime			
Doripenem			
Imipenem			89
Meropenem			84
Amikacin			90
Aztreonam			
Gentamicin	83		89
Ciprofloxacin	73	74	65
Minoscylcline			
Tigecycline			
Trimethoprim-sulfamethoxazole		47	39
Levofloxacin	74	68	
Colistin			
Oxacillin	72		
Erythromycin	78	70	
Clindamycin	71	58	
Linezolid	100	100	
Daptomycin	100	100	
Teicoplanin	84	89	
Vancomycin	100	67	
Benzylpenicillin	24	11	
Tetracycline		82	
Tigecycline	89	74	

Most of the *Staphylococcus* culture isolates were sensitive to linezolid (100%), daptomycin (100%), tigecycline (89%)

In our study, most of the *Enterococcus* culture isolates were sensitive to linezolid (100%), daptomycin (100%), teicoplanin (89%), and tigecycline (74%)

DISCUSSION

The majority of the PWD in this study were over 50 years old (63%). This could be a sign of comorbidities such neuropathy, peripheral vascular disease, and kidney illnesses being more common in this age group. A study by King et al. in 1998 also mentioned that the majority of people with diabetic foot were in 45–64 years.[8]

In the present study, most of the *Enterobacteriaceae* culture isolates were sensitive to amikacin (90%), imipenem (89%), meropenem (84%)

Most of the *Pseudomonas* culture isolates were sensitive to amikacin (90%), imipenem (72%), meropenem (70%).

Higher male prevalence is comparable with study by Harrison and Lederberg. This might be because men engage in more outdoor physical activity than women, especially in hot, humid environments, with poor foot care.[9]

Our study found that the majority of DFI patients reported having an advanced grade of infection, Wagner Grade III and above. This is frequently ascribed to the public's

Comment [FG15]: Not clear, unfortunately I can't read. Please replace with a better and higher resolution figure

Comment [FG16]: Should include the MARI results

Comment [FG17]: It would be beneficial to present the antibiogram in a graph

Comment [FG18]: The ref you cited is 25 years ago, there more recent and better studies conducted that should be cited

and medical professionals' lack of knowledge on foot care.

S. aureus was the single most frequent pathogen (26%) followed by *E. coli* (20%). Other studies have also found the same (study by Abdulrazaq et al.) In contrast, another study carried out by Ako-Nai et al. [10] showed *E. coli* as the frequent bacterial pathogen, while *P. aeruginosa* was reported as the most common pathogen by Shankar et al.[11] Source of infection, use of antibiotic drug for treatment, sample collection method, and different types of infection can influence pathogen diversity in DFI.

While GPC were more prevalent in Grades I and II, Gram-negative bacilli and mixed infections were more evident in Grades III and IV, suggesting that Gram-negative infections increase the severity and render patients more likely to require limb amputation.

These days, the rising threat of MDR pathogens and related consequences in developing nations worries clinical microbiologists and doctors. [12] In the current investigation, 91 percent of the bacteria (VRE 33.33 percent, MRSA 48.14 percent, and ESBL 77.67 percent) were resistant to three or more antibiotics. In

contrast to an Iranian study by Japoni et al, these rates are much higher. These isolate infections are more challenging to treat.

Frequent hospitalization, frequent use of broad-spectrum antibiotics, insufficient surgical source reduction, chronic wounds, irrational use of antibiotics, and the transmission of resistance genes via transport methods are possible causes of MDR. Clinicians should use antibiotics judiciously, on time, and in sufficient amounts, and the relevant organizations should periodically monitor drug intake in order to improve the condition and lower the rate of amputation.

Clinicians should switch to using narrower spectrum therapy depending on culture report. To reduce infection sources, a sufficient and prompt surgical intervention is necessary. These aid in lowering the excessive and careless use of antibiotics.

CONCLUSION

This study demonstrated that Gram-negative aerobes like *S. aureus* were the most frequent microbes found in diabetic foot ulcers.

In the DFI cases, monomicrobial infection was more prevalent than polymicrobial infection.

Comment [FG20]: You need to justify

Formatted: Font: Italic

Formatted: Font: Italic

Comment [FG19]: Please italic all the bacteria spp mentioned in this manuscript.

Formatted: Font: Italic

Comment [FG21]: S aureus is Gram positive.

Formatted: Font: Italic

MDR organisms were alarmingly prevalent in the PWD and in people with foot ulcers.

According to local sensitivity patterns ideal empirical antibiotics combination for Diabetic foot ulcers in our institution is Linezolid and Amikacin which is most effective in cohort of patients with presentation of infections associated with DFU. In the present study, 91% of the bacteria were resistant to three or more antibiotics. Thus, indiscriminate use of antibiotics and chances of subsequent development of antibiotic resistance can also be reduced with proper knowledge of antibiotics sensitivity.

REFERENCES

1. Lipsky BA. A report from the international consensus on diagnosing and treating the infected diabetic foot. *Diabetes/metabolism research and reviews*. 2004 May;20(S1):S68-77.
2. Singh N, Armstrong DG, Lipsky BA. Preventing foot ulcers in patients with diabetes. *Jama*. 2005 Jan 12;293(2):217-28.
3. Richard JL, Sotto A, Lavigne JP. New insights in diabetic foot infection. *World journal of diabetes*. 2011 Feb 2;2(2):24.
4. Bronze MS, Khardori R, editor. *Diabetic foot infections treatment and management*. Medscape; 2016. Available form: <http://emedicine.medscape.com/article/237378-treatment>

5. Zubair M, Malik A, Ahmad J. Clinico-bacteriology and risk factors for the diabetic foot infection with multidrug resistant microorganisms in north India. *Biol Med*. 2010;2(4):22-34.
6. Garcia-Garrote F, Cercenado E, Bouza E. Evaluation of a new system, VITEK 2, for identification and antimicrobial susceptibility testing of enterococci. *Journal of clinical microbiology*. 2000 Jun 1;38(6):2108-11.
7. David HP, Pincus DH. Microbial identification using the BioMérieux VITEK® 2 System. BioMérieux, Inc. Hazelwood, MO, USA. 2009.
8. King H, Aubert RE, Herman WH. Global burden of diabetes, 1995–2025: prevalence, numerical estimates, and projections. *Diabetes care*. 1998 Sep 1;21(9):1414-31.
9. Lederberg J, Harrison PF, editors. *Antimicrobial resistance: issues and options*.
10. Ako-Nai AK, Ikem IC, Akinloye OO, Aboderin AO, Ikem RT, Kassim OO. Characterization of bacterial isolates from diabetic foot infections in Ile-Ife, Southwestern Nigeria. *The Foot*. 2006 Sep 1;16(3):158-64.
11. Shankar EM, Mohan V, Premalatha G, Srinivasan RS, Usha AR. Bacterial etiology of diabetic foot infections in South India. *European journal of internal medicine*. 2005 Dec 1;16(8):567-70.
12. Japoni A, Vazin A, Hamedi M, Davarpanah MA, Alborzi A, Razaatpour N. Multidrug-resistant bacteria isolated from intensive-care-unit patient samples. *Brazilian Journal of Infectious Diseases*. 2009;13:118-22.

Comment [FG22]: Improper conclusion and lack of specificity. It is advice to rewrite a stronger conclusion and recommendation.

Comment [FG23]: Check and rewrite the format of the reference listed