

Anti-hyperlipidemic Effect of Methanol Seed Kernel Extract of *Mangifera indica* on Wistar Rats fed High Fat Diet

Abstract

The aim of this study was to evaluate the hyperlipidemic effect of methanol seed kernel extract of *Mangifera indica* on wistar rats. Mango seed kernels were dried at room temperature before being ground to fine powder. 500 g of mango seed kernel powder was soaked in 500 ml of methanol and shaken intermittently for 72h after which the extract was concentrated. Twenty five adult male wistar rats were divided into five groups of five rats each. **Group I** was the normal control and was fed only normal rat chow. **Groups II-V** was induced hyperlipidemia, however, while Groups II was not treated with the extract (negative control), **Groups III and IV** were treated with 150 and 350 mg/kg body weight of methanol seed kernel extracts of *Mangifera indica* and **Group V** the standard drug (atorvastatin). Treatment lasted for 21 days after which rats were sacrificed and blood sample collected subsequently analysed via standard procedures. Hyperlipidemia was characterized by increased levels of total cholesterol and Low Density Lipoprotein (LDL). Oral administration of methanol seed kernel extract of *Mangifera indica* significantly ($P < 0.05$) reduced the aforementioned indices to levels which though were significantly ($P < 0.05$) higher than that reported for the normal control group. On the other hand, it was observed that the levels of high density lipoprotein (HDL) and triacylglyceride (TG) in the negative control (**Group II**) were significantly low but increased following oral administration of extract in a dose dependent manner. It was also observed that extract reduced the weight of hyperlipidemic rats.

Keywords: Lipoprotein, Hyperlipidemia, *Mangifera indica*, Atorvastatin, Kernel

Introduction

Hyperlipidemia is a secondary metabolic dysregulation orchestrated by enhanced levels of plasma lipid including primarily total cholesterol, triglyceride and Low Density Lipoprotein alongside decreased High Density Lipoprotein reason for which atherosclerosis is initiated and progressed [1]. Increased serum levels of triglyceride, cholesterol and LDL have been identified as core risk factors for the untimely development of cardiovascular diseases like hypertension and coronary

Comment [PM1]: Pure ?

Comment [PM2]: Ne female why ?

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Bold

heart disease [2] and have been traced to increased uptake of lipid via the gut or enhanced endogenous synthesis of the said molecules.

Formatted: Font: Italic

The use of plants with therapeutic significance has been an integral component of the human health care system since prehistoric times and interest in the practice has improved tremendously, evident by the fact that an estimated 80% global population depend on it to be relieved of one disease or the other [3].

Mango is of the genus *Mangifera* consisting of 30 species of tropical fruiting trees in the flowering family of *Anacardiaceae*. Botanically, it is called *Mangifera indica*. Its application in Ayurvedic medicine dates back to more 4000 years ago. Ayurveda was able to establish that the various parts of this tree have varied medicinal properties. Parts of the tree is rich in mangiferin, a polyphenolic antioxidant with, anti-lipid peroxidation, immunomodulation, cardiogenic, hypotensive, wound healing and anti-degenerative and anti-diabetic properties [4].

Although, it has been established that different parts of *Mangifera indica* (mango) have unique therapeutic ability, it is yet to be known whether or not the seed could be useful in the treatment of hyperlipidemia, hence, the imperativeness of this study is determined.

Materials and Methods

Collection and processing of plant material

Mango (*Mangifera indica*) seeds were identified at the herbarium unit of the department of Biological Science, Ahmadu Bello University Zaria Kaduna State. The seed kernel were subsequently dislodged and afterwards dried at room temperature for ten days. The dried seed kernels were pulverized with the aid of mortar and pestle and sieved to obtain a fine powder. The 500g of powdered plant sample was introduced into a conical flask containing 500mL of

methanol and shaken intermittently for 3 days. The extract was dried in water bath below 40°C.

describe final extract ... no methanol trace ?....

Comment [PM3]: Pure?

Comment [PM4]: No try other solvent?
Ethanol ?

Comment [PM5]: Temperature ?why not
heat it?

Animals

Adult male wistar rats weighing 120-160g were procured from the animal house of the Department of Science Laboratory Technology, AkanuIbiam Federal Polytechnic UnwanaAfikpo, Ebonyi State. The rats were housed in well ventilated cages under standard laboratory conditions and were maintained at ambient temperature and relative humidity respectively. They were fed growers mash (Vital feeds Nigeria Ltd) and provided with water. Acclimatization lasted for two weeks after which experiment commenced.

Comment [PM6]: No female why ?

Comment [PM7]: What happens to the rats
after the experiment?

Preparation of High Fat Diets (HFD)

The high fat diet was prepared by mixing 60 ml of cholesterol with 5 g of rat chow feed. (Hassarajani *et al.*, 2007).

Comment [PM8]: ?

Animal grouping

A total number of twenty five adult male wistar rats were for this study. Animals were divided into five groups of rats per group.

Group I (Normal group): animals were allowed access to rat chow and water

Group II (Negative control): animals were induced hyperlipidemia without treatment

Group III: Hyperlipidemic rats were administered with 150mg/kg of MSKE.

Comment [PM9]: define

Group IV: Hyperlipidemic rats were administered with 350mg/kg of MSKE.

Group V: Hyperlipidemic rats were administered with 4 mg/kg of atorvastatin.

Treatment lasted for 21 days treatment after which rats were anaesthetized by exposure to chloroform, sacrificed and blood sample collected in plain bottles and afterwards centrifuged at 500 rpm for 10 minutes.

Biochemical assays for lipids

Cholesterol, HDL and triacylglyceride levels were estimated from serum by CHOD-PAP according to the method of Devi and Sharma, (2004), respectively. LDL and HDL were calculated using the method by Johnson *et al.* (1997). While the artherogenic index was calculated by using the method described by Muruganandan et al. (2005)

Body Weight Measurement

The weight of the rats was determined prior to hyperlipidemia induction, after induction and after treatment. The changes in bodyweight was calculated and recorded.

Statistical analysis

Data generated from the study was analyzed using statisticssoftware IBM SPSS Statistics 21. Data were expressed as mean \pm standardeviation (SD). The results were considered as significant at $P<0.05$. Mean values were compared using one way analysis of variance (ANOVA).

Table 1: Lipid Profile of Hyperlipidemic Rats administered with Methanol Leaf Extract of *M. indica*

Comment [PM10]: explain with few sentences

Groups	Treatment	TC (mg/dl)	HDL (mg/dl)	LDL (mg/dl)	TG (mg/dl)
Group I	Normal control (NC)	60.23±2.37 ^d	19.11±5.09 ^a	56.12±2.41 ^d	24.62±4.01 ^{bc}
Group II	Negative Control	76.10±3.89 ^c	13.21±2.01 ^{ab}	58.50±3.29 ^d	19.12±1.85 ^a
Group III	150mg/kg MLEMI	64.21±5.54 ^{bc}	21.13±2.32 ^b	41.20±4.86 ^b	23.20±1.64 ^c
Group IV	300mg/kg MLEMI	63.10±2.50 ^a	26.12±2.30 ^c	43.5±4.498 ^{bc}	28.21±1.03 ^b
Group V	Atovarstatin	60.30±2.30 ^b	22.12±3.21 ^b	23.21±1.98 ^a	28.91±3.24 ^a

Results are expressed as mean ± standard deviation of three determinations. Values with the same superscript in column are significantly at P≤0.05.

Formatted: Font: 9 pt

Table 1 & 2 in resultats discussion part

Table 2: Effect of Methanol Leaf Extract of *M. indica* on the weight of Hyperlipidemic Rats

Comment [PM11]: explain with few sentences

Groups	Treatment	Wt. before induction	Wt. after induction	Wt. after treatment
Group I	Normal control (NC)	145.62±2.34	180.12±3.78	150.65±6.30
Group II	Negative Control	110.11±6.39	161.29±3.75	130.63±5.80
Group III	150mg/kg MSKE	102.43±5.34	147.13±4.57	104.12±3.28
Group IV	300mg/kg MSKE	128.56±3.46	185.60±5.01	150.21±3.46
Group V	Atovarstatin	120.22±2.32	176.34±3.60	138.34±6.90

Results are expressed as mean ± standard deviation of three determinations.

Formatted: Font: 9 pt

Result and Discussions

Comment [PM12]: must be improve

For the levels of lipid to be elevated, it is either, the amount of lipid absorbed through the gut is increased or the endogenous synthesis of lipid is enhanced. Thus, in order to reduce

hyperlipidemia, it is either the endogenous synthesis is blocked or absorption is decreased. Table 1 shows the lipid profile of hyperlipidemic rats administered with methanol leaf extract of *Mangifera indica* (mango) indicating that feeding high fat diet on animals significantly increased the level of total cholesterol and Low Density Lipoprotein (LDL). However, oral administration of methanol mango seed kernel extract significantly ($P < 0.05$) reduced the aforementioned indices to the levels which though were significantly ($P < 0.05$) higher than that reported for the normal control group. On the other hand, it was observed that the levels of high density lipoprotein (HDL) and Triacylglyceride (TG) in the negative control (Group II) were significantly low but increased following oral administration of extract in a dose dependent manner. This may be attributed to enhanced inhibition of intestinal absorption of cholesterol, interference with lipoprotein production, increased expression of hepatic LDL receptors and their protection which culminate to leading to increased elimination of LDL-C from the blood and its increased degradation of cholesterol in the body all of which translate to declined serum LDL-C levels which may have also reduced serum cholesterol (TC) levels (Brown and Goldstein, 1981). This is consistent with the findings of Khyati et al (2010) which showed that aqueous leaf extract of *Mangifera indica* significantly ($P < 0.05$) reduced TC, TG, LDL-C, VLDL and significant increase in HDL-C ($p < 0.05$). Table 2 shows the effect of treatment with MLEMI on the weight of hyperlipidemic rats indicating that treatment with extract reduced the weight of hyperlipidemic rats.

Formatted: Font: 12 pt

Comment [PM13]: necessary to discuss

Formatted: Font: 12 pt

Conclusion

References

Comment [PM14]: add references and recent references

- [1] Ghule BV, Ghante MH, Saoji AN, Yeole PG, Hypolipidemic and antihyperlipidemic effects of *Lagenaria siceraria* (Mol.) fruit extracts, *Indian J Exp Biol* 44, 905-909, 2006.
- [2] Ansarullah, Jadeja RN, Thounaojam MC, Patel V, Devkar RV, Ramachandran AV, Antihyperlipidemic potential of a polyherbal preparation on triton WR 1339 (Tyloxapol) induced hyperlipidemia: A comparison with lovastatin, *Int J Green Pharm* 3, 119-124, 2009.
- [3] Shaik A, Elumalai A, Eswaraiah MC. An updated review on hepatoprotective medicinal plants. *Journal of Drug Delivery & Therapeutics*. 2012; 2(2):1-3.
- [4] Shah KA, Patel MB, Shah SS, Chauhan KN, Parmar PK, Patel NM. Antihyperlipidemic activity of *Mangifera indica* leaf extract on rats fed with high cholesterol diet. *Der Pharmacia Sinica* 2010; 1(2):156-161.
- [5] Muruganandan S, Scrivivasan K, Gupta S, Gupta PK, Lal J. Effect of mangiferin on hyperglycemia and atherogenicity in streptozotocin diabetic rats. *J Ethnopharmacol* 2005; 97:497-501.
- [6] Brown, M.S and Goldstein, J.L (1981) *N Engl J Med* 305: 515.
- [7] Khyati A. Shah, Mandev B. Patel, Shreya S. Shah, Kajal N. Chauhan, Parul K. Parmar, Natavarlal M. Patel. Antihyperlipidemic activity of *Mangifera indica* l. leaf extract on rats fed with high cholesterol diet *Der Pharmacia Sinica*, 2010, 1 (2): 156-161