

## Original Research Article

# Screening For Potential Lignocellulolytic Enzymes-Producing White Rot Fungi

### ABSTRACT

**Introduction:** The three major parts of lignocellulose are cellulose, hemicellulose and lignin which can be broken down by cellulase, xylanase and laccase, respectively, thereby making the reducing sugar in lignocellulose available for industrial processes.

**Aims:** This work aimed to screen for white-rot fungi with the potential of producing cellulase, xylanase and laccase which are vital in breaking lignocellulosic substrates.

**Methodology:** Some white rot fungi were screened for their abilities to produce cellulase, xylanase and laccase on potato dextrose agar supplemented separately with 1 % carboxyl methyl cellulose (CMC), 1 % of xylan, and 0.1 % of 2,2'-azino-bis (3-ethylbenzthiazoline-6-sulphonic acid), respectively and enzymes relative activity were determined.

**Results:** The highest average relative activity of cellulase ( $1.07 \pm 0.073$ ) was recorded with *Lentinus squarrosulus* while the same average relative xylanase activity (1.13) was produced by both *Lentinus squarrosulus* and *Pleurotus ostreatus*. *Pleurotus tuber-regium* did not produce cellulase and xylanase. The highest average relative laccase activity ( $1.43 \pm 0.286$ ) was produced by *Pleurotus tuber-regium* followed by *Pleurotus ostreatus* ( $1.24 \pm 0.162$ ) and the least was by *Lentinus squarrosulus* ( $1.12 \pm 0.134$ ).

**Conclusion:** *Pleurotus ostreatus* and *Lentinus squarrosulus* produced cellulase, xylanase and laccase which are important in breaking down lignocellulose. *Pleurotus ostreatus* and *Lentinus squarrosulus* could be employed to break down lignocellulose.

**Keywords:** *Pleurotus ostreatus*, *Lentinus squarrosulus*, *Pleurotus tuber-regium*, white rot fungi, cellulase, xylanase, laccase, lignin modifying enzymes

### 1. INTRODUCTION

Lignocelluloses are wastes in the environment which consist of three main components such as cellulose, hemicellulose and lignin [1,2]. Cellulose is made up of hexoses while hemicellulose contains hexoses and pentoses. Reducing sugars in lignocellulolytic substrates that could be converted to value-added products are not readily available for bioconversion until lignocelluloses are pretreated. There are three main types of pretreatment: chemical, physical and biological [1]. Some of the advantages of the biological pretreatment method are low energy consumption, low cost and environmentally friendly procedures [1,3]. Biological pretreatment is the use of organisms or their metabolites to break down lignocellulose. Organisms suitable for breaking down lignocellulose to simple sugar should be able to produce lignocellulolytic enzymes needed to break down the three main components of lignocellulose.

Cellulases are enzymes that break down the cellulolytic part of lignocellulosic substrates to simple sugars while xylanases are involved in the conversion of the hemicellulose portion of lignocellulose to sugar [4,5]. Lignases are used to degrade lignin that binds lignocellulose together. Production of three main lignocellulolytic enzymes by a single organism indicates

Comment [srd1]: Check the sentence

that the organisms could be used to break down lignocellulose yielding simple sugar that could be utilized for other value-added products.

There are different groups of fungi involved in the degradation of lignocellulosic substrates which are white-rot fungi, brown-rot fungi and soft-rot fungi [6]. White rot and brown rot fungi are Basidiomycetes while soft rot fungi are Ascomycetes [6,7]. White rot fungi is a group of fungi that breaks down lignocellulosic substrates leaving a white fibrous appearance on degraded substrates. This group of fungi produce lignocellulolytic enzymes (cellulase, xylanase and ligninolytic enzymes) which are important to many industries. Ligninolytic enzymes are also referred to as lignin-modifying enzymes, lignases or ligninases. Laccase, manganese peroxidase, lignin peroxidase and versatile peroxidase are examples of lignin-modifying enzymes. White rot fungi have been used for the degradation of lignocellulose by some researchers [1,2,8]. This work aimed at screening some white rot fungi for their ability to produce lignocellulolytic enzymes (cellulase, xylanase and lignin-modifying enzymes) on plate.

## **2. MATERIAL AND METHODS**

### **2.1 Collection of White Rot Fungi**

Three white rot fungi (*Pleurotus ostreatus*, *Pleurotus tuber-regium* and *Lentinus squarrosulus*) were collected from the Department of Botany, University of Ibadan, Ibadan. They were screened for their ability to produce cellulase, hemicellulase and lignase/laccase.

### **2.2 Screening of White Rot Fungi for Enzymes Production**

#### **2.2.1 Screening of White Rot Fungi for Cellulase Production**

Potato Dextrose Agar was prepared and supplemented with 1 % of Carboxyl Methyl Cellulose (CMC). It was sterilized at 121 °C and 1.05kg cm<sup>-2</sup> for 15 minutes. It was dispensed into sterile Petri dishes and allowed to solidify. Each plate was inoculated with *Pleurotus ostreatus*, *Pleurotus tuber-regium* and *Lentinus squarrosulus* separately and incubated at 28 ±2 °C for five days. There were five sets for each white rot fungus. Each set was taken every 24 hours, flooded with 2% (w/v) aqueous congo red and left for 15 minutes. Excess stain was poured off after 15 minutes and washed with distilled water and the appearance of a clear area around colonies against a red colour for undegraded CMC indicates cellulase production [9]. Relative cellulase activities were determined by dividing the diameter of hydrolysed CMC by the diameter of the organism.

#### **2.2.2 Screening of White Rot Fungi for Xylanase Production**

Xylan is the main component of hemicellulose. One percent (1% w/v) xylan was used to supplement potato dextrose agar and was sterilized at 121 °C and 1.05kg cm<sup>-2</sup> for 15 minutes. It was dispensed into sterile Petri dishes and allowed to gel. *Pleurotus ostreatus*, *Pleurotus tuber-regium* and *Lentinus squarrosulus* were inoculated into each plate separately and incubated at 28 ±2 °C for five days. There were five sets for each organism. Each set was taken every 24 hours and flooded with iodine stain (0.25% w/v aqueous I<sub>2</sub> and KI) and left for 5 minutes. The stain was poured off the plate after 5 minutes and was washed with distilled water. The appearance of a clear zone against a blue/reddish purple colour shows xylanase activities [9]. Relative xylanase activities were determined by dividing the diameter of hydrolysed xylan by the diameter of the organism.

#### **2.2.3 Screening of White Rot Fungi for Lignase Production**

Tannic acid agar was prepared by supplementing potato dextrose agar with one percent tannic acid. It was sterilized at 121 °C and 1.05kg cm<sup>-2</sup> for 15 minutes. It was dispensed into sterile Petri dishes and allowed to solidify. Each plate was inoculated separately with *Pleurotus ostreatus*, *Pleurotus tuber-regium* and *Lentinus squarrosulus* and incubated at 28 ±2°C. Growth and colour were observed every 24 hours for five days. A brown oxidation zone around colonies indicates lignin degradation [9]. Relative lignase activities were determined by dividing the diameter of degraded lignin by the diameter of the organism.

#### **2.2.4 Screening of White Rot Fungi for Laccase Production**

Laccase is one of the enzymes involved in lignin degradation. Potato dextrose agar was supplemented with 0.1% (w/v) of 2,2'-azino-bis (3-ethylbenzthiazoline-6-sulphonic acid) (ABTS). It was sterilized at 121 °C and 1.05kg cm<sup>-2</sup> for 15 minutes, dispensed into sterile Petri dish, allowed to solidify and inoculated separately with *Pleurotostreatus*, *Pleurotus tuber-regium* and *Lentinus squarrosulus* and incubated at 28 ±2°C. The plates were observed every 24 hours for five days for the development of green or purple colouration around the colonies indicating the production of laccase [9]. Relative laccase activities were determined by dividing the diameter of oxidized ABTS by the diameter of the organism.


### 3. RESULTS

The three white rot fungi (*Lentinus squarrosulus*, *Pleurotostreatus* and *Pleurotus tuber-regium*) screened for cellulase, xylanase, lignase and laccase showed the ability to produce two or all four enzymes. The appearance of a clear area around fungus growth against a red colour of undegraded carboxymethyl cellulose (CMC) as shown by *Lentinus squarrosulus* indicated the ability of the organism to produce cellulase on the plate (Plate 1). A xylanase-producing *Lentinus squarrosulus* on potato dextrose agar supplemented with xylan is shown in Plate 2. The appearance of a clear area around the organism against a blue/reddish purple colour showed xylanase activities.


Plate 3 shows *Pleurotostreatus* with the ability to degrade lignin on tannic agar. The brown colouration around the growing fungus confirmed its ability to produce lignase. Lignase combines all the enzymes involved in lignin degradation which are known as lignin-modifying enzymes. Laccase-producing *Pleurotus tuber-regium* on potato dextrose agar supplemented with 2, 2'-azino-bis (3-ethylbenzthiazoline-6-sulphonic acid) (ABTS) is shown in Plate 4. The development of purple colouration indicated the ability of the fungus to produce laccase.

The three white rot fungi (*Lentinus squarrosulus*, *Pleurotostreatus* and *Pleurotus tuber-regium*) were able to produce cellulase, xylanase, lignase and laccase except for *Pleurotus tuber-regium* that could not produce cellulase and xylanase as shown in Table 1. The relative lignase activity of *Lentinus squarrosulus* ranged from 1.15 (96 hours) to 1.43 (24 hours). An increase in relative lignase activity of *Pleurotostreatus* (1.10 – 1.72) and *Pleurotus tuber-regium* (1.10 – 2.40) was observed with increase in the period of incubation with the least and highest at 24 and 120 hours of incubation, respectively. Relative laccase activity of *Lentinus squarrosulus* and *Pleurotostreatus* decreased with an increase in days of incubation from 1.36 to 1.03 and 1.40 to 1.06, respectively whereas that of *Pleurotus tuber-regium* increased in the first 72 hours of incubation from 1.10 to 1.82 and thereafter decreased. A decrease in relative cellulase activity of both *Lentinus squarrosulus* and *Pleurotostreatus* with an increase in the day of incubation was recorded.


Relative xylanase activity of *Lentinus squarrosulus* ranged from 1.01 to 1.40 while that of *Pleurotostreatus* ranged from 1.02 to 1.40. *Pleurotus tuber-regium* did not produce both cellulase and xylanase on the plate. Average relative activities of lignase and laccase by *Pleurotus tuber-regium* (1.80 ± 0.537 and 1.43 ± 0.286) > *Pleurotostreatus* (1.49 ± 0.260 and 1.24 ± 0.162) > *Lentinus squarrosulus* (1.27 ± 0.103 and 1.12 ± 0.134), respectively. The highest average relative cellulase activity (1.07 ± 0.073) was obtained by *Lentinus squarrosulus* and the same average relative xylanase activity (1.13) was recorded by *Lentinus squarrosulus* and *Pleurotostreatus*.




**Plate 1: Cellulase-Producing *Lentinus squarrosulus* on Carboxymethyl Cellulose Agar**



**Plate 2: Xylanase-Producing *Lentinus squarrosulus* on Potato Dextrose Agar Supplemented with Xylan**



**Plate 3: Lignase-Producing *Pleurotus ostreatus* on Tannic Acid Agar**



**Plate 4: Lacase-Producing *Pleurotus tuber-regium* on Potato Dextrose Agar Supplemented with 2,2'-azino-bis (3-ethylbenzthiazoline-6-sulphonic acid) (ABTS)**

**Table 1: Relative Activities of Lignocellulolytic Enzymes of Some White Rot Fungi on Plate**

White Rot Fungi	Period	Lignocellulolytic Enzymes			
		Lignase	Laccase	Cellulase	Xylanase
<i>Lentinus squarrosulus</i>	24	1.43	1.36	1.19	1.40
	48	1.27	1.08	1.08	1.14
	72	1.27	1.10	1.03	1.06
	96	1.15	1.05	1.03	1.06
	120	1.22	1.03	1.01	1.01
	Average	1.27 0.103	± 1.12 0.134	± 1.07 0.073	± 1.13 0.156
<i>Pleurotus ostreatus</i>	24	1.10	1.40	1.09	1.40
	48	1.35	1.38	1.07	1.14
	72	1.61	1.28	1.03	1.06
	96	1.67	1.08	1.02	1.02
	120	1.72	1.06	1.02	1.02
	Average	1.49 0.260	± 1.24 0.162	± 1.05 0.032	± 1.13 0.159
<i>Pleurotus tuber-regium</i>	24	1.10	1.10	-	-
	48	1.40	1.20	-	-
	72	1.95	1.82	-	-
	96	2.15	1.50	-	-
	120	2.40	1.52	-	-
	Average	1.80 0.537	± 1.43 0.286	± -	-

Key: - : Enzyme not detected

#### 4. DISCUSSION

The white rot fungi used in this work had the ability to produce hydrolytic (cellulase and xylanase) and oxidative (laccase) enzymes. Production of hydrolytic enzymes by *Pleurotostreatus* and *Lentinus squarrosulus* was confirmed by their cellulose-degradation potential in the formation of clear halos around the fungi plugs against a pink Congo red-cellulose complex. An indication of their abilities to break down cellulose to simple sugars as reported by Olanbiwoninu and Fasiku, [10]. Production of cellulase by *Pleurotostreatus* and *Lentinus squarrosulus* as observed in this study has been reported by some researchers to be due to the ability of the organisms to excrete hydrolysing enzymes which effectively broke down the cellulolytic component of lignocellulose [11-19]. However, the inability of *Pleurotus tuber-regium* to produce cellulase in the course of this work could be an influence of environmental factors or the genetic makeup of the organism which might make the degradation of cellulose difficult.

In the hydrolysis of xylan, the clear zone against the blue-black colour observed is an indication of the fungal abilities to produce hemicellulase [5,20,21]. The fungi effectively converted xylan to hexoses and pentoses which resulted in clear zones around xylanase producing mushrooms on xylan agar. Xylanase production by both *Pleurotostreatus* and *Lentinus squarrosulus* could have been determined by the genetic makeup of the organisms because many researchers have reported using these organisms to produce xylanase. Akhmedova and Rashidova [18], Majumder [22] and, Han *et al.* [23] produced xylanase with *Pleurotostreatus* while Vichitrakaet *al.* [19], Pukahutaet *al.* [24] and, Isikhuemhenet *al.* [25] used *Lentinus squarrosulus* to produce xylanase.

White-rot fungi are efficient in degrading lignin compounds because of their ability to produce lignin-modifying enzymes such as laccase and others. Production of lignase on tannic acid agar by *Pleurotostreatus*, *Pleurotus tuber-regium* and *Lentinus squarrosulus* in this work might be due to their abilities to utilise tannic acid which resulted in the brown coloration observed on the plates as earlier reported by Pointing [9]. Gramsset *al.* [26] explained that laccase-producing organisms oxidized 2, 2'-Azino-bis (3-ethylbenzthiazoline-6-sulphonic acid) (ABTS) that was colourless to ABTS<sup>2+</sup> (purple) and purple coloration that appeared on medium supplemented with ABTS in this work indicated abilities of mushroom to produce laccase. The oxidative capability of lignin-modifying enzymes of *Pleurotostreatus* and *Lentinus squarrosulus* recorded in this research has been confirmed by many researchers [27-30].

#### 5. CONCLUSION

*Pleurotostreatus* and *Lentinus squarrosulus* produced the three main lignocellulolytic enzymes (cellulase, xylanase and lignin-modifying enzymes) while *Pleurotus tuber-regium* produced cellulase and lignase. *Pleurotostreatus* and *Lentinus squarrosulus* with the potential of producing lignocellulolytic enzymes could be utilized to degrade lignocellulosic substrates releasing reducing sugars that are important for bioconversion.

#### REFERENCES

1. Fasiku SA, Wakil SM. Pretreatment of maize straw with *Pleurotostreatus* and *Lentinus squarrosulus* for bioethanol production using *Saccharomyces cerevisiae*. Novel Research in Microbiology Journal. 2021;5(6):1480-1493.

- 2 Fasiku SA, Wakil SM. Screening of factors responsible for conversion of maize straw into bioethanol. *Journal of Microbiology, Biotechnology and Food Sciences*. 2022;12(2):e5901.
- 3 Wagner AO, Lackner N, Mutschiechner M, Prem EM, Markt R, Illmer P. Biological Pretreatment Strategies for Second-Generation Lignocellulosic Resources to Enhance Biogas Production. *Energies (Basel)*. 2018; <https://doi.org/10.3390%2Fen11071797>
- 4 Fasiku SA, Ogunsola OF, Fakunle A, Olanbiwoninu AA. Isolation of Bacteria with Potential of Producing Extracellular Enzymes (Amylase, Cellulase and Protease) from Soil Samples. *Journal of Advances in Microbiology*. 2020;20(3):21-26.
- 5 Fasiku SA, Bello MA, Odeniyi OA. Production of xylanase by *Aspergillus niger* GIO and *Bacillus megaterium* through solid-state fermentation. *Access Microbiology*. 2023; <https://doi.org/10.1099/acmi.0.000506.v3>
- 6 Isroi, Millati R, Syamsiah S, Niklasson C, Cahyanto MN, Lundquist K, Taherzadeh MJ.. Biological Pretreatment of Lignocelluloses with White-Rot Fungi and its Application: A Review. *Bioresources*. 2011;6(4):5224-5259.
- 7 Hatakka A. Biodegradation of lignin. *Biology, Chemistry, Biotechnology, Application*. Vol. 1. Lignin, Humic Substances and Coal, M. Hofrichter and A. Steinbuchel (eds.), Wiley-WCH, 2001;129-180.
- 8 Ibarra-Islas A, Lopez PMG, Hernandez JEM, Leon SH, Armenta S, Arce-Cervantes O, Lopez JE. Use of Nutchells wastes in the production of Lignocellulolytic Enzymes by White rot fungi. *Agriculture, Agribusiness and Biotechnology*. 2023;66:e23210654
- 9 Pointing SB. Qualitative methods for the determination of lignocellulolytic enzyme production by tropical fungi. *Fungal Diversity*. 1999;2:17 – 33.
- 10 Olanbiwoninu AA, Fasiku SA. Production of bacterial amylases and cellulases using sweet potato (*Ipomoea batatas*. (L.) Lam.) peels. *African Journal of Biochemistry Research*. 2015;9(9):104-109.
- 11 Atri NS and Sharma SK. Qualitative Estimation of Cellulases and Lignin Modifying Enzymes in Five Wild *Lentinus* Species Selected from North West India. *Academic Journal of Plant Sciences* 2011;4(4):105-109.
- 12 Khalil MI, Hoque MM, Basunia MA, Alam N. Production of cellulase by *Pleurotostreatus* and *Pleurotussajor-caju* in solid state fermentation of lignocellulosic biomass. *Turkish Journal of Agriculture Forestry*. 2011;35:333-341.
- 13 Karthikeyan P. Optimization of cellulase enzyme production from *Pleurotostreatus* and *Calocybe indica*. *Research Article Biological Sciences*. 2015;5(1):11-16.
- 14 Debnath G, Das P, Saha AJ. 'Extracellular Cellulase Produced By *Lentinus Squarrosulus* A Wild Edible Mushroom of Tripura'. *International Journal of Current Advanced Research*. 2018;07(1), pp. 8877-8880.
- 15 Premkumar G, Muthuramkumar S, Kannan M, Varatharaju G, Rajarathinam K. Cellulase and Xylanase Production by *Pleurotus* sp. on Mixed Substrate System. *Journal of Biotechnology and Biomaterials*. 2018;8(2):1-4.
- 16 Huang L, Sun N, Ban L, Wang Y, Yang H. Ability of different edible fungi to degrade crop straw. *AMB Express*. 2019;9(1):4.
- 17 Datsomor O, Yan Q, Opoku-Mensah L, Zhao G, Miao L. Effect of Different Inducer Sources on Cellulase Enzyme Production by White-Rot Basidiomycetes *Pleurotostreatus* and *Phanerochaetechrysosporium* under Submerged Fermentation. *Fermentation*. 2022;8:561. <https://doi.org/10.3390/fermentation8100561>
- 18 Akhmedova ZR, Rashidova NT. Enzymatic Activity of wood degrading-Basidiomycetes *Pleurotostreatus* UZBI-105 in the various carbon sources. *Spectrum Journal of Innovation, Reforms and Development*. 2022;6:87-91.
- 19 Vichitraka A, Somboon P, Tantratian S, Onmankhong J, Sirisomboon P, Pornchaloempong P, Pukahuta C, Pornpukdeewattana S, Krusong W, Charoenrat T. Application of baby corn husk as a biological sustainable feedstock for the production of

- cellulase and xylanase by *Lentinus squarrosulus* Mont., Bioresource Technology Reports. 2023;21:101341, <https://doi.org/10.1016/j.biteb.2023.101341>.
- 20 Yu X, Atalla RH. The complex of xylan and iodine: the induction and detection of nanoscale order. Carbohydrate Research. 2005;340:981-988.
- 21 Mohammad NS, Ariffin ZZ. *Aspegillus sydowii* strain SCAU066 and *Aspergillus versicolor* Isolate BAB-6580: Potential source of xylanolytic, cellulolytic and Amyololytic Enzymes. Science Letters. 2020;14(2):15-23.
- 22 Majumder K. Production of xylanase and CMCase by *Pleurotostreatus* in Polyurethane foam based solid state fermentation. International Journal of Advanced Scientific Research and Management. 2019;4(4):7-11.
- 23 Han M, An Q, He S, Zhang X, Zhang M, Gao X, Wu Q, Bian L. Solid-state fermentation on poplar sawdust and corncob wastes for lignocellulolytic enzymes by different *Pleurotostreatus* strains. BioResources. 2020;15(3):4982-4995.
- 24 Pukahuta C, Suwanarit P, Shianagawa E, Hoshida H, Nishizawa. Combination of Laccase, Xylanase and Cellulase in Lignocellulose Degradation by White Rot Fungi, *Lentinus polychrous* Lev. and *L. squarrosulus* Mont. Kasetsart Journal – Natural Science. 2004;38:65-73.
- 25 Isikhuemhen OS, Mikiashvili NA, Adenipekun CO, Ohimain EI, Shahbazi G. The tropical white rot fungus, *Lentinus squarrosulus* Mont.: lignocellulolytic enzymes activities and sugar release from cornstalks under solid state fermentation. World Journal of Microbiology and Biotechnology 2012;28:1961-1966.
- 26 Gramss G. Reappraising a Controversy: Formation and Role of the Azodication (ABTS<sup>2+</sup>) in the Laccase-ABTS Catalyzed Breakdown of Lignin. Fermentation. 2017;3:27.
- 27 Alvarez-Cervantes J, Sanchez C, Diaz R, Diaz-Godinez G. Characterization of production of laccases, cellulases, and xylanases of *Pleurotostreatus* grown on solid-state fermentation using an inert support. Revista Mexicana de IngenieriaQuimica. 2016;15(2):323-331.
- 28 Kumar VV, Venkataraman S, Kumar PS, George J, Rajendran DS, Shaji A, Lawrence N, Saikia K, Rathankumar AK. Laccase production by *Pleurotostreatus* using cassava waste and its application in remediation of phenolic and polycyclic aromatic hydrocarbon-contaminated lignocellulosic biorefinery wastewater. Environmental Pollution. 2022;309:119729.
- 29 Franco PCI, Shiraishi IS, Dekker RFH, Barbosa-Dekker AM, Borsato D, Angelelli KB, Evaristo GPC, Simionato JI, Daniel JS. Optimization of laccase production by *Pleurotostreatus* Florida and evaluation of metabolites generated during Kraft lignin biotransformation. Waste Biomass Valor. 2023; <https://doi.org/10.1007/s12649-022-02029-9>
- 30 Mathur A, Dubey S, Prasad R, Singh RP. Mycelial and secretome proteomic dynamics of *L. squarrosulus* AF5 in azo dye degradation, Journal of Environmental Chemical Engineering. 2023;11(2):109374, <https://doi.org/10.1016/j.jece.2023.109374>.