

**Effect of Phosphorus and Molybdenum on growth and yield of cowpea (*Vigna unguiculata* L.)**

**ABSTRACT**

A field experiment was conducted during *Zaid* season of 2022 at Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology And Sciences, Prayagraj, Uttar Pradesh, India to determine the “Effect of phosphorus and molybdenum on the growth and yield of summer Cowpea (*Vigna unguiculata* L.)”. The results showed that the treatment 9 [Phosphorus (50kg/ha) + Molybdenum (2 kg/ha)] recorded significantly higher plant height (73.43 cm), higher number of nodules per plant (17.66), higher plant dry weight (23.07 g/plant), higher crop growth rate (18.87g/m<sup>2</sup>/day) maximum number of pods per plant (15.10), maximum number of seeds per pod (14.67), higher seed index (16.03 g), higher seed yield (1550.27 kg/ha) and higher stover yield (1615.50 kg/ha). Similarly, maximum gross return (94954.8 INR/ha), higher net return (65670.3 INR/ha) and highest Benefit Cost ratio (2.24) was also recorded in treatment 9 [Phosphorus (50kg/ha) + Molybdenum (2kg/ha)] as compared to other treatments.

**Key words:** Phosphorus, Molybdenum, Growth, Yield, Economics.

## Introduction

Cowpea is commonly grown in sub-tropical regions that are moderately humid and warm and more drought resilient however, it is not tolerant to frost and waterlogging. Seeds of cowpea are nutritious and cheap sources of quality protein, vitamins, iron and phosphorus as an excellent substitute for eggs, meat and other protein-rich foods thus, they are a significant part of the human diet. It also plays an important role in providing soil nitrogen to cereal crops (such as maize, millet and sorghum) when grown in rotation, especially in areas where poor soil fertility is a problem. It does not require a high rate of nitrogen fertilization; its roots have nodules in which soil bacteria called Rhizobia inhabit and help to fix nitrogen from the air into the soil in the form of nitrates (**Sheahan, 2012**). The protein in cowpea seeds is rich in amino acids like lysine, phenyl amine, leucine and tryptophan compared to cereal grains (**Gad and Kandil, 2013**). The protein content present in cowpea is about 25-30% in mature seeds, 4-5% in immature pods and 3-5% in green leaves (**Bressani and Elias, 1980**). The ripe seed on average contains 22% protein, 1.4% fat, 59.1% carbohydrate, and 3.7% ash. The energy value is 340 kcal/ 100 g.

In India, pulses are grown nearly in 25.43 m ha with an annual production of 17.28 m t and productivity of 679 kg/ha. The per capita availability of pulses in India is 35.5g/day as against the minimum requirement of 70g/day/capita as advocated by the Indian Council of Medical Research. Cowpea is grown across the world on an area of 14.5 m ha of land planted each year and the total annual production is 6.2m t. In India during 2020-21 cowpea grew about 13.3 m ha with an annual production of 8.06 m t and productivity of 596 kg/ha. Some of the states like Uttar Pradesh are about 2.38 m ha with an annual production of 2.56 and productivity of 1079 kg/ha major producer of cowpea in India as advocated by the Ministry of Agriculture and Farmers Welfare (GOI, 2020-21).

Phosphorus is critical to cowpea yield because it is reported to stimulate growth, initiate nodule formation as well as influence the efficiency of rhizobium-legume symbiosis (**Haruna and Aliyu, 2011**). It is required in large quantities in young cells such as shoot and root tips to increase metabolism and promote rapid cell division. It also aids in flower initiation, seed and fruit development (**Ndakidemi and Dakora, 2007**). Phosphorus plays key roles in many plant processes such as energy metabolism, nitrogen fixation, synthesis of nucleic acids and membranes, photosynthesis, respiration and enzyme regulation. (**Karikari et al., 2015**).

Molybdenum also playing important role in the structural interring of the cell wall and cell

membrane and synthesis of protein as well as nitrogen fixation. Molybdenum is required for the formation of the nitrate reductase enzyme and in the legume, it plays an additional role in symbiotic nitrogen fixation. The nitrogen-fixing enzyme, nitrogenase is a compound of molybdenum without adequate quantities of this element, nitrogen fixation can't occur. Thus, the application of Mo increased the yield but also improved the quality (Movalia *et al.*, 2020). Molybdenum also has a function of a co-factor, i.e., structural and catalytic function, in several enzymes which have been discovered in higher plants. Although molybdenum does not take part in nitrogen metabolism directly, some researchers showed that it still plays an important stimulation role in the process of nitrogen metabolism. Keeping in view the above facts, the present investigation was undertaken to find out "Effect of phosphorus and molybdenum on the growth and yield of summer Cowpea (*Vigna unguiculata* L.)"

#### **MATERIALS AND METHODS:**

The experiment was conducted during *zaid* season of 2022 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj (U.P). The soil of the field constituting a part of central Gangetic alluvium is neutral and deep. The soil of the experimental field was sandy loam in texture, nearly neutral in soil reaction (pH 7.8), low level of organic carbon (0.62%), available N (225 kg/ha), P (38.2 kg/ha), K (240.7 kg/ha) and zinc (2.32 mg/kg). The treatment consisting of three levels of phosphorus *viz.*, 30, 40, 50 kg/ha and three levels of molybdenum *viz.*, 1, 1.5, 2 kg/ha. The experiment was laid out in RBD with 10 treatments each replicated thrice. The treatment combinations are T1 - phosphorus (30kg/ha) + molybdenum (1 kg/ha), T2 - phosphorus (30kg/ha) + molybdenum (1.5 kg/ha), T3 - phosphorus (30kg/ha) + molybdenum (2 kg/ha), T4 - phosphorus (40kg/ha) + molybdenum (1 kg/ha), T5 - phosphorus (40kg/ha) + molybdenum (1.5 kg/ha), T6 - phosphorus (40kg/ha) + molybdenum (2 kg/ha), T7 - phosphorus (50kg/ha) + molybdenum (1 kg/ha), T8 - phosphorus (50kg/ha) + molybdenum (1.5 kg/ha), T9 - phosphorus (50kg/ha) + molybdenum (2 kg/ha), T10 – Control (25:50:25NPK kg/ha). The growth parameters and yield, production was recorded at harvest from randomly selected plants in each plot. The data was computed and analyzed by following statistical method of Gomez and Gomez (1984).

## RESULTS AND DISCUSSION

### Growth parameters

#### Plant height (cm)

At 60DAS, higher plant height (73.43 cm) was recorded in treatment 9 [Phosphorus (50kg/ha) + Molybdenum (2 kg/ha)] which was significantly superior over all other treatments. However, treatment 6 [Phosphorus (40kg/ha) + Molybdenum (2 kg/ha)] is statistically at par with treatment 9 [Phosphorus (50kg/ha) + Molybdenum (2kg /ha)]. The significant and highest plant height observed with the application of phosphorus (50 kg/ha) might be due the role of phosphorus in cell division (**Namakka et al., 2017**). Further, higher plant height was found to the fact that molybdenum (2kg/ha) increases availability of nitrogen which helps in the process of nitrogen assimilation. Similar findings also reported by **Chatterjee and Bandyopadhyay (2015)**.

#### Number of Nodules/plant

At 60DAS, higher number of Nodules /Plant (7.89) was recorded in treatment 9 [Phosphorus (50kg/ha) + Molybdenum (2 kg/ha)], which was significantly superior over all other treatments. However, treatment 6 [Phosphorus (40kg/ha) + Molybdenum (2 kg/ha)] is statistically at par with treatment 9 [Phosphorus (50kg /ha) + Molybdenum (2kg /ha)]. Application of phosphorus showed significant and higher number of nodules/plant might due to it plays role in energy metabolism, nitrogen fixation, synthesis of nucleic acid and membranes, photosynthesis, respiration and enzyme regulation (**Nkaa et al.,2014**). Further, Mo (20 kg/ha) application significantly increased the number of nodules/plant due to its vital role in synthesis and activity of molybdo enzymes such as nitrogen assimilation enzyme – nitrate reductase and the nitrogen fixing enzyme - nitrogenase, key regulatory component for initiation of nodulation. These results were in conformity with those of **Franco and Munns (1981)**.

#### Plant Dry Weight (g)

At 60DAS, higher plant dry weight (23.07 g) was recorded in treatment 9 [Phosphorus (50kg/ha) + Molybdenum (2 kg/ha)], which was significantly superior over all other treatments. However treatment 6 [Phosphorus (40kg/ha) + Molybdenum (2 kg/ha)] is statistically at par

with treatment 9 [Phosphorus (50kg /ha) + Molybdenum (2kg /ha)]. Maximum plant dry weight was increased significantly with application of molybdenum (2kg/ha) this might be due to higher availability of nutrients, synthesis of more carbohydrates and their translocation to different plant parts resulted in increased vegetative growth including the reproductive structures which in turn increase dry weight of the plant. Similar results have been reported by **Harireddy et al (2021)**.

#### **Crop growth rate (g/m<sup>2</sup>/day)**

At 45-60DAS, higher crop growth rate (11.78 g /m<sup>2</sup>/day) was recorded in treatment 9 [Phosphorus (50kg/ha) + Molybdenum (2 kg/ha)], which was significantly superior over all other treatments. However, treatment 1 [Phosphorus (30kg/ha) + Molybdenum (1 kg/ha)], treatment 2 [Phosphorus (30kg/ha) + Molybdenum (1.5 kg/ha)], treatment 3 [Phosphorus (30kg/ha) + Molybdenum (2 kg/ha)], treatment 4 [Phosphorus (40kg/ha) + Molybdenum (1 kg/ha)], treatment 5 [Phosphorus (40kg/ha) + Molybdenum (1.5 kg/ha)], treatment 6 [Phosphorus (40kg/ha) + Molybdenum (2 kg/ha)] were statistically at par with treatment 9 [Phosphorus (50kg /ha) + Molybdenum (2kg /ha)]. The maximum crop growth rate was observed by the application of molybdenum (2 kg/ha) this might be due to better accumulation of dry matter throughout the plant vegetative and reproductive phase, which enhances the physiological and metabolic activity and growth by assimilating the available nutrients at higher rate, facilitating more photosynthesis. A similar trend was reported by **Harshini et al (2022)**.

#### **Yield parameters**

##### **Number of pods/plants**

Treatment with application of treatment 9 [Phosphorus (50kg/ha) + Molybdenum (2 kg/ha)] was recorded higher number of pods/plant (15.10) which was significantly superior over all other treatments. However, treatment 6 [Phosphorus (40kg/ha) + Molybdenum (2 kg/ha)] is statistically at par with treatment 9 [Phosphorus (50kg/ha) + Molybdenum (2 kg/ha)]. The higher number of pods/plant with application of phosphorus (50 kg/ha) might be due to it favourably influenced the plant vigour, morphology and metabolic processes. (**Chatterjee and Bandyopadhyay. 2015**). Further higher number of pods/plant was observed with application of molybdenum (2kg/ha) might have been possible due to increased availability of nitrogen leads to biological nitrogen fixation that induces plant growth to produce huge biomass and number of pods/plant. Similar results were reported by **Mahesh et al., (2021)** in black gram.

##### **Number of seeds/pods**

Treatment with application of treatment 9 [Phosphorus (50kg/ha) + Molybdenum (2 kg/ha)] was recorded higher number of seeds/pods (14.67) which was significantly superior over all other

treatments. However, treatment 6 [Phosphorus (40kg/ha) + Molybdenum (2 kg/ha)] is statistically at par with treatment 9 [Phosphorus (50kg/ha) + Molybdenum (2 kg/ha)]. Application of phosphorus (50kg/ha) significantly increased number of seeds/pod might be due to its role in translocation of assimilates to the pods being a constituent of protoplasm. Similar results were reported by **Shilpa et al., (2011)**.

#### **Seed index.**

Treatment with application of treatment 9 [Phosphorus (50kg/ha) + Molybdenum (2 kg/ha)] was recorded higher number of test weight (16.03 g) which was significantly superior over all other treatments. However, treatment 6 [Phosphorus (40kg/ha) + Molybdenum (2 kg/ha)] is statistically at par with treatment 9 [Phosphorus (50kg/ha) + Molybdenum (2 kg/ha)]. Significant and higher seed index was with application of phosphorous (50kg/ha) might be due to increase photosynthetic activity of leaves, translocation of photosynthates from source to sink. Similar result was observed by **Singh et al., (2018)** in green gram. Further increase in seed index with the application of molybdenum (2kg/ha) might be due to the increase in phosphorus contents of the seed (phospholipids and phosphorus - protein) and also its involvement in many physiological processes in plants. Similar results was reported by **Tahir et al., (2014)** in black gram.

#### **Seed yield (kg/ha)**

Treatment with application of treatment 9 [Phosphorus (50kg/ha) + Molybdenum (2 kg/ha)] was recorded higher seed yield (1550.27 kg/ha) which was significantly superior over all other treatments. However, treatment 6 [Phosphorus (40kg/ha) + Molybdenum (2 kg/ha)] is statistically at par with treatment 9 [Phosphorus (50kg/ha) + Molybdenum (2 kg/ha)]. The application of Phosphorus (50 kg/ha) significantly increases seed yield might be due to its role in enhancing photosynthesis apparatus. (**Namakka et al., 2017**). Further, increase in seed yield with application of molybdenum (2kg/ha) might be due to improved rhizobium activity, nitrogen fixation, vegetative growth and yield components. Similar results were recorded by **Chatterjee and Bandyopadhyay (2015)**.

#### **Straw yield (kg/ha)**

Treatment with application of treatment 9 [Phosphorus (50kg/ha) + Molybdenum (2 kg/ha)] was recorded higher stover yield (1615.50 kg/ha) which was significantly superior over all other treatments. However, treatment 6 [Phosphorus (40kg/ha) + Molybdenum (2 kg/ha)] is statistically at par with treatment 9 [Phosphorus (50kg/ha) + Molybdenum (2 kg/ha)]. The

significant and higher straw yield increased with the application of phosphorus (50kg/ha) might be due to it enhanced the survival and multiplication of microorganisms, improved nitrogen fixation, transport of sugars and better uptake, assimilation of available nutrients by the plants during the entire growth period. Similar results were recorded by **Anand *et al.*, (2022)** in black gram. Further, higher straw yield was recorded with application of molybdenum (2kg/ha) might be due to it increased quantities of micronutrients in cowpea grain and stover, which may have been caused by plant leaves utilizing available micronutrients immediately. Zn, Fe, and Mo concentrations in cowpea grain and stover were raised. Similar findings were recorded by **Dhaliwal *et al.*, (2022)**.

### **Harvest index (%)**

Treatment with application of treatment 8 [Phosphorus (50kg/ha) + Molybdenum (1.5 kg/ha)] was recorded higher straw yield (49.05 %) which was significantly superior over all other treatments. However, treatment 3 [Phosphorus (30kg/ha) + Molybdenum (2 kg/ha)], treatment 4 [Phosphorus (40kg/ha) + Molybdenum (1kg/ha)], treatment 5 [Phosphorus (40kg/ha) + Molybdenum (1.5 kg/ha)], treatment 6 [Phosphorus (40kg/ha) + Molybdenum (2 kg/ha)], treatment 7 [Phosphorus (50kg/ha) + Molybdenum (1kg/ha)], treatment 9 [Phosphorus (50kg/ha) + Molybdenum (2 kg/ha)] were statistically at par with treatment 8 [Phosphorus (50kg/ha) + Molybdenum (1.5 kg/ha)]. The significant and higher harvest index was observed with application of phosphorus (50 kg/ha) might be due to development of extensive root system to extract more water and nutrient from soil (**Pramanik and Singh 2003**). Further higher harvest index was with application of molybdenum (2kg/ha) might be due to improved cell activities, enhanced cell multiplication, enlargement and luxuriant growth. Similar result was reported by **Kokani *et al.*, (2015)** in black gram.

### **Economics:**

The result showed that maximum gross return (94954.8 INR/ha), higher net return (65670.3 INR/ha) and higher benefit cost ratio (2.24) was recorded in treatment 9 [Phosphorus (50kg/ha) + Molybdenum (2 kg/ha)] as compared to other treatments.

Higher net returns, gross returns and B:C ratio was recorded with the application of phosphorus (50kg/ha) might be due to greater increase in grain and straw yield resulted higher benefit cost ratio. These results are in conformity with those observed by **Khanna *et al.*, (2019)**.

**Table 1: Effect of Phosphorus and Molybdenum on growth attributes of cowpea.**

S. No	Treatments	Plant height (cm)	Number of nodules/Plant	Plant dry weight(g)	CGR (g/m <sup>2</sup> /day)
1.	Phosphorus 30kg/ha + Molybdenum 1kg/ha	64.54	14.11	18.02	15.94
2.	Phosphorus 30kg/ha + Molybdenum 1.5kg/ha	67.02	14.78	18.63	15.57
3.	Phosphorus 30kg/ha + Molybdenum 2kg/ha	70.89	15.56	20.40	17.79
4.	Phosphorus 40kg/ha + Molybdenum 1kg/ha	69.31	15.33	19.20	16.79
5.	Phosphorus 40kg/ha + Molybdenum 1.5kg/ha	71.86	16.45	20.45	17.03
6.	Phosphorus 40kg/ha + Molybdenum 2kg/ha	73.17	17.44	22.24	18.41
7.	Phosphorus 50kg/ha + Molybdenum 1kg/ha	70.95	16.11	20.25	16.53
8.	Phosphorus 50kg/ha + Molybdenum 1.5kg/ha	72.15	16.89	20.79	17.44
9.	Phosphorus 50kg/ha + Molybdenum 2kg/ha	73.43	17.66	23.07	18.87
10.	Control (25:50:25NPK kg/ha)	61.85	13.33	16.20	14.13
	<b>F-test</b>	S	S	S	S
	<b>Sem±</b>	0.40	0.24	0.28	0.48
	<b>CD at 5%</b>	1.19	0.72	0.85	1.45

**Table 2: Effect of phosphorus and molybdenum on yield attributes and yield of cowpea.**

S.No index	Treatment Combination	Number of Pods/plant	Number of Seeds/pod	Seed index	Seed yield	Stover yield	Harvest
1.	Phosphorus 30kg/ha + Molybdenum 1 kg/ha	11.17	11.43	11.50	883.13	1090.77	44.77
2.	Phosphorus 30kg/ha + Molybdenum 1.5kg/ha	11.57	11.97	12.41	954.67	1158.50	45.18
3.	Phosphorus 30kg/ha + Molybdenum 2 kg/ha	13.20	12.70	13.86	1197.60	1273.33	48.45
4.	Phosphorus 40kg/ha + Molybdenum 1 kg/ha	11.20	12.47	13.03	1070.33	1190.33	47.54
5.	Phosphorus 40kg/ha + Molybdenum 1.5kg/ha	14.17	12.83	14.03	1313.67	1497.07	46.74
6.	Phosphorus 40kg/ha + Molybdenum 2 kg/ha	14.47	13.73	15.00	1476.80	1572.57	48.00
7.	Phosphorus 50kg/ha + Molybdenum 1 kg/ha	13.93	13.00	13.30	1268.43	1394.97	47.60
8.	Phosphorus 50kg/ha + Molybdenum 1.5kg/ha	13.80	13.07	14.09	1430.90	1486.80	49.05
9.	Phosphorus 50kg/ha + Molybdenum 2 kg/ha	15.10	14.67	16.03	1550.27	1615.50	48.96
10.	Control (25:50:25NPK kg/ha)	10.27	11.67	10.66	756.63	964.73	43.94
	<b>F Test</b>	S	S	S	S	S	S
	<b>SEm±</b>	0.29	0.39	0.40	30.33	27.96	0.85
	<b>CD (P=0.05)</b>	0.88	1.18	1.21	90.12	83.09	2.54

**Table 3: Effect of phosphorus and molybdenum on economics of cowpea.**

S.No	Treatment Combination	Total cost of Cultivation (INR/ha)	Gross returns	Net returns	B: C ratio
1.	Phosphorus 30kg/ha + Molybdenum 1 kg/ha	24564.5	54296.52	29732.02	1.21
2.	Phosphorus 30kg/ha + Molybdenum 1.5 kg/ha	25668.5	58670.4	33001.9	1.28
3.	Phosphorus 30kg/ha + Molybdenum 2 kg/ha	26784.5	73383.8	46599.3	1.73
4.	Phosphorus 40kg/ha + Molybdenum 1 kg/ha	25814.5	65648.2	39833.7	1.54
5.	Phosphorus 40kg/ha + Molybdenum 1.5 kg/ha	26918.5	80616.68	53698.18	1.99
6.	Phosphorus 40kg/ha + Molybdenum 2 kg/ha	28034.5	90495.08	62460.58	2.22
7.	Phosphorus 50kg/ha + Molybdenum 1 kg/ha	27064.5	77779.56	50715.06	1.87
8.	Phosphorus 50kg/ha + Molybdenum 1.5 kg/ha	28168.5	87637.92	59469.42	2.11
9.	Phosphorus 50kg/ha + Molybdenum 2 kg/ha	29284.5	94954.8	65670.3	2.24
10.	Control (25:50:25 NPK kg/ha)	24844.5	46555.68	21711.18	0.87

\*Data was not subjected to statistical analysis.

## CONCLUSION

Based on the above findings it can be concluded that application of Phosphorus 50kg/ha along with Molybdenum 2kg/ha has performed better in growth parameters and yield attributes of cowpea (*Ankur Gomati-VU 89*) and also proven profitable. Since the findings are based on one season, further trails are needed to confirm the results.

## REFERENCES

- Anand, A., Umesha, C. and Sanodiya, L.K. (2022).** Effect of Phosphorus and Molybdenum on yield and economics of Black gram (*Vigna mungo* L.). *The Pharma Innovation Journal*. **11**(5):1417-1420.
- Bressani, R. and Elias, L.G. (1980).** Nutritional value of legume crops for human and animal. **135**.
- Chatterjee, R. and Bandyopadhyay, S. (2015).** Effect of boron, molybdenum and bio-fertilizer on growth and yield of cowpea (*Vigna unguiculata* L. Walp.) in acid soil of eastern Himalayan region. *Journal of the Saudi Society of Agricultural Sciences*.
- Dhaliwal S.S., Sharma, V., Shukla, A.K., Kaur, J., Verma, V., Kaur, M., Singh, P., Brestic, M., Gaber, A. and Hossain, A. (2022).** Interactive Effects of Molybdenum, Zinc and Iron on the Grain Yield, Quality, and Nodulation of Cowpea (*Vigna unguiculata* (L.) Walp.) in North-Western India. *Molecules* 2022, **27**, 3622.
- Franco, A.A. and Munns, D.N. (1981).** Response of *Phaseolus vulgaris* L. to molybdenum under acid conditions. *Soil Sci. Soc. Am. J.* **45**:1144-1148.
- Gad, N. and Kandil, H. (2013).** Evaluate the effect of molybdenum and different nitrogen levels on cowpea (*Vigna unguiculata*). *Journal of Applied Sciences Research*. **9**(3):1490-1497.
- Harireddy, Y.V. and Dawson, J. (2021).** Effect of bio-fertilizers and levels of *vermicompost* on

- growth and yield of cowpea (*Vigna unguiculata* L.). *The Pharma Innovation Journal*. **10(6)**:985-988.
- Harshini, K., Singh, V., George, S.G. and Singh, S.K. (2022).** Effect of Molybdenum and Bio-fertilizers on Growth and Yield of Cowpea (*Vigna unguiculata* L.). *International Journal of Plant & Soil Science*. **34(12)**: 92-98.
- Haruna, I.M. and Aliyu, L. (2011).** Yield and economic returns of sesame (*Sesamum indicum* L.) as influenced by poultry manure, nitrogen and phosphorus at Samaru, Nigeria. *Elixir Agric*. **39**:4884-4887.
- Karikari, B., Arkorful, E. and Addy, S. (2015).** Growth, Nodulation and yield response of cowpea to phosphorus fertilizer application in Ghana. *Journal of Agronomy*. **14(4)**:234-240.
- Khanna, R., Singh, A., Singh, S., Singh, V., Kumar, P. and Singh, S.P. (2019).** Effect of varying phosphorus levels and rhizobium inoculation on growth and yield of cowpea (*Vigna unguiculata* L.). *Progressive Research – An International Journal Society for Scientific Development in Agriculture and Technology*. **14 (4)**: 314-319.
- Kokani, J.M., Shah, K.A., Tandel, B.M. and Bhimani, G.J. (2015).** Effect of FYM, Phosphorus and Sulphur on yield of Summer Black gram and Post-Harvest Nutrient status of soil. *The Bioscan an International Quarterly Journal of Life Sciences*. **10(1)**:379-383.
- Mahesh, K., Umesha, C., Sanodiya, L.K. and Kumar, S.C. (2021).** Impact of Phosphorus levels and application of molybdenum of Black gram (*Vigna mungo* L.) on growth and yield attributes. *Biological Forum-An International Journal*. **13(3)**:12-15.
- Movalia, J., Donga, S. and Parmar, K.B. (2020).** Effect of boron and molybdenum on summer green gram (*vigna radiata* L.) (GM-4) under medium black calcareous soils: A Review. *SSRN Electronic Journal* · DOI: 10.2139/ssrn.3585103.
- Namakka, A., Jibrin, D.M., Hamma, I.L. and Bulus, J. (2017).** Effect of Phosphorus Levels on Growth and Yield of Cowpea (*Vigna unguiculata* (L.) Walp.) in Zaria, Nigeria. *Journal of Dryland Agriculture*. **3(1)**: 85 - 93.

- Ndakidemi, P.A. and Dakora, F.D. (2007).** Yield components of nodulated cowpea (*Vigna unguiculata*) and maize (*Zea mays*) plants grown with exogenous phosphorus in different cropping systems. *Aust. J. Exp. Agric.* **47**:583-589.
- Nkaa, F.A., Nwokeocha, O.W. and Ihuoma, O. (2014).** Effect of phosphorus fertilizer on growth and yield of cowpea (*Vigna unguiculata*). *IOSR Journal of pharmacy and Biological Sciences(IOSR-JPBS)*. **9(5)**:74-82.
- Pramanik, K. and Singh, R.K. (2003).** Effect of levels and mode of phosphorus and bio-fertilizer on chickpea under dryland conditions. *Indian journal of Agronomy*. **48**:294-296.
- Sheahan, C.M., (2012).** Plant guide for cowpea (*Vigna unguiculata*). *USDA- Natural Resources Conservation Service*, Cape Max Plant Materials Center, Cape May, NJ.
- Shilpa, H.D., Wali, S.Y., Hanamant, M.H. and Amrutha, T.G. (2015).** *International Journal of Technical Research and Applications*. **3(5)**:224-226.
- Singh, R., Singh, V., Singh, P. and Yadav, R.A. (2018).** Effect of phosphorus and PSB on yield attributes, quality and economics of summer green gram (*Vigna radiata* L.). *Journal of Pharmacognosy and Phytochemistry*. **7(2)**:404-408.
- Tahir, M., Sher, A. and Majeed, M.A. (2014).** Effect of molybdenum on yield and quality of black gram (*Vigna mungo* L.). *Pakistan Journal of Life and Social Sciences*. **12(2)**: 101-105.