

## Short Research Article

### Effect of planting geometry and nutrient management on yield, economics and quality of dwarf rice bean (*Vigna umbellata*) under rainfed condition

#### ABSTRACT

Rice bean (*Vigna umbellata*) is one of the under-utilized crops which can become a promising pulse crop. A field experiment was conducted during *kharif* season of 2020 at Research Farm, College of Agriculture, Central Agricultural University, Imphal to study the effect of planting geometry and nutrient management on yield, economics and quality of dwarf rice bean under rainfed condition. The experiment was laid out in Factorial Randomized Block Design (FRBD) with three replications. Results indicate that planting geometry at 60 cm × 10 cm ( $S_2$ ) and nutrient management ( $F_2$ ) i.e. application of 60 kg  $P_2O_5$ /ha with seed treatment of molybdenum + phosphate solubilizing bacteria (PSB) recorded significantly influenced the yield and quality of dwarf rice bean and gave the maximum gross return, net return and B:C ratio. It can be inferred that integration of nutrient management i.e. involving phosphatic fertilizer with molybdenum and phosphate solubilizing bacteria ( $F_2$ ) coupled with higher planting geometry ( $S_2$ ) i.e. (60 cm × 10 cm) significantly influenced the yield and quality of dwarf rice bean.

**Keywords:** Rice bean, planting geometry, nutrient management, yield, quality

#### Introduction

Pulses play a vital part in Indian agriculture. India is a leading pulse growing country and forms an essential part of cropping system of the farmers around the country. The present average per capita consumption of pulses in India was 14 kg per year against the WHO recommendation of 20 kg per year. India with 28 Mha of pulses cultivation area with the annual productivity at 885 kg/ha, has also increased significantly over last ~~05~~ five year (GOI, 2022). Pulses have the capacity to fix the soil atmospheric nitrogen in their root system, which enables the plants to meet the nitrogen requirement on its own. They are known to render significant impact on soil health and considered to be key component for sustainable agriculture (Laishram *et al.*, 2020). By 2050, the domestic requirements for pulse would be 26.50 M tons, necessitating stepping up production by 81.50%, i.e., 11.9 million tons additional produce at 1.86 % annual growth rates (Singh *et al.*, 2015). Thus, there is need to

**Comment [AAS1]:** Write information about treatment details in short in abstract. Write result in the comparative form such as increase in studied parameter over control or inferior treatment or recommended practices. Concluding statement should contain quantifiable information than saying/ writing just best treatment name. Is it nutrient management or Integrated nutrient management? Please made needful correction in title and over entire MS.

**Comment [AAS2]:** It should be specific- add "for NEH region" after "promising pulse crop"

**Comment [AAS3]:** Avoid repetition of words in title and keywords; write the nutrient sources, name of quality parameters, Location/region "NEH".

**Comment [AAS4]:** Lacking review of literature on same line (Selected crop explored in selected region before on national level); Clearly define the research gap for the present investigation.

**Comment [AAS5]:** Check spell of tonns?

increase the production and productivity of pulses in the country by more intensive interventions.

Rice bean (*Vigna umbellata*) is one of the under-utilized pulse crops which can become a promising pulse crop. This crop is a native of South and South East Asia from the Himalayas to South China and Indonesia (Chandel and Pant, 1982). In India, as cultigens, rice bean is mainly confined to the tribal regions of the North Eastern Hills, Western and Eastern Ghats in peninsular India (Arora *et al.*, 1980). In the North Eastern Region and also in the hilly regions of both North Bengal and Sikkim, rice bean is grown predominantly under rain fed condition in a mixed farming system, under shifting cultivation or in kitchen gardens and backyards. Despite being a crop that is thought to be underutilised, rice beans are valued for their high nutritional content in Manipur [state of India](#).

Planting geometry and nutrient management are of immense significance which is to be managed properly for higher rice bean production. Inappropriate planting times, high population density, and spatial configuration are some of the major causes of low production (Devi, 2021). Apart from non-existence of economically rational and socially optimum cropping pattern and crop diversification; biotic factors play major role in limiting rice bean production in rain-fed agriculture. Plant geometry plays an important role in the dominance and suppression during the process of competition. Ideal plant geometry is precious and important for better and efficient utilization of available plant growth resources in order to get maximum productivity in crops (Rana, 2011). Increase in yield can be ensured by maintaining appropriate plant population through different planting patterns. The influence of spacing on yield and quality of rice bean is still lacking in most of the rice bean growing areas in our country.

The North Eastern [Hilly region of](#) India has more than 80% hilly area of pH less than 5.5 with acid in nature and poor availability of P, Mo, Ca and Mg as well as the toxicity of Al, Fe, and Mn has been blamed for the soil's low productivity. Phosphorus serves a master nutrient for legume crop; it is involved in several plant functions, including energy storage and transfer, photosynthesis, transformation of sugars and starches, etc. Phosphorus application in rice bean promotes seed yield and dry matter production by enhancing root proliferation and nodulation. Yet, in acidic soil conditions, phosphorus is one of the main minerals that are limited for leguminous crops. Molybdenum a trace element found in the soil is required for growth of most biological organisms including plants (Agarwala *et al.*, 1978).

Phosphate solubilizing bacteria (PSB) play fundamental roles in biogeochemical phosphorus cycling in agro-ecosystems. It mobilize insoluble form of phosphorus in the rhizosphere into available form. And hence, application of such naturally occurring organic inputs as a whole improved the soil support system and ensured nutrient availability which was ultimately reflected in yield performance of the crop (Zaidi *et al.*, 2017). Keeping the above points in view, a field experiment was conducted to find out suitable planting geometry and nutrient management practices to assess their effect on yield and quality and economics of growing dwarf rice bean under rainfed condition.

### 1. **Materials and Methods**

A field experiment was conducted during *kharif* season of 2020 at Research Farm, College of Agriculture, Central Agricultural University, Imphal to study the effect of planting geometry and nutrient management on yield, quality and economics of dwarf rice bean (*Vigna umbellata*) under rainfed condition. The experimental site is located at 24.45°N latitude and 93.56°E longitudes at an elevation of 790 m above mean sea level. The soil of the experimental site was acidic with pH 5.1, available nitrogen 231.03 kg/ha, available phosphorous 10.95 kg/ha, available potash 150.00 kg/ha, available molybdenum 0.025 mg/kg and organic carbon 1.8 %. The experiment was laid out in Factorial Randomized Block Design (FRBD) with three replications consisting of three planting geometry treatments *viz.* (S<sub>1</sub>: 30 cm × 10 cm, S<sub>2</sub>: 45 cm × 10 cm and S<sub>3</sub>: 60 cm × 10 cm) and four nutrient management treatments *viz.* (F<sub>0</sub>: Seed treatment with molybdenum + PSB; F<sub>1</sub>: 20 kg P<sub>2</sub>O<sub>5</sub>/ha + seed treatment with molybdenum + PSB; F<sub>2</sub>: 40 kg P<sub>2</sub>O<sub>5</sub>/ha + seed treatment with molybdenum + PSB and F<sub>3</sub>: 60 kg P<sub>2</sub>O<sub>5</sub>/ha + seed treatment with molybdenum + PSB). The total rainfall received during the crop season was 977.1 mm.

The land was properly prepared by tractor driven cultivator and leveller. Adequate drainage channels were made in the field to drain the excess water as and when required. A uniform dose of 20 kg N/ha through urea and 20 kg K<sub>2</sub>O/ha through muriate of potash was applied to all the plots. Phosphorus was applied as per treatments through single superphosphate. The required quantity of different fertilizers as per treatments weighted separately and mixed together. After this, all fertilizers of each plot were applied in the rows at the time of sowing nearly 2 cm below the seed.

The rice bean variety used was a local variety and were treated with the mixture of 0.2% sodium molybdate per kg of seed just before sowing and 20 ml of phosphate solubilizing

**Comment [AAS6]:** Provide information about climatic condition and prevailing weather during crop growing season. Mention sowing time, sowing method and crop variety: Is seed rate is same in all three spacing? Provide the procedure name and reference followed for nitrogen analysis of grain.

**Comment [AAS7]:** Is this style is accepted? Or "kg ha<sup>-1</sup>"; Check it do needful correction.

bacteria (PSB) per kg of seed as per treatment. Sowing was done manually using 15 kg seeds/ha in the rows. The crop was infested with several weed species during early growth stage; hence, one hand weeding was done manually in all plots with the help of khurpi at 20 days after sowing (DAS). One spray of Emamectin benzoate 5 % (Proclaim) @ 1 liter/ha was given as a preventive measure at 30 DAS to control aphids and leafhoppers during early vegetative stage. Plant protection measures were provided as and when required.

The crude protein content was determined by multiplying percentage of nitrogen content in seed of rice bean with a factor of 6.25 (Piper, 1966).

$$\text{Crude protein \%} = \text{Nitrogen \%} \times 6.25$$

The crude protein yield in q/ha under a particular treatment was calculated by multiplying crude protein content with dry matter yield (q/ha) of the respective treatments

$$\text{Crude protein yield (kg/ha)} = \frac{\text{Crude protein content (\%)} \times \text{seed yield (q/ha)}}{100}$$

The economics of the experiment was worked out from the cost of inputs and income from the seed yield. The gross and net returns as well as the cost benefit ratio were worked out by using the formulae:

$$\text{Gross returns (Rs/ha)} = \text{Cost of seed yield of ricebean}$$

$$\text{Net returns (Rs/ha)} = \text{Gross returns} - \text{total cost of cultivation}$$

$$\text{Cost benefit ratio} = \frac{\text{Gross returns}}{\text{Total cost of cultivation}}$$

All the data obtained were statistically analyzed by the method of analysis of variance to test the significance of the treatment effects as well as result interpretation as given by Gomez and Gomez (1984).

## 2. Results and Discussion

### *Effect of planting geometry and nutrient management on yield (q/ha) of dwarf rice bean*

Perusal of data revealed that the effect of planting geometry and nutrient management significantly influenced the seed yield and stover yield of rice bean (Table 1). Increasing the planting geometry up to 60 cm × 10 cm (S<sub>3</sub>) gave the highest seed yield (14.66 q/ha), stover yield (89.71 q/ha) and harvest index (33.12%). Planting geometry (S<sub>1</sub>) i.e. 30 cm × 10 cm gave the lowest yield. This might be ascribed to less inter-plant competition and better

**Comment [AAS8]:** Avoid repetition/ duplicity of results and reading of table in text; Avoid writing S1, S2, S3.F0 ...F4, etc after treatment name (for entire MS); Combine table 1 and 2; Information on price of grain per quintal? Is price of stover is considered for calculation of returns? Results need to be discussed for protein yield and economics in more elaborative than just writing the "results are in conformity with ....."

availability of growth resources, such as light, moisture, nutrients, and space for one another in broader planting geometry, which had improved yield-attributing features and ultimately raised the yield of rice bean. These results are in conformity with the findings of Goud *et al.* (2016) and Ramanjaneyulu *et al.* (2017). Among the different nutrient management practices,

Treatments	Seed yield (q/ha)	Stover yield (q/ha)	Harvest index (%)	Gross return (Rs/ha)	Net return (Rs/ha)	B:C
------------	-------------------	---------------------	-------------------	----------------------	--------------------	-----

the highest seed yield (16.01 q/ha), stover yield (105.18 q/ha) and harvest index (33.37%) were recorded in (F<sub>3</sub>) *i.e.* 60 kg P<sub>2</sub>O<sub>5</sub>/ha + seed treatment with molybdenum+ PSB and it was found lowest in seed treatment with molybdenum + PSB (F<sub>0</sub>). Such increase in yield might be due to the increase in vegetative growth with the increasing doses of phosphorus. This result is in agreement with those reported by Bhuiyan *et al.* (2016) and Nissa *et al.* (2017).

Integration of phosphatic fertilizer with molybdenum and phosphate solubilizing bacteria significantly influenced the seed yield and stover yield of rice bean. The treatment combination (S<sub>3</sub>F<sub>3</sub>) *i.e.* 60 cm × 10 cm and 60 kg P<sub>2</sub>O<sub>5</sub>/ha + seed treatment with molybdenum + PSB gave the highest seed yield (17.10 q/ha), stover yield (105.84 q/ha) and harvest index (37.70%) while the least was observed in treatment combination of 30 cm × 10 cm and seed treatment with molybdenum + PSB (S<sub>1</sub>F<sub>0</sub>). Such increase in the yield with the application of phosphorus is obvious due to acidic condition of the soil where the availability of phosphorus was limiting and in the medium range. Further, the seed treatment with molybdenum might have enhanced the nitrogen fixation resulting in better growth and yield. The optimum activities of phosphate solubilizing bacteria also increase the availability of phosphorus to the plant which ultimately increases the yield of rice bean. These findings are in line with those obtained by Yadav and Yadav (2011) and Manohar (2014).

***Effect of planting geometry and nutrient management on economics (Rs/ha) of dwarf rice bean***

The data presented in (Table 1) revealed that the planting geometry (S<sub>3</sub>) *i.e.* 60 cm × 10 cm gave the maximum gross return (117240 Rs/ha), net return (71335 Rs/ha) and B:C (2.55). The lowest was for the planting geometry (S<sub>1</sub>) *i.e.* 30 cm × 10 cm. Among the different nutrient management practices, the maximum gross return (128054 Rs/ha), net return (80877 Rs/ha) and B:C (2.72) was found in (F<sub>3</sub>) *i.e.* 60 kg P<sub>2</sub>O<sub>5</sub>/ha + seed treatment with molybdenum+ PSB and it was found lowest in seed treatment with molybdenum + PSB (F<sub>0</sub>).

<b>Planting geometry</b>						
30 cm × 10 cm (S <sub>1</sub> )	13.39	78.81	29.11	106960	60305	2.29
45 cm × 10 cm (S <sub>2</sub> )	14.07	83.72	31.81	112320	66165	2.45
60 cm × 10 cm (S <sub>3</sub> )	14.66	89.71	33.12	117240	71335	2.55
S. Em (±)	0.29	2.55	0.73	-	-	-
C.D. (P = 0.05)	0.60	5.29	1.51	-	-	-
<b>Nutrient management</b>						
Seed treatment with molybdenum + PSB (F <sub>0</sub> )	11.71	65.91	28.91	93360	48058	2.07
20 kg P <sub>2</sub> O <sub>5</sub> /ha + seed treatment with molybdenum+ PSB (F <sub>1</sub> )	12.71	75.45	29.31	101654	55728	2.22
40 kg P <sub>2</sub> O <sub>5</sub> /ha + seed treatment with molybdenum+ PSB (F <sub>2</sub> )	15.71	89.79	33.80	125627	79076	2.70
60 kg P <sub>2</sub> O <sub>5</sub> /ha + seed treatment with molybdenum+ PSB (F <sub>3</sub> )	16.01	105.18	33.37	128054	80877	2.72
S. Em (±)	0.33	2.94	0.84	-	-	-
C.D. (P = 0.05)	0.688	6.10	1.75	-	-	-
<b>Interaction (Planting geometry × Nutrient management)</b>						
S <sub>1</sub> F <sub>0</sub>	11.27	59.83	28.06	90080	44361	1.9
S <sub>1</sub> F <sub>1</sub>	12.65	74.07	28.62	101120	54777	2.1
S <sub>1</sub> F <sub>2</sub>	14.19	76.92	29.62	113440	66472	2.4
S <sub>1</sub> F <sub>3</sub>	15.41	104.42	30.13	123200	75607	2.5
S <sub>2</sub> F <sub>0</sub>	11.53	64.10	29.30	91440	45721	2.0
S <sub>2</sub> F <sub>1</sub>	12.67	75.78	30.02	101280	54937	2.1
S <sub>2</sub> F <sub>2</sub>	16.55	89.74	35.65	132400	85432	2.8
S <sub>2</sub> F <sub>3</sub>	15.53	105.27	32.29	124160	76567	2.6
S <sub>3</sub> F <sub>0</sub>	12.32	73.79	29.36	98560	52841	2.1
S <sub>3</sub> F <sub>1</sub>	12.82	76.50	29.29	102560	56217	2.2
S <sub>3</sub> F <sub>2</sub>	16.38	102.71	36.13	131040	84072	2.7
S <sub>3</sub> F <sub>3</sub>	17.10	105.84	37.70	136800	89207	2.8
S. Em (±)	0.57	5.10	1.46	-	-	-
C.D. (P = 0.05)	1.19	10.58	3.03	-	-	-

**Table 1. Effect of planting geometry and nutrient management on yield and economics of dwarf rice bean**

The treatment combination (S<sub>3</sub>F<sub>3</sub>) i.e. 60 cm × 10 cm and 60 kg P<sub>2</sub>O<sub>5</sub>/ha + seed treatment with molybdenum + PSB gave the maximum gross return (136800 Rs/ha), net return (89207 Rs/ha) and benefit cost ratio (2.8); while the least was observed in treatment combination of 30 cm × 10 cm and seed treatment with molybdenum + PSB (S<sub>1</sub>F<sub>0</sub>). The increase in net return might be due to increase in yield of dwarf rice bean and relatively low production cost per unit of yield under the treatment. The higher benefit cost ratio in treatments S<sub>3</sub> and F<sub>3</sub> (i.e. 60 cm × 10 cm and 60 kg P<sub>2</sub>O<sub>5</sub>/ha + seed treatment with molybdenum + PSB) might be due to the higher gross and net return with comparative lower

cost of cultivation in compare with other treatments. The result is in conformity with the findings of Rajeshkumar *et al.* (2017) and Vikram *et al.* (2018).

#### ***Effect of planting geometry and nutrient management on quality of dwarf rice bean***

The effect of planting geometry and nutrient management significantly influenced the crude protein content and protein yield of rice bean (Table 2). Increasing the planting geometry upto 60 cm × 10 cm (S<sub>3</sub>) gave the highest crude protein content (20.32%) and protein yield (42.56 kg/ha) of rice bean. It was found lowest in planting geometry 30 cm × 10 cm (S<sub>1</sub>). Wider spacing has less plant population than the narrow spacing so the nutrient content in seed is more in the wider spacing as there is less competition among the plants. This result is in conformity with the findings of Rath *et al.* (1997).

Among the nutrient management practices, the highest crude protein content (21.67%) and crude protein yield (49.77 kg/ha) were recorded in 60 kg P<sub>2</sub>O<sub>5</sub>/ha + seed treatment with molybdenum+ PSB (F<sub>3</sub>) and it was found lowest in seed treatment with molybdenum + PSB (F<sub>0</sub>). The higher protein content in this treatment may be due to adequate availability of phosphorus besides the contribution from the phosphate solubilizing bacteria which play a major role in protein synthesis. This observation was supported by the findings of Singha and Sarma (2001) and Gupta *et al.* (2012).

Integration of phosphatic fertilizer with molybdenum and phosphate solubilizing bacteria significantly influenced the crude protein content and crude protein yield of rice bean. The treatment combination (S<sub>3</sub>F<sub>3</sub>) *i.e.* 60 cm × 10 cm and 60 kg P<sub>2</sub>O<sub>5</sub>/ha + seed treatment with molybdenum + PSB gave the highest crude protein content (23.26%) and crude protein yield (48.80 kg/ha) while the least was observed in treatment combination of 30 cm × 10 cm and seed treatment with molybdenum + PSB (S<sub>1</sub>F<sub>0</sub>). The results are in agreement with the finding of Gupta *et al.* (2012).

On the basis of findings, it was concluded that a wider crop spacing of 60 cm × 10 cm along with a higher dose of phosphorus 60 kg P<sub>2</sub>O<sub>5</sub>/ha application with seed treatment of molybdenum + phosphate solubilizing bacteria (PSB) proved to be superior over narrow spacing and low dose phosphorus application by giving best results in terms of yield, economics and quality of growing dwarf rice bean under rainfed conditions of Manipur.

**Comment [AAS9]:** Write the conclusion in quantifiable form and not in way it was written (It is clear from this conclusion how much advantages has been expect from best treatment over control or other treatment.

**Table 2. Effect of planting geometry and nutrient management on yield of rice bean**

Treatments	Crude protein content (%)	Crude protein yield (kg/ha)
<b>Planting geometry</b>		
30 cm × 10 cm (S <sub>1</sub> )	19.32	36.58
45 cm × 10 cm (S <sub>2</sub> )	19.91	41.87
60 cm × 10 cm (S <sub>3</sub> )	20.32	42.56
S. Em (±)	0.35	1.23
C.D. ( <i>P</i> = 0.05)	0.72	2.54
<b>Nutrient management</b>		
Seed treatment with molybdenum + PSB (F <sub>0</sub> )	17.61	27.53
20 kg P <sub>2</sub> O <sub>5</sub> /ha + seed treatment with molybdenum+ PSB (F <sub>1</sub> )	18.92	35.35
40 kg P <sub>2</sub> O <sub>5</sub> /ha + seed treatment with molybdenum+ PSB (F <sub>2</sub> )	21.20	48.69
60 kg P <sub>2</sub> O <sub>5</sub> /ha + seed treatment with molybdenum+ PSB (F <sub>3</sub> )	21.67	49.77
S. Em (±)	0.40	1.42
C.D. ( <i>P</i> = 0.05)	0.84	2.93
<b>Interaction (Planting geometry × Nutrient management)</b>		
S <sub>1</sub> F <sub>0</sub>	17.44	24.00
S <sub>1</sub> F <sub>1</sub>	18.65	30.78
S <sub>1</sub> F <sub>2</sub>	20.19	44.77
S <sub>1</sub> F <sub>3</sub>	20.99	46.75
S <sub>2</sub> F <sub>0</sub>	17.43	24.79
S <sub>2</sub> F <sub>1</sub>	19.28	37.79
S <sub>2</sub> F <sub>2</sub>	22.16	51.15
S <sub>2</sub> F <sub>3</sub>	20.77	53.76
S <sub>3</sub> F <sub>0</sub>	17.94	33.81
S <sub>3</sub> F <sub>1</sub>	18.82	37.49
S <sub>3</sub> F <sub>2</sub>	21.25	50.15
S <sub>3</sub> F <sub>3</sub>	23.26	48.80
S. Em (±)	0.70	2.45
C.D. ( <i>P</i> = 0.05)	1.45	5.08

## References

Agarwala SC, Sharma CP, Farooq S, Chatterjee C. Effects of molybdenum deficiency on the growth and metabolism of corn plants raised in sand culture. *Canadian Journal of Botany*. 1978; 56:1905-1909.

Arora R, Chandel KPS, Joshi BS, Pant KC. Rice bean: Tribal pulse of eastern India. *Econ Bot*. 1978;34(3): 260-263.

**Comment [AAS10]:** Avoid citing old references and cross check the reference for their presence in both in text and reference list; Strictly follow the reference writing style of the journal both for citing references in the text and enlisting them in the reference list.

Bhuiyan MMH, Rahman MM, Afroze F, Sutradhar GNC, Bhuiyan MSI. Effect of phosphorus, molybdenum and *Rhizobium* inoculation on growth and nodulation of mungbean. J Soil Nat. 2008;2(2): 25-30.

Chandel KPS, Pant KC. Genetic resources of *Vigna* species in India, their distribution, diversity and utilization in crop improvement. Annals of Agricultural Sciences. 1982;3(1-2):19-34.

Devi KM, Luikham E, Singh LN, Devi NS, Singh NG. Influence of planting geometry and nutrient management on growth, nodulation and yield of dwarf Rice bean (*Vigna umbellata*) under rainfed condition. The Pharma Innovation Journal. 2021;10(9): 1766-1770.

Gomez KA, Gomez AA. Statistical Procedure for Agricultural Research, 2nd Ed. John Wiley and Sons, New York. 1984.

Goud VV, Konde NM, Patil AN. Effect of planting geometry on growth, nutrient uptake and yield of medium duration pigeonpea hybrid under rainfed condition. PKV Res J. 2016;40(1): 10-14.

Government of India. Annual Report: 2021-2022. Ministry of Agriculture & Farmers Welfare, Directorate of Pulses Development. DPD/Pub./TR/45/2021-22. 2022.

Gupta SC, Gangwar S. Effect of molybdenum, iron and microbial inoculants on symbiotic traits, nutrient uptake and yield of chickpea. Journal of Food legumes. 2012;25(1): 45-9.

Keerthi MM, Babu R, Joseph M, Amutha R. Optimizing plant geometry and nutrient management for grain yield and economics in irrigated greengram. American Journal of Plant Sciences. 2015;6: 1144-1150.

Khanda CM, Mahapatra AK, Misra PK. Response of ricebean (*Vignaumbellata*) to row spacing and phosphorus under rainfed condition. Annals of Agricultural Research. 2001;22: 481-484.

Laishram B, Singh TB, Kalpana A, Wangkheirakpam M, Chongtham SK, Singh WJ. Effect of Salicylic Acid and Potassium Nitrate on Growth and Yield of Lentil (*Lens culinaris* L.) under Rainfed Condition. Int J Curr Microbiol App Sci. 2020;9(11): 2779-2791.

Manohar MS. Effect of sulphur and molybdenum on growth and productivity of fenugreek. M.Sc. Thesis, SKNAU, Jobner. 2014.

Nissa un Seerat, Bashir S, Dar SA, Baba JA, Hakeem SA, Wani RA, Mughal, M.N, Basu YA. Effect of *Rhizobium* and PSB in combination with phosphorus on the enrichment of soil fertility and its effect on yield of green gram (*Vigna radiata* L.). Int J Curr Microbiol Appl Sci. 2017;6(11): 3648-3652.

Piper CS. Soil and plant analysis. Hans Publication, Bombay. 3rd Ed., pp. 49-54. 1966.

Prasad G, Bhat BB, Kar BC, Mishra SN. Response of ricebean (*Vigna umbellata*) to spacing and levels of nitrogen. Indian Journal of Agronomy. 1994;39(3): 485-487.

Rahman MM, Bhuiyan MMH, Sutradhar GNC, Rahman MM, Paul AK. Effect of phosphorus, molybdenum and *Rhizobium* inoculation on yield and yield attributes of mungbean. International Journal of Sustainable Crop Production. 2008;3(6):26-33.

Rajeshkumar S, Durairaj SN, Kannan V. Effect of crop geometry and foliar nutrition on growth and yield of irrigated blackgram (*Vigna mungo* L.). International Journal of Current Microbiology and Applied Sciences. 2017;6(11): 4084-4094.

Ramanjaneyulu, A.V., Indudhar Reddy, K., Spandana Bhatt, P., Neelima, T.L, Srinivas, A. Influence of pigeon pea varieties, N levels and planting methods on yield and economics under rainfed conditions. Legum Res. 2017;40(5): 911-915.

Rana MM, Chowdhury SH, Bhuiya MS. Effects of plant population and bio-fertilizer on the growth parameters of three summer mungbean (*Vigna radiata* L.) cultivars. Bangladesh Journal of Agricultural Research. 2011;36: 537-542.

Rath BS, Tripathy RK, Nayak SP, Kar BC. Effect of row spacing on yield and seed quality of ricebean (*Vigna umbellata*) varieties. Indian Journal of Agronomy. 1997;42(4):717-719.

Singh AK, Singh SS, Prakash V, Kumar S, Dwivedi SK. Pulses production in India: Present status, bottleneck and way forward. Journal of Agri Search. 2015;2(2):75-83.

Singh DP, Rajput AL, Singh SK. Response of French bean to spacing and nitrogen levels. Indian Journal of Agronomy. 1996;41:608-610.

Singha DD, Sarma A. Response of blackgram (*Phaseolus mungo*) to phosphorus fertilization and *Rhizobium* inoculation in the hill soils of Assam. Annals of Agricultural Research. 2001;22(1): 151-153.

Vikram S, Ajeev K, Manohar L, Mahesh K. Chickpea cultivation governing by phosphorus application in calcareous soil. *International Journal of Current Microbiology and Applied Sciences*. 2018;7(02):3419-3425.

Yadav SL, Yadav LR. Effect of phosphorus and biofertilizers on productivity of garden pea under semiarid conditions. *J Food Legum*. 2011;24(3): 248-249.

Zaidi A, Khan MS, Rizvi A, Saif S, Ahmad B, Shahid M. Role of phosphate-solubilizing bacteria in legume improvement. In: *Microbes for Legume Improvement*, Springer, Chambridge, pp 175-197. 2017.

UNDER PEER REVIEW