

MICROBIOLOGICAL EVALUATION OF FRESH CATFISH (*Clarias gariepinus*) OBTAINED FROM SELECTED MARKETS AND PONDS IN MINNA METROPOLIS

ABSTRACT

Microbiological evaluation of fresh catfish (*Clarias gariepinus*) sold in three different markets and ponds in Minna Metropolis were carried out. The following bacteria were discovered *Escherichia coli*, *Bacillus spp*, *Staphylococcus spp*, *Pseudomonas species*, *Streptococcus species*, *Shigella species* and *Enterobacter species*. Bosso market skin *Escherichia coli*, *Staphylococcus aureus*, *Aspergillus niger*, *Streptococcus spp*, *Rhizopus species*. Kure market *Pseudomonas aeruginosa*, *Escherichia coli*, *Aspergillus*, *Staphylococcus spp*, *Aspergillus*, *Bacillus subtilis*. Mobil market, *Escherichia coli*, *Rhizopus spp*, *Staphylococcus aureus*, *Aspergillus flavus*. Bosso ponds *Enterobacter spp*, *Pseudomonas spp*. Mobil pond *Streptococcus spp*. Kure pond *Escherichia coli*. Bosso pond (Skin) 3.8×10^2 Gills 3.0×10^2 (Satisfactory) Kure pond (Skin) 2.4×10^2 Satisfactory Gills 2.6×10^2 (Satisfactory) Mobil pond The bacteria load count is ranged from 4.0×10^5 cfu/mL and 1.64×10^3 cfu/mL. The fungi isolated were *Rhizopus spp* and *Aspergillus niger*. The fungi count from the catfish sample is ranged from 0.5×10^1 cfu/mL and 2.0×10^3 cfu/mL. This shows that bacteria and fungi contaminations occurred in catfish obtained from three different markets location in Minna, Niger state.

Keywords: Microbiological, catfish (*Clarias gariepinus*), markets, ponds.

1.0 INTRODUCTION

The last two decades have seen appreciable increase in global fish trade and the need to enforce safety standards and regulations on imported consignment especially from developing nations fraught with unacceptable levels of microbiological contamination (ICMSF, 2005).

Contamination concern has been on high loads of unsuspected spoilage by microorganisms like *Salmonella sp*, *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Escherichia coli*. (Gram *et al.*, 2000). Many spp of fish (approximately 6850) live in fresh water lakes and rivers. Fish is mainly thought of as a source of proteins (Daniele *et al.*, 2006). Fishing in Minna metropolis is mainly done by the Gwaris and Gbagyl tribes and the women are responsible for the sale especially in Shiroro and Gurara locals in Niger state of Nigeria. The microorganisms associated with fresh catfish in harveste from rivers and ponds in Minna is essential in order to establish an overview of the bacteriological pathogens associated with fresh fish in Niger and compare with other parts of the world (Daniel *et al.*, 2006)

Comment [EM1]: In writing your abstract, I suggest you write base on the following:
Aim of the study
Summary of the methodology
Summary of your results
Conclusion
Note: you may not write each of the above bullet, but make sure you explain them all.

Comment [EM2]: Check the name of the organism

Comment [EM3]: spp.

Comment [EM4]: You should rewrite your abstract, and used full stop, and comma where necessary

Comment [EM5]: These are just common key words, use standard key words that will distinguished your work.

Comment [EM6]: Rewrite your introduction to note more than one page.

Comment [EM7]: No full stop before the citation.

Comment [EM8]: Check this and add space before the word fishing, also, remove space in the bracket.

Comment [EM9]: Check spelling

Comment [EM10]: ????

Comment [EM11]: Delete

Spoilage patterns of fish have been well documented (Botta, 2007) its size. Fresh fish spoilage and high perish ability are primarily due to large amount of non-protein nitrogen (like free amino acids), volatile nitrogen bases (ammonia, creatine, taurine, uric acid, carnosine and histamine) which support post mortem bacteria growth (Jay *et al.*, 2005).

Comment [EM12]: Delete space

Acuff and Finnereported that heating, freezing and exposure to high concentration of salt lead to chemical and physical changes, and therefore digestibility increased due to protein denaturation, but content of thermo labile compound and polyunsaturated fatty acid is often reduced.

Comment [EM13]: Add the year

Fish have been known to harbor micro flora. These are microorganisms and are found mostly in their gills and gut. Microbes may become intimately associated with the external surfaces of the fish or may be incorporated into the resident micro flora (Austin and Austin, 1987). The catfish species in Nigeria are *Clariasgariepinus*, *Heterobanchusbidorsalis*, *Clariasheterobranchus hybrid (Heteroclarias)*, and *Clariasnigrodigitatus*.

Heterobranchus species is the most commonly cultured fish in south eastern part of Nigeria; African catfish is popular in the market and has great potential to boost the rapidly growing Nigerian aquaculture (Olaleye and Adewumi 2010). Generally, the fact that fish farming affords best utilization of land and provides income to an investor; it is one of the cheapest sources of animal protein.

Fish spoilage is one of the greatest problems affecting the fishing industry world wideand attempts are being made to reduce the spoilage to the barest minimum. About 20 to 50% of domestic fish productions are wasted through spoilage in tropical developing countries Eyo (2001)

Comment [EM14]: worldwide

The broad objective of this work is to evaluate the microorganisms associate with cat fish (*Clariasgariepinus*) sample obtained from selected ponds in Minna metropolis. To estimate the microbial load of fresh catfish obtained from the pond and market in MinnaMetropolis, to isolate and identify micro-organisms associated with fresh catfish obtained from the ponds andMarkets location in Minna metropolis.

2.0 MATERIALS AND METHODS

2.1 Sources of materials and preliminary handling

Comment [EM15]: add the following:
2.1 Study area
2.2 sample collection

Samples of fresh catfish *Clarias gariepinus*. Fresh fish were randomly selected and purchased from Kure market, Mobil fish market and Bossomarket on various occasions, 9.am in a specific time of the day. These samples were conveyed in different polythene bags surround by ice block beside the polythene bags to the laboratory where the practical was carried out and the sample was kept in the freezer for about 2 hours before analysis were carried out on each species.

2.2 Method of analysis

The bacteriological media and reagent used were prepared and sterilized by autoclaving at temperature of about 121°C for 15min. Sterile water (100 mL) was poured into each polythene bags containing one of each species and was used to thoroughly wash the skin of the fish. And 1.0 mL of wash water was diluted to 10 fold in phosphate buffered saline. Various culture media are namely: Nutrient agar, MacConkey agar, and potato dextrose agar (P.D.A) were used. Others were test tube, string (injection needle), cotton wool, standing cork, and Petri-dishes. The gill were aseptically removed and weighed in various sample and recorded and was similarly treated like the skin.

2.2.1 Procedure for preparing the media

2.2.2 Nutrient agar

Nutrient agar was used for cultivation of bacteria, it is called general purpose agar medium. it contain beef extract 3.0g, agar-agar 15g, gelysate peptone 5.0g, final pH of 6.8. Therefore 28g of nutrient agar was dissolved in one liter (1000ml) of distill water, and then sterilized at 121°C for 15 min.

2.2.3 Potato dextrose agar (PDA)

Twenty four grams of P.D.A powder were dissolved in 500 mL of distilled water allow it to dissolved gently and autoclave it at 121°C for 15 min.

2.2.4 MacConkey agar

24g of MacConkey agar were dissolved in 500 mL of distilled water and autoclave it at 121°C for 15 min.

2.3.0 Serial Dilution Method

9.0 mL of distilled water was dispensed into test-tube sterilize by autoclaving at 121°C for 15 min and allow it cool, five in number for each sample carried out (I.e. skin and gill for every sample). 1.0 mL of sample washed from the skin was dispensed into the test-tube serially and mixed it thoroughly and label 10^{-1} and introduced into the second test-tube. It is repeated up to 10^{-5} for each sample.

Comment [EM16]: Media preparation is a general procedure that must be done before isolation of microorganisms in any research setting. Hence, it is not required in the manuscript and must be remove.

Comment [EM17]: Acknowledge the source of the methodology

1.0 mL was also pipette from each test-tube and place in petri-dish and 15.0 mL of nutrient agar was added to it and rocked on the table to mix, and allow to gel. Then the Petri-dishes were taken to incubator incubated at 30°C for 72 h. The plates were placed in colony counter and recorded.

2.4 Fungi

Serial dilution for fungi culture procedure is the same with the above but potatoes dextrose agar was used instead of nutrient or MacConkey agar.

2.5.1 Total and viable plate counts

The plate count method is most frequently employed for the enumeration of bacterial cells in a given sample. The plate count can be performed by using the pour plate method. In these methods overcrowding of colonies on plate should be avoided as this leads to inaccuracies in the count. Normally, plates with count ranging from 30 to 300 colonies are considered for population estimation. Therefore select a dilution which yields fewer than 300 colonies per plate since with colonial concentrations exceeding this count will usually be depressed to an unknown degree by overcrowding and micro antagonism.

2.5.2 Enumeration of micro organisms

To determine the number of microorganism in the fish sample is very important, in determined the safety of the fish product depending on knowing the level of microorganism present in particular fish product. Sample are taken for total number of living and dead cells, or by estimation of the number of living organism per mL this technique is used to assess the quality of the sample.

2.6. Bacteria Gram Stain Reaction

2.6.1 How to Gram Stain

Gram staining is a quick procedure used to look for the presence of bacteria in tissue samples and to characterize bacteria as Gram-positive or Gram-negative, based on the chemical and physical properties of their cell walls.

The Gram stain is always been done as the first step in diagnosis of a bacterial infection.

2.6.2 Steps to be followed in Gram Staining

Add 1-2 drops of the tissue sample onto a glass slide



Comment [EM18]: Remove the space

Comment [EM19]: Remove the space

Comment [EM20]: 2.5.1 represents the enumeration of microorganisms, and the method used enumerate both living and death cells.

Comment [EM21]: 2.6 Identification of Microorganisms
2.6.1 Identification of Bacteria
The bacteria isolated in the study will be identified using standard techniques. Gram's staining techniques and certain biochemical techniques such as indole test, citrate utilization test, motility test, Voges-Proskauer test, as described by et al. (yeay).
2.6.2 Identification of Fungi

Note: you should provide the citation to all the methodologies used in the study.

Also, for identification of fungi

Comment [EM22]: Delete

Comment [EM23]: Delete

Heats fix the smear, by quickly passing it two to three times through a flame



Flood the smear with crystal violet / gently rinse off the crystal violet with tap water



Flood the smear with iodine / gently rinse off the iodine with tap water



Decolorize by adding alcohol or acetone to the smear while holding the slide at an angle to allow the decoloriser to drain



Gently rinse off excess decolorized with tap water



Flood the smear with safranin counter stain / gently rinse off excess safranin with tap water



Drain slide and allow it to air dry



Examine the slide under the light microscope

Fig 1: Flow diagram of gram staining bacteria.

Comment [EM24]: Remove. It is not necessary

2.6.4 Test for Gram-positive and Gram-negative Bacterial Identification

Gram Staining Procedure

In this process most bacteria have one of these two types of cell walls; we can use this difference as a feature that can be identified using the Gram stain. The Gram stain is a differential stain that uses two dyes to differentiate between the two basic bacterial cell wall types. The inoculate bacterial sample was taken from the skin and gill which was incubated.

First a bacterial smear must be heat fixed to a microscope slide. A smear is a sample of bacteria suspended in a small amount of water on a slide. That sample is then dried using heat. The heat kills the bacteria and attaches the sample to the slide so that it does not easily wash away.

The Gram staining procedure goes as follows:

- Flood the slide with Crystal Violet (the primary stain).
- After 1 minute, rinse the slide with water.
- Flood the slide with Iodine (Iodine is a mordant that binds with Crystal violet and is then unable to exit the Gram positive peptidoglycan cell wall.)

- After 1 minute, rinse the slide with water.
- Flood the slide with Acetone Alcohol. (Alcohol is a decolorizer that will remove the stain from the Gram-negative cells.)
- After 10 or 15 seconds, then rinse the slide with tap water. (Do not leave the decolorized for too long or it may remove stain from the Gram-positive cells as well.)
- Flood slide with Safranin (the counterstain or additional stain for microscope specimen).
- After 1 minute, rinse the slide with water.
- Gently blot the slide dry. It is now ready to be viewed under oil immersion (1000x TM) with a bright-field compound microscope.

After this staining procedure, the Gram positive (+) cells will appear purple, having retained the primary stain. The Gram negative (-) cells will appear pink; having retained the counter stain after the primary stain was removed by the decolorizer.

Comment [EM25]: Delete

2.6.5 Indole test:

Indole test is employed to determine the ability of isolate to break down the amino acid.

Material: test tube contains 1% peptone water, bacterial culture, kovac's reagent.

Inoculate the peptone water with the bacterial culture.

Incubate for 48 hours at 37^oc. Add 0.5ml of kovac's reagent and shake gently.

Presence of red color indicates indole (positive).

Absence of red color indicates not indole (negative).

2.6.6 Citrate Utilization test.

Tests for the ability of bacteria to convert citrate (an intermediate of the Krebs cycle) into oxaloacetate (another intermediate of the Krebs cycle) and also help in identification of some members of the family Enterobacteriaceae. In this media, citrate is the only carbon source available to the bacteria. The media will turn a bright blue as a result of an increase in the pH of the media. To inoculate this slant, use the transfer loop. Inoculate the citrate agar plate with organism and inoculate at 30^oc for 72 hours for organism to grow.

2.6.8.1 MR-VP (Methyl red-Voges Proskauer)

This test is used to determine two things. The MR portion (methyl red) is used to determine if glucose can be converted to acidic products like lactate, acetate, and formate. The VP portion is used to determine if glucose can be converted to acetoin.

These tests are performed by inoculating a single tube of MR-VP media with a transfer loop and then allowing the culture to grow for 3-5 days. After the culture is grown, about half of the culture is transferred to a clean tube. One tube of culture will be used to conduct the MR test, the second tube serves as the VP test.

MR (Methyl red) test:

Methyl red is added to the MR tube. A red color indicates a positive result (glucose can be converted into acidic end products such as lactate, acetate, and formate). A yellow color indicates a negative result; glucose is converted into neutral end products


2.6.8.2 VP (Voges Proskauer) test:

First alpha-naphthol (also called Barritt's reagent A) and then potassium hydroxide (also called Barritt's reagent B) are added to the VP tube. The culture should be allowed to sit for about 15 minutes for color development to occur. If acetoin was produced then the culture turns a red color (positive result); if acetoin was not produced then the culture appears yellowish to copper in color (a negative result).

2.6.9 Catalase:

This test is can be used to detect the enzyme catalase. This enzyme is responsible for protecting bacteria from hydrogen peroxide (H_2O_2) accumulation, which can occur during aerobic metabolism. If hydrogen peroxide accumulates, it becomes toxic to the organism. Catalase will breaks H_2O_2 down into water and O_2 . To perform the catalase test simply smear a small amount of the test organism onto the lid of a Petri plate/culture dish. Then add a drop of hydrogen peroxide to the smear. If bubbles become visible (these would be the O_2 bubbling up) then the test is positive and you can conclude that the organism makes catalase. A lack of bubbles indicates the absence of catalase. Because most aerobic organism make catalase while anaerobic organism cannot.

Equation.



Smear small amount of test organism onto slide.

Then add a drop of hydrogen peroxide to the smear and watch for the bubble

Presence of gas bubbling up is positive

Absence of gas bubbling up is negative.

2.6.10 Oxidase test

To perform this test simply swab some of your test culture into one of the boxes on an oxidase dry slide. If a color change to purple or blue is evident at 30 sec-1 minute then the result is positive. It is important that the test is read by one minute to ensure accurate results (avoid false negatives and false positives). If it remain unchanged is negative.

2.6.11 Coagulase test

This test use to differentiate positive from negative bacterial in the presence of coagulation. The enzyme coagulase clots plasma by converting fibrinogen to fibrin.

Procedure

- A drop of physiological saline was placed on each clean slide.
- Emulsify a colony of the test organism with wire loop on each of the drops to make thick suspensions.
- Add a drop of plasma or serum to the suspensions and mix gently. Look for clumping of organism within 10 seconds
- Presence of agglutination give positive.
- Absence of agglutination give negative.

2.6.12 Motility test

Some bacterial genera can move actively from one place to another; others that are not endowed with the requisite locomotory structure such as flagella cannot move. The ability to test this is important in their identification as it helps to differentiate mobile from non mobile organisms. The determination of motility among bacterial species can be demonstrated by hanging drop or stab culture in motility medium. A positive motility test is indicative of the presence of locomotory organ in the test isolate.

Comment [EM26]: Acknowledge the source of the methodologies

Comment [EM27]: 3.0 Results

3.1 Microbiological Load of Catfish sample from skin and gills

Table 1 shows the result for the control experiment. Table 2 and 3 showed the microbial loads, microscopic and biochemical characteristics of the bacterial associate with catfish in three market locations in minna. The total coliform unit count (cfu/ml) of fish sample in Kure market is 1.64×10^3 cfu/ml and 7.2×10^4 cfu/ml getting from skin and gills. The total coliform counts on skin and gills of fish sample in Mobil market were 6.0×10^6 cfu/ml and 4.8×10^4 cfu/ml. (Daniel *et al.*, 2016) The total coliform unit counts in Bosso market were 5.6×10^4 cfu/ml and 4.0×10^5 cfu/ml getting from skin and gills. The total number of fungi colony count found in catfish sample from both skin and gills found in Kure market were 2.0×10^3 cfu/ml and 1.0×10^2 cfu/ml. while fungi colony found in Mobil market were 0.1×10^4 cfu/ml and 0.7×10^2 cfu/ml from both skin and gills. And fungi colony found in Bosso markets were 1.3×10^2 cfu/ml and 0.5×10^1 cfu/ml from both skin and gills respectively. In summary therefore, Table 4 shows the bacterial identified from isolate were *Escherichia coli*, *Staphylococcus spp*, *Streptococcus spp*, *Bacillus*, *Shigella*, *Enterobacterspp* and *Pseudomonas*. While Table .5 shows the fungi species isolated namely; were *Aspergillus spp* and *Rhizopus*. These are summarized on table 5.

Comment [EM28]: You need to described each results separate under separate headings as:
3.1.1: for Table 1
3.1.2: for Table 2
3.2.3: for Table 3
3.2.4: for Table 4

Comment [EM29]: Delete the citation is not used when describing the results. You can only use it when discussing the results of the study.

Comment [EM30]: Done mention a Table twice e.g. Table 5

Table 1: Microbial counts of fish samples from three various ponds and Control

Location	Part isolated	Coliform count	Standard
Organization of			
Of Ponds	from	(cfu/ml)	Nigeria, AL(4×10^2 cfu/ml)
Bosso pond	Skin	3.8×10^2	Satisfactory
	Gills	3.0×10^1	Satisfactory
Kure pond	Skin	2.4×10^2	Satisfactory
	Gills	2.6×10^2	Satisfactory
Mobil pond	Skin	4.6×10^3	Marginal
	Gills	3.2×10^1	Satisfactory

Comment [EM31]: What is this???

Comment [EM32]: This table is scattered. You need to rearrange it

Data are means of triplicate determinations cfu/ml = colony forming unit per meal; AL = acceptable level, Satisfactory $< 10^3$; Marginal $< 10^5$; Unsatisfactory $> 10^5$

UNDER PEER REVIEW

Table 2: Microbial count of fresh catfish (*Clarias gariepinus*) obtained from three different locations.

S/No.	Location Of market	Part of fish (cfu/ml)	Coliform count (cfu/ml)	Fungi Count(SON), AL (4×10^2 cfu/ml)	
1.	Bosso Market	Skin	5.6×10^5	1.3×10^2	Marginal
2.		Gills	4.0×10^5	0.5×10^1	Marginal
3.	Kure Market	Skin	1.64×10^3	2.0×10^3	Unsatisfactory
4.		Gills	7.2×10^4	1.0×10^2	Unsatisfactory
5.	Mobil Market	Skin	6.0×10^6	0.7×10^4	Unsatisfactory
6.		Gills	4.8×10^4	0.7×10^2	Marginal

Key; Mkt = market. Cfu/ml = colony forming unit per meal

Table 3: CHARACTERIZATION AND IDENTIFICATION OF BACTERIA ISOLATED

Comment [EM33]: Present the table in landscape. Also, use insert to insert the table.

MCA	Cells	colonial morphology		Biochemical test carried							
		of market	from	shape	Elevation	form	Gram	Motility	MR	VP	
Indole	Citrate	Catalase	Coagulase	micro-organism	Reaction	test	test	test	test	test	test
BOSMKT _{MN}	Skin	Pink	Rods	Convex circular	-	-	+	-			
+	-	+	-	Bacilli spp							
	Gills	White	Rods	Convex Irregular	+	+	+	-	+		
+	+	+									
KURMKT _{MN}	Skin	Pink	Cocci	Convex spindle	+	+	-	+	+	-	
+	+	Staphylococcus									
	Gills	Pink	Rods	Convex spindle	+	+	+	-	+		
+	+	+									
MOBMKT _{MN}	Skin	Pink	Cocci	Convex circular	-	-	-	+	-		
-	-	-									
	Gills	White	Rods	Convex circular	+	+	+	-	+		
+	+	+									
BOSPOND	Skin	Green	Rods	Spherical	-	+	+	-	-		
+	-	-									
	Gills	Pink	Cocci	Convex spindle	+	-	-	-	+		
-	-	+	+								
KURPOND	Skin	Green	Cocci	Convex spindle	+	-	-	+	-	+	
+	+										
	Gills	Pink	Rods	Convex spindle	+	+	-	-	+		
-	-	+	+								
MOBPOND	Skin	Pink	Rods	Convex spindle	+	-	-	-	+		
-	-	+	-								

- Gills Pink Rods Convex spindle + - - +
- - + +

Key; BOSMKT_{MN}__ Bosso market Minna Niger State, KURMKT_{MN}__ Kure market Minna Niger State, MOB_{MN}MKT__ Mobil market Minna Niger State, MCA__Microscopic appearance

UNDER PEER REVIEW

Table 4: Summary of the entire microorganism identified from the skin and gills of catfish

Comment [EM34]: Please redesign the table using insert icon. Also, names of the microorganisms should be written correctly.

Location	Part of the fish	Bacterial	Fungi
Bosso market	Skin	<i>Escherichia coli</i>	-
	Gills	<i>Staphylococcus spp</i>	<i>Aspergillus niger</i>
	Skin	<i>Streptococcus spp</i>	<i>Rhizopus spp</i>
	Gills	-	
Kure market	Skin	<i>Pseudomonas aeniginosa</i>	
	Gills	<i>Escherichia coli</i>	
	<i>Aspergillus niger</i>		
	Gills	<i>Staphylococcus spp</i>	<i>Aspergillus niger</i>
Mobil market	Skin	<i>Escherichia coli</i>	<i>Rhizopus spp</i>
	Gills	<i>Staphylococcus aureus</i>	
	<i>Aspergillus flavus</i>		
Bosso pond	Gills	<i>Enterobacterspp</i>	
	Skin	<i>Pseudomonas spp</i>	-
Mobil pond	Skin	<i>Streptococcus spp</i>	<i>Rhizopus spp</i>
	Gills	<i>Shigella spp</i>	
Kure pond	Skin	<i>Escherichia coli</i>	-
	Gills	<i>Staphylococcus spp</i>	<i>Aspergillus niger</i>

Table 5: Morphological Characteristics and Identification of Mould Isolated in Catfish

Part of the fish	Colour of Characteristics	Colour of Probable	Nature of	Shape and kind	Presence of	Appearance
Sporangiophore	Aerial Hyphae of Sporehead	Substrate Organism	Hyphae	of Asexual	Special	
Haphae	Spore	Structure	or Conidiophores			
Gills	Whitish black	Brown	Septatate	Oval greenish	foot cell	long erect
Swell into	<i>Aspergillus</i>					
		Conidia nucleate	<i>spp</i>	Present	Septate conidia	Multi-
Skin conidia	Dark green	Purplish red	Septed	Brown	Absent	Micro-
	Short-columer <i>Rhizopore</i>					

Comment [EM35]: Check your table. It is generally not clear, and does not explain any result.

Discussion

Catfish contamination at various markets

The result shown in Table 3 reveals that catfish species purchased from the various markets in Minna metropolis were highly contaminated with various bacterial species such as *Escherichia coli*, *Staphylococcus spp*, *Streptococcus spp*, *Bacillus*, *Shigella*, *Enterobacterspp* and *Pseudomonas* and fungi species were *Aspergillus spp* and *Rhizopus*. This could have resulted due to the dirty environment of the markets or from the fish seller/ handlers themselves through their containers, sneezing and picking of noses. This is in consistence with the observation of Daniel *et al*, (2006) who reported higher concentration of bacterial at various markets at Makurdi as compared to fresh fish samples caught at River Benue. However the total coliform count was highest on the skin than the gills from catfish obtained from Bosso market compared to that of Mobil and Kure market. This could be due to the concentration of broken soak ways and poor sewage disposal systems available there.

Comment [EM36]: Your discussion should cover all the results obtained in the study. In other word, all the results of the study must be discuss, and compare with the findings of other people.

Comment [EM37]: Heading is not required here. Hence, it should be deleted.

Comment [EM38]: Avoid repetition of your result by referring to a particular table or figure. You need to compare your result with the results of at least two researchers in the same or related topic.

Mould and Fungi contamination at various markets

Comment [EM39]: Delete

Aspergillus niger and *Rhizopus* spp probably contaminated the catfish sample through the air and soil, micro-organisms are ubiquitous, (Adam and Moss, 2000). These were not isolated from the central expert i.e. the catfish obtained from ponds. Adequate and proper cleaning detergents used by the fish reared may have ensured that there was no fungal nor mould contamination. *Aspergillus* spp and *Rhizopus* spp contaminated catfish samples from Bosso, Mobil and Kure markets. This could be due to inadequate personal hygiene.

CONCLUSION

The microbial load of the catfish (*Clarias gariepinus*) varies according to difference in location and distance from purchasing source. This has shown clearly from the data generated, the following species of organisms are the major contaminants, such species are *Escherichia coli*, *Staphylococcus* spp, *Streptococcus* spp, *Bacillus*, *Shigella*, *Enterobacter* spp and. This contamination of fish occurs probably due to conveyance of fish to the various markets or the material in which the fish is being kept on, it may also be as a result of exposure of fish to flies and insects. This contamination may also result from dusty atmosphere and dirty environment, by heaps of refuse dump all over the place by the market seller.

Comment [EM40]: ????

Comment [EM41]: Conclude your study by writing based on the finding of the study.

Recommendation

The following recommendation will help to foster fish farming and deterioration of fish and its products. The proper sanitation should be implemented in the fresh fish market in order to maintain the hygienic environment. Awareness campaigns need to be conducted at fish selling locations on mode of fish handling to reduce contamination. Further studies need to be carried out in order to ascertain the point of entry of contaminants with a view of curtailing it.

REFERENCES

- Abe, H. and E. Okuma (1991). Rigor mortis progress of carp acclimated to different water temperatures, *Nippon Suisan Gakkaishi*, 57, 2095-2100.
- Adams, M.R. and Moss, M.O. (1995). *Food microbiology*, Royal Society of Chemistry, Science Park, Cambridge pp 17, 121-122.
- Alverson, D.L., M.H. Freeberg, J.G. Pope, S.A. Murawski (1994). A global assessment of fisheries by-catch and discards. FAO Fish. Tech. Pap. No. 339. FAO, Rome.
- Anthoni, U., T. Borresen, C. Christophersen, L. Gram, and P.H. Nielsen (1990). Is trimethylamine oxide a reliable indicator for the marine origin of fishes. *Comp. Biochem. Physiol.* 97B, 569-571.
- Anthonio, O.R. and Akinwumi, J.A.: Supply and distribution of fish in Ibadan, Nigeria. *Geog. J.*

- 14(2): 16 (1991).
- Assiah, Van Eer, O.: Small Scale Fresh Water Fish Farming. Agromisa Press, Netherlands, Pp 1-3 (1997). F.A.O.: Fish for Food and Employment. Food and Agricultural Organization, Rome (1991).
- Azam, K., I.M. Mackie and J. Smith (1990). Effect of stunning methods on the time of onset, duration and resolution of rigor in rainbow trout (*Salmo gairdneri*) as measured by visual observation and analysis for lactic acid, nucleotide-degradation products and glycogen. In: Chilling and freezing of new fish products. Sci. Tech. Froid. 1990-3. Proceedings of the meeting of Commission C2 I.I.F.-I.I.R. Aberdeen. 351-358.
- Benso, H.J. (1990). Microbiological Application: A Laboratory Manual in general Microbiology, 5th ed. Win C. Brown Publishers, USA. pp. 288.
- Borresen, T. (1992). Quality aspects of wild and reared fish. In: H.H. Huss, M. Jacobsen and J. Liston (eds.) Quality Assurance in the Fish Industry. Proceedings of an International Conference, Copenhagen, Denmark, August 1991. Elsevier, Amsterdam, 1-17.
- Botta, J.R., K.M. Kennedy, J.W. Kiceniuk, and J. Legrow, (1992). Importance of redfish level, fish size and roe content to the quality of roe capelin. Int. J. Food Sci. Technol. 27, 93-98.
- Boyd, L.C., D.P. Green, F.B. Giesbrecht, and M.F. King (1993). Inhibition of oxidative rancidity in frozen cooked fish flakes by tert-butylhydroquinone and rosemary extract. J. Sci. Food Agric. 61, 87-93.
- Bremner, H. A. (1992). Fish flesh structure and the role of collagen - its postmortem aspects and implications for fish processing. In: H.H. Huss, M. Jacobsen and J. Liston (eds.) Quality Assurance in the Fish Industry. Proceedings of an International Conference, Copenhagen, Denmark, August 1991. Elsevier, Amsterdam, 39-62.
- Chiba, A., M. Hamaguchi, M. Kosaka, T. Tokuno, T. Asai, and S. Chichibu (1991). Quality evaluation of fish meat by ³¹P-phosphorus-nuclear magnetic resonance. J. Food Sci. 56, 660-664.
- Clucas I.J. (1991). Design and trials of ice boxes for use on fishing boats in Kakinada, India. Bay of Bengal Programme BOBP/WP67. Madras, India.
- Dalgaard, P. (1993). Evaluation and prediction of microbial fish spoilage. Ph.D. Thesis. The Technological Laboratory of the Danish Ministry of Fisheries and the Royal Veterinary and Agricultural University, Denmark.
- Dalgaard, P. (1994). Qualitative and quantitative characterization of spoilage bacteria from packed fish. Int. J. Food Microbiol. (In press).
- Dalgaard, P. (1994). Modeling of microbial activity and prediction of shelf life for packed fresh fish. Int. J. Food Microbiol. (In press).

- Dalgaard, P., L. Gram, and H.H. Huss (1993). Spoilage and shelf life of cod fillets packed in vacuum or modified atmospheres. *Int. J. Food Microbiol.* 19, 283-294.
- Dalgaard, P. and H.H. Huss (1994). Mathematical modelling used for evaluation and prediction Of microbial fish spoilage. In: D.E.Kramer, F. Shahidi and Y. Jones (eds.) *Proceedings of the Symposium New Developments in Seafood Science and Technology*, CIFST, Vancouver, Canada.
- DANIDA (1989). *Environmental Issues in Fisheries Development*. DANIDA, Danish Ministry of Foreign Affairs, Copenhagen.
- Devaraju, A.N. and T.M.R. Setty (1985). Comparative study of fish bacteria from tropical and cold/temperate marine waters. In: Reilly, A. (ed.) *Spoilage of tropical fish and product development*. FAO Fish. Rep. (317) Suppl., 97-107.
- DiChristina, T.J. and E. F. DeLong (1993). Design and application of rRNA-targeted oligonucleotide probes for the dissimilatory iron- and manganese-reducing bacterium *Shewanella putrefaciens*. *Appl. Environ. Microbiol.* 59, 4152-4160.
- DiChristina, T.J. and E. F. DeLong (1994). Isolation of anaerobic respiratory mutants of *Shewanella putrefaciens* and genetic analysis of mutants deficient in anaerobic growth on Fe²⁺. *J. Bacteriol.* 176, 1464-1474.
- Etienne, M. and N. Bregeon (1992). A quick enzymatic quantitative analysis of histamine in tuna by microplate reader. Proc. of the 22nd annual meeting of the Western European Fish Technology Association.
- F.A.O.: F.A.O.'s Newsletter (2000). Anthonio, O.R. and Akinwumi, J.A.: Supply and distribution of fish in Ibadan, Nigeria. *Geog. J.*, 14(2): 16 (1991).
- FAO (1993a), *FAO Yearbook: Fishery Stat.* Vol. 72 and 73. FAO. Rome.
- FAO (1993b). *The State of Food and Agriculture 1993*. FAO Agric. Ser. 26.
- FAO (1993c). *Aquaculture production 1985-1991*. FAO Fisheries circular No. 815, Rev. 5.
- FAO (1994). *Review of the state of world marine fishery resources*. FAO Fish. Tech. Pap. 335.
- Farber, J. M. (1991). Microbiological aspects of modified-atmosphere packaging technology. A review. *J. Food Prot.* 54, 58-70.
- Ferich, G.N. and Hendrie, M.S. (1985). Bacteria associated with diseases of fish. In: *Isolation and Identification of Microorganism of medical and Veterinary Importance*.
- Freeman D.W. and J.O. Heamsberger (1993). An instrumental method for determining Rancidity in frozen catfish fillets. *J. Aquat. Food Prod. Technol.* 2, 35-50.
- Gill, T.A. (1990). Objective analysis of seafood quality. *Food Rev. Int.* 6, 681-714.

- Gill, T.A. (1992). Biochemical and chemical indices of seafood quality. In: H.H. Huss, M. Jacobsen and J. Liston (eds.) *Quality Assurance in the Fish Industry. Proceedings of an International Conference, Copenhagen, Denmark, August 1991.* Elsevier, Amsterdam, 377-388.
- Gill, T.A., J. Conway, and J. Evrovski (1992). Changes in fish muscle proteins at high and low temperature. In: G.J. Flick and R.E.Martin (eds.) *Advances in seafood biochemistry-composition and quality,* Technomic Publishing, Lancaster, Pennsylvania, 213-231.
- Gram, L. (1992). Evaluation of the bacteriological quality of seafood, *J. Food Microbiol.* 16, 25-39.
- Haard, N.F. (1992). Technological aspects of extending shelf life of seafood: A review. *J.Aqua. Food Prod. Technol.* 1, 9- 27.
- Ihekoronye, A.I. and Ngoddy, P.O. (1985). *Integrated Food science and Technology for the Tropics.* Macmillan Publishers, London pp 333-342.